Victoria University of Wellington

School of Biological Sciences

A thesis submitted to the Victoria University of Wellington in fulfilment of the
requirements for the degree of Doctor of Philosophy

Big fish, little fish:
a multi-omics investigation of growth traits in silver trevally
(*Pseudocaranx georgianus* – araara)

Doctoral Dissertation of:
Noemie Valenza-Troubat

Primary Supervisor:

Dr. Peter Ritchie

Secondary Supervisor:

Dr. Maren Wellenreuther

April 2021

“No one supposes that all the individuals of the same species are cast in the same actual mould. These individual differences are of the highest importance for us, for they are often inherited.”

Charles Darwin,
On the Origin of Species by Means of Natural Selection,
or the Preservation of Favoured Races in the Struggle for Life, 1859

Life will always be more work. The only thing you can do is make it more fun.

Abstract

Understanding the relationship between DNA sequence variation and the diversity of observable traits across the tree of life is a central research theme in biology. In all organisms, most traits vary continuously between individuals. Explaining the genetic basis of this quantitative variation requires disentangling genetic from non-genetic factors, as well as their interactions. The identification of causal genetic variants yields fundamental insights into how evolution creates diversity across the tree of life. Ultimately, this information can be used for medical, environmental and agricultural applications. Aquaculture is an industry that is experiencing significant global growth and is benefiting from the advances of genomic research. Genomic information helps to improve complex commercial phenotypes such as growth traits, which are easily quantified visually, but influenced by polygenes and multiple environmental factors, such as temperature. In the context of a global food crisis and environmental change, there is an urgent need not only to understand which genetic variants are potential candidates for selection gains, but also how the architecture of these traits are composed (e.g. monogenes, polygenes) and how they are influenced by and interact with the environment. The overall goal of this thesis research was to generate a genome-wide multi-omics dataset matched with exhaustive phenotypic information derived from a F₀-F₁ pedigree to investigate the quantitative genetic basis of growth in the New Zealand silver trevally (*Pseudocaranx georgianus*). These data were used to identify genomic regions that co-segregate with growth traits, and to describe the regulation of the genes involved in response to temperature fluctuations. The findings of this research helped gain fundamental insights into the genotype–phenotype map in an important teleost species and understand its ability to dynamically respond to temperature variations. This will ultimately support the establishment of a genomics-informed New Zealand aquaculture breeding programme.

Chapter 1 of this thesis provides an overview of how genes interact with the environment to produce various growth phenotypes and how an understanding of this is important in aquaculture. This first chapter provides the deeper context for the research in subsequent data chapters.

Chapter 2 describes the study population, the collection of phenotypic and genotypic data, and a first description of the genetic parameters of growth traits in trevally. A combination of Whole Genome Sequencing (WGS) and Genotyping-By-Sequencing (GBS) techniques

were used to generate 60 thousand Single Nucleotide Polymorphism (SNP) markers for individuals in a two-generation pedigree. Together with phenotypic data, the genotyping data were used to reconstruct the pedigree, measure inbreeding levels, and estimate heritability for 10 growth traits. Parents were identified for 63% of the offspring and successful pedigree reconstruction indicated highly uneven contributions of each parent, and between the sexes, to the subsequent generation. The average inbreeding levels did not change between generations, but were significantly different between families. Growth patterns were found to be similar to that of other carangids and subject to seasonal variations. Heritability as well as genetic and phenotypic correlations were estimated using both a pedigree and a genomic relatedness matrix. All growth trait heritability estimates and correlations were found to be consistently high and positively correlated to each other.

In **Chapter 3**, genotypic and phenotypic data were used to carry out linkage mapping and a genome-wide association study (GWAS) to map quantitative trait loci (QTLs) associated with growth differences in the F₁ population. A linkage map was generated using the largest family, which allowed to scan for rare variants associated with the traits. The linkage map reported in this thesis is the first one for the *Pseudocaranx* genus and one of the densest for the carangid family. It included 19,861 SNPs contained in 24 linkage groups, which correspond to the 24 trevally chromosomes. Eight significant QTLs associated with height, length and weight were discovered on three linkage groups. Using GWAS, 113 SNPs associated with nine traits were identified and 29 genetic growth hot spots were uncovered. Two of the GWAS markers co-located with the QTLs discovered with the linkage mapping analysis. This demonstrates that combining QTL mapping and GWAS represents a powerful approach for the identification and validation of loci controlling complex phenotypes, such as growth, and provides important insights into the genetic architecture of these traits.

Chapter 4, the last data chapter, investigates plasticity in gene expression patterns and growth of juvenile trevally, in response to different temperatures. Temperature conditions were experimentally manipulated for 1 month to mimic seasonal extremes. Phenotypic differences in growth were measured in 400 individuals, and the gene expression patterns of the pituitary gland and the liver were compared across treatments in a subset of 100 individuals, using RNA sequencing. Results showed that growth increased 50% more in the warmer compared with the colder condition, suggesting that temperature has a large impact on the metabolic activity associated with growth. We were able to annotate 27,887 gene models and found 39 differentially expressed genes (DEGs) in the pituitary, and 238 in the liver. Of these, 6 DEGs

showed a common expression pattern between the tissues. Annotated blast matches of all DEGs revealed genes linked to major pathways affecting metabolism and reproduction. Our results indicate that native New Zealand trevally exhibit predictable plastic regulatory responses to temperature stress and the genes identified provide excellent for selective breeding objectives and studied how populations may adapt to increasing temperatures.

Finally, **Chapter 5** discusses the implications, future directions, and application of this research for trevally and other breeding programmes. It more broadly highlights the insights that were gained on the genetic architecture of growth, and the role of temperature in interacting and modulating genes involved in plastic growth responses.

Acknowledgements

Undertaking this PhD research has been the best decision of my life, so far. It involved moving to Nelson, New Zealand, learning three different coding languages, diving deep into the whole new field that quantitative genetics was for me at the time and managing to write a thesis in another language. It also involved making this place home, creating a new family, discovering the wonders of this country, and spending my spare time tramping, sailing, climbing, skiing, and laughing. Over anything else, these 3.5 years have taught me how much community and support are essential. This is team work. On both a professional and personal level. This section is not so much an acknowledgment, but merely the expression of my deep gratitude for being surrounded by extraordinary people, without whom none of this would have been possible. Or might have, but in a very painful way. Most of you are not just colleagues anymore, but friends. And not just friends anymore, but family.

To Maren. Thank you for believing I was able to pull this work together. During the numerous moments of doubts I've had since starting, the fact that you chose me as a student, that you thought I could do this, has always helped me. Thank you for being a role model to me and to women in science. During my education, most of my teachers have been males and I realise that I was missing someone to look up to. Thank you for showing me that women can be high achievers, extremely successful scientists, but also friends, mothers and fit and fun. Thank you for creating our team, for being an amazing leader and also much more than a supervisor.

To Peter. Thank you for always having an encouraging smile, a sympathetic ear, and generally great life advice. Our philosophical chats and you sharing your experiences have helped me navigate tough decisions and keep a clear mind as to what I want.

To my science big brothers. Peter, you have been a confidant, a shoulder, and a guide. Thank you for being so sensitive and knowing how to listen when I needed to talk. Thank you for sharing your experiences, your advice and your help. I am humbled by your wisdom. You also have sent me countless dubious memes and laughed at many of my questionable jokes, so brownie points for that. Matt, thank you for never being tired of me, your “annoying little sister”, always answering my endless questions about your life during your thesis, listening to my rants about veganism, and work and life in general. Thank you for eating lunch disturbingly

early with me, for always being up for a beer (but not at the Freehouse) and bearing me as a flatmate.

To Marcus. Thank you for showing me (some of) the many ropes you know. Although it annoyed me at first, the point you made about teaching me from the beginning rigorously reproducible workflows, change control, simulations or basic coding etiquette have really shaped my way of working and I couldn't be more grateful. Thank you for killing the thousands of jobs I would send simultaneously to the servers by mistake, keeping your cool while I bombarded you with millions of questions about statistics and algorithms, sharing yummy vegan curry recipes and more broadly having been the best coding mentor I could ask for.

To Elena. Mil gracias por no haber tenido miedo frente al desafío que representaba mi proyecto... You taught me that rigorous work is also synonym of laughter and kindness. And thank you for sending me cute cat videos or funny memes to cheer me up, those really helped!

To Sara, for being a brilliant scientist, colleague, ski buddy and friend. Working with you has been a pleasure and I hope this is just the start of a long collaboration.

To my science little brothers at VUW, Tom and Yvàn. I loved every visit, rant, video call and cheeky Friday drinks with you two. By the way, you two shouldn't work so late.

To my amazing APG team: Igor, David, Dafni, Christina, Erin, Nick. And the rest of PFR staff, Alex, Bodhi, Roy, Therese, Warren, Scottie, Steve, Marc, Josh, David chocolatine.

To my Nelson family. My heart sisters: Christie, Victoria, Alison, Camille, Marama, Gemma, Karen. My wāhine. My confidants, supporters, adventurers. Sisterhood takes all its sense with you. Being known and vulnerable is amazing! To Ben and Benjah. And to the rest of my many friends here.

A ma famille. Loin physiquement mais à jamais plus près dans mon cœur. A Maman, parce que tu as créé entre nous un lien qui dépasse mère-fille. A Papa, et ta curiosité contagieuse pour les sciences et tes questions dans la voiture. Sans toi je ne serais pas ici aujourd'hui. A Lara, pour ton soutien indéfectible. A Lucas et Paul, qui me manquez chaque jour. Aux deux meilleures grand-mères du monde, Nanique et Calie.

A Julien, je sais que de là où tu es, tu es fier de moi.

From the bottom of my heart, thank you.

External contributions

This research was funded by the programme “Accelerated breeding for enhanced seafood production” (C11X1603), funded by the New Zealand Ministry of Business, Innovation, and Employment (MBIE) Endeavour Fund. All fish used in this study were supplied by the New Zealand Institute for Plant & Food Research Limited. All laboratory work, analysis and writing were conducted by the author with the help of supervisors Drs Peter Ritchie and Maren Wellenreuther. Additional contributors are mentioned below:

- The Genotyping-by-Sequencing and RNA-sequencing methods applied in this thesis were supervised by Dr. Elena Hilario from Plant & Food Research in Auckland.
- The trevally genome assembly used was supplied by Roy Storey and Marcus Davy at Plant & Food Research in Te Puke, and is now published in Ruigrok et al. (2021).
- The heritability models were reviewed and improved by Dr Fiona Hely at the agribusiness consulting company ABACUSBIO.
- Maintenance of trevally fish populations for this thesis was largely carried out by staff at Plant & Food Research in Nelson.

Contents

1	General Introduction.....	1
1.1	A brief history of genetics.....	1
1.2	Research questions	4
1.3	Genetic influence on traits.....	6
1.3.1	Heritability and additive genetic variance	7
1.3.2	Gene pleiotropy: the Swiss army genes.....	8
1.3.3	Epistasis and other context-dependent effects.....	9
1.3.4	QTL mapping	10
1.3.5	Genome-wide association studies.....	11
1.4	Environmental influence on traits	12
1.4.1	Phenotypic plasticity.....	13
1.4.2	Gene expression, the intermediate phenotype	15
1.4.3	RNA-sequencing and statistical frameworks	15
1.5	Growth is a complex trait	17
1.5.1	Endocrine regulation of somatic growth	17
1.5.2	The impact of temperature.....	19
1.5.3	Genomic insights to mitigate the effects of temperature variations	21
1.6	Wider relevance of this research	22
1.6.1	Aquaculture under climate change	22
1.6.2	The role of aquaculture in New Zealand	23
1.6.3	Trevally (<i>Pseudocaranx georgianus</i>) – araara	24
1.7	Aims and thesis structure	25
1.7.1	Specific objectives	26
1.7.2	Thesis structure.....	26
2	Evaluating new species for aquaculture: a genomic dissection of growth in New Zealand silver trevally (<i>Pseudocaranx georgianus</i>).....	29
2.1	Abstract	30
2.2	Introduction	30
2.3	Methods.....	33
2.3.1	Populations studied and holding conditions	33

2.3.2	Tissue sampling, DNA extraction and library preparations	34
2.3.3	Genotyping data quality checking and processing	35
2.3.4	Pedigree reconstruction and F ₀ sex prediction	36
2.3.5	Inbreeding calculations	37
2.3.6	Phenotyping, trait estimations and phenotypic correlations	37
2.3.7	Trait heritability and genetic correlations	39
2.3.8	Ethics statement	40
2.4	Results	40
2.4.1	Filtering allowed to increase quality of calls tenfold in offspring	40
2.4.2	Reconstruction of the pedigree showed skewed parental contributions	41
2.4.3	Unbalanced inbreeding levels detected within families	43
2.4.4	Growth pattern in trevally is seasonally influenced and is under family-effects	43
2.4.5	Genetic correlations, phenotypic correlation and trait heritability	46
2.5	Discussion	48
2.5.1	Future directions and management implications	50
3	Unravelling the complex genetic basis of growth in New Zealand silver trevally (<i>Pseudocaranx georgianus</i>)	53
3.1	Abstract	54
3.2	Introduction	54
3.3	Methods	56
3.3.1	Study population	56
3.3.2	Phenotyping and trait estimation	57
3.3.3	Genotyping and variant calling	57
3.3.4	Linkage map construction and QTL identification	58
3.3.5	Genome-Wide Association Study	58
3.3.6	Ethics statement	59
3.4	Results	59
3.4.1	Phenotypic values of growth traits	59
3.4.2	QTL mapping revealed rare variants associated with growth	61
3.4.3	Genome-wide association found strongly associated common SNPs	66
3.5	Discussion	67
3.5.1	Future directions	72

4	Differential expression analyses reveal extensive transcriptional plasticity induced by temperature in New Zealand silver trevally (<i>Pseudocaranx georgianus</i>).....	75
4.1	Abstract	76
4.2	Introduction	77
4.3	Methods.....	79
4.3.1	Fish holding procedures, experimental setup and trait phenotyping	79
4.3.2	RNA extraction, Illumina library preparation and sequencing.....	80
4.3.3	Genome annotation and completion rate	80
4.3.4	Sequencing data processing	81
4.3.5	Differential expression analysis.....	82
4.3.6	Gene ontology.....	82
4.3.7	Ethics statement	83
4.4	Results	83
4.4.1	Gain in weight increased in the high treatment	83
4.4.2	Raw sequencing data and quality statistics.....	83
4.4.3	Functional annotation and filtering.....	85
4.4.4	Differential gene expression analysis revealed extensive transcriptional plasticity	86
4.4.5	Gene ontology to reveal molecular pathways.....	89
4.5	Discussion	91
4.5.1	Future directions	94
5	General Discussion	97
5.1	Overview	97
5.2	Summary of key findings	98
5.2.1	Data quality is of utmost importance.....	98
5.2.2	Reconstruction of the study pedigree and evaluating inbreeding level	99
5.2.3	Trait heritability is a key determinant in selective breeding programmes	100
5.2.4	Linkage maps and QTL mapping allow to understand the genetic architecture of traits.....	101
5.2.5	Applying GWAS analyses to fine map QTLs	103
5.2.6	Growth is strongly influenced by temperature	105
5.3	Future research and recommendations	107
5.3.1	The complexity of decision making and resulting biases.....	107

5.3.2	Aquaculture breeding programmes	108
5.3.3	Enhancing our understanding of the genetic architecture of growth.....	110
5.3.4	Climate change and implications for trevally	112
5.4	Conclusions	115
A	Supplementary material.....	117
A.2	Chapter 2	118
A.3	Chapter 3	131
A.4	Chapter 4	135
B	Appendix 1: Iteration process for QTL mapping.....	180
	Bibliography.....	184

Glossary

Additive genetic variation: The variance in a trait that is due to the effects of each individual allele being added together, without any interactions with other alleles or genes.

Allele: One of two or more forms of a given gene variant.

BUSCO: Benchmark Universal Single-Copy Orthologue. An assessment tool which provides quantitative measures of genomic data completeness in terms of expected gene content. Used to gauge the quality and completeness of genome assemblies and their annotations.

Contig: Set of overlapping DNA fragments that together represent a consensus region of the DNA.

Continuous (or quantitative) trait: Distribution of phenotypes varying along a continuum. Individuals differ by small degrees (examples: height, blood pressure, reaction time, learning ability).

Coverage: Read depth per locus, per individual.

DEGs: Differentially expressed genes. Comparison of abundance of gene transcripts between sample group types or within a transcriptome.

Discrete (or qualitative) trait: Traits occurring in distinct categories: e.g. presence or absence (examples: albinism, cystic fibrosis, Huntington's disease), or classes (colour, sex).

Mendelian traits	Complex traits
Discrete traits	Continuous traits
Single or few major genes	Polygenically inherited
Genes have a major effect on phenotype	Each gene has a minor effect on phenotype
Location and function often known	Location and effect often unknown
Environmental effects are limited	High influence of the environment

Dominant genetic variance: The phenotypic variance contributed from dominant gene actions that covers the influence of recessive alleles at particular loci.

Epistasis: The interaction between genes at two or more loci. The expression of one gene modifies the phenotype of another gene.

GBS: Genotyping-by-Sequencing. Uses restriction enzymes to reduce genome complexity and allows to genotype multiple DNA samples at once. This technique targets only a small percentage of the genome which allows multiplexing (the sequencing of multiple DNA samples at once). This technique does not require previous knowledge of variation.

Gene: Specific fragment of DNA that codes for a protein.

Genetics: The study of specific and limited numbers of genes, or parts of genes, that have a known function.

Genetic architecture: A description of all the genetic and environmental factors that influence a trait.

Genome: All of the genetic material of an organism. Includes both coding and non-coding regions.

Genomics: The study of the entirety of the genome, including genes, regulatory elements, transposable elements, other repeats.

Genotype: An individual's collection of genes. In this thesis however, genotype is used to refer to the two alleles inherited for a particular gene.

Linkage group: Set of genes known to be grouped and inherited together (except for crossing-over). Eventually the group covers an entire chromosome.

Locus (loci): Region(s) of the genome. Can be a gene or a molecular marker.

N50: The sequence length of the shortest contig at 50% of the total genome length.

Non-additive variance: The variance associated with epistasis and dominance. The expression of an allele is conditional on the expression of another allele.

Phenotype: Observable trait or characteristic.

QTL: Quantitative Trait Loci. Genes controlling quantitative traits. One trait may be controlled by multiple QTLs, each of which segregate according to Mendelian laws.

Scaffold: A portion of the genome sequence reconstructed from end-sequenced whole-genome shotgun clones. Scaffolds are composed of contigs and gaps.

Variance: Measure of variability, partitioned into components attributable to different sources of variation. It is calculated as the expectation of the squared deviation of a random variable from its mean.

Variant: A specific region of the genome which differs between two genomes.

WGS: Whole Genome Sequencing. Performed on massively parallel sequencing technologies in order to retrieve enough DNA fragments to cover the whole span of the genome of interest

1 General Introduction

1.1 A brief history of genetics

Since prehistoric times, humans have noticed that offspring are similar to their parents and used this observation to improve crops and animals through selective breeding. Genetics, as the modern field of science seeking to understand this process, really began in the mid-19th century with the publication of two major pieces of work. In 1858, Charles Darwin and Alfred Wallace formulated simultaneously a ground-breaking, yet controversial theory: evolution driven by natural selection (Darwin, 1858; Darwin et al., 1858). By explaining that species evolve over time, that new species come from pre-existing ones, and that all species share a common ancestor, they were shedding light on the mechanisms behind the diversity of all living beings and how the environment drives selection. The evidence they used to support their theory convinced an increasing number of fellow scientists that species indeed evolved by descent with modification from their ancestors. However, the processes driving these changes continued to be fiercely debated,

and natural selection as a causal driver only received widespread approval much later by the scientific community. Shortly after the formulation of this foundational theory, Gregor Mendel published his work on the transmission of qualitative (or discrete) traits in peas through heritable units, now known as genes (Mendel, 1866). Using field experiments and a statistical framework, he developed fundamental laws for how genes govern the inheritance of traits. However, after Mendel's discovery, researchers were finding that qualitative traits were rare, and that most of the heritable units were actually quantitative (or continuous). In 1918, Ronald Fisher published "The Correlation Between Relatives on the Supposition of Mendelian Inheritance", where he introduced the term variance and explained the idea that continuous variation amongst phenotypic traits could be the result of the combination of many discrete genes of weak effects, and thus be the result of Mendelian inheritance. This was the first step towards establishing quantitative genetics, or the study of continuous phenotypes. In 1930, he published a second seminal piece of work, "The Genetical Theory of Natural Selection", mathematically unifying Mendelian genetics and evolutionary selection by demonstrating that the latter could change allele frequencies in a population. Pushed to its extreme, this type of model is referred to as the 'infinitesimal model', and it laid the foundations for the growth of quantitative genetics in the 20th century (Lynch & Walsh, 1999). In the 1940s, a fundamental effort took place across the scientific community to weave together the fields of genetics, developmental physiology, ecology, cytology and mathematics together into a framework that was termed the 'Modern Evolutionary Synthesis'. This framework provided a holistic unification of several, until then, standalone concepts. It allowed scientists to understand interactions between the genetic makeup of an individual and the environment, and how this impacts trait evolution in a more integrated way. By disentangling how evolutionary selection shapes individual phenotypes, populations and species, scientists started to gain a deeper understanding about how microevolution connects to macroevolution.

Although Fisher's work retained its primordial influence, the importance of the infinitesimal model remained a theoretical abstraction for most of the 20th century. For many, only a few loci (< 10) could approximate infinitesimal predictions (McGuffin & Huckle, 1990; Thoday & Thompson, 1976) and it was entirely unclear how many genetic regions actually affected complex traits (Risch et al., 1999; Visscher & Goddard, 2019). As an example, a review of quantitative genetics published in 1989 stated: "we still do not know whether the number of loci responsible for most genetic variation is small (5–20) or

large (100 or more)” (Barton & Turelli, 1989). In the 1990s, hundreds of studies on complex traits in human genetics, plant and animal breeding and evolutionary genetics were published, which were only powered to detect large-effect loci. This was the consequence of the lack of number of observation and technical limitations to screen the genome of the time. Typical examples include: foundational reviews on association mapping in human diseases, which computed the power to detect common variants with relative risks between the alternate homozygotes ranging from 2.25 to 16 (Risch & Merikangas, 1996); or breeding studies finding QTL with major effects on commercial traits (e.g. Coppieters et al., 1998; Nezer et al., 1999; Perretant et al., 2000). These genetic golden nuggets (with such dramatic effects as to behaving like Mendelian genes) led researchers to hope that large-effects alleles were the primary cause for evolution (Rockman, 2012).

The advent of the genomics era in the 21st century and the ability to generate thousands of genome-wide markers completely transformed our ability to study the genetic basis of complex traits (Claussnitzer et al., 2020; Nadeem et al., 2018). While studies showed the power of new technical and statistical approaches to identify significant and replicable signals, it quickly became clear that most variants generally explained only small fractions of the heritability of traits (Rockman, 2012). The limited power to detect causative loci became known as the “missing heritability” problem: a mystery that was largely resolved by work showing that most of the heritability is due to the presence of many sub-significant causal variants and their interactions (Zuk et al., 2012). Recent work is now showing that for most quantitative traits, the bulk of the heritability is spread widely and surprisingly uniformly across the genome (Yang et al., 2011; Loh et al., 2015; O’Connor, 2020). In fact, most complex traits are enormously polygenic, with various studies estimating > 10,000 or even > 100,000 causal variants per trait (O’Connor et al., 2019; Zhang et al., 2018).

As the number of loci found associated with traits increases, it is becoming more and more apparent that many genetic effects depend on the variation at other loci and that they are influenced by the environment (Bazakos et al., 2017; Mackay et al., 2009). Studies with a higher number of phenotypic measurements are associated with more alleles compared with studies with fewer measurements (Ashton et al., 2017). Alongside an increase in genotyping capacity, more precise technologies to define phenotypes and the incorporation of additional information about environmental conditions are revealing additional genetic interactions and enable more accurate estimation of effect sizes

(Robinson et al., 2014). The integration of multiple types of datasets and the ever growing computing power available are now allowing to develop and combine methods to study genotype-phenotype relationships. These results are used in many areas: disease risk prediction (Kamps et al., 2017), individual therapeutic treatments in human populations (Govender et al., 2020), adaptive evolution modelling (Waldvogel et al., 2020), and selective breeding programmes for economically important species (Tan et al., 2020).

1.2 Research questions

The overarching research questions addressed in this thesis were: what is the genetic basis of growth, what are the underlying associated variants, which molecular pathways are involved, and how are they influenced by the environment? To address these questions, this thesis used the teleost species silver trevally (*Pseudocaranx georgianus*) as a model and aimed to assess the heritability of various growth traits in a captive population, investigate genetic and phenotypic correlations, screen for QTLs associated with growth and assess the role of temperature and its effect on phenotypic plasticity and gene expression (Figure 1.1).

Figure 1.1: General thesis flow diagram.

1.3 Genetic influence on traits

Quantitative traits are complex phenotypes that do not behave like Mendelian traits: the genetic segregation of a single gene cannot explain their pattern of inheritance. Instead, these traits are the result of the interactions of many loci of small additive effects. Hence a fundamental, yet difficult, question to solve in quantitative genetics is: how much does each individual locus contribute to the total variance of a phenotype? As we will see, the relationship between genotype and phenotype is not straightforward and many mechanisms intervene along the way to give shape to organisms.

The fundamental goal of quantitative genetics (i.e. to investigate the loci underlying variation in heritable phenotypes, their effect size, architecture, mechanisms of action and interaction, and the influence of environmental variables) was summarised in the 1990s by a key equation in the field (Falconer & Mackay, 1996; Lynch & Walsh, 1999):

$$V_P = V_G + V_E$$

Here, the total phenotypic variance (V_P) is partitioned into the variance explained by all the genetic effects (V_G) and the environmental effects (V_E) influencing the trait. Under this view, all deviations from genotype values are considered as “environmental.” Over the years, following the discovery of epigenetic regulations and with a growing recognition of phenotypic plasticity, this equation was updated to:

$$V_P = V_G + V_E + V_{G \times E} + V_R \quad \text{Equation 1}$$

which accounts for noise (V_R) and also includes the explicit recognition of a genotype–environment interaction ($V_{G \times E}$) (Scheiner & Goodnight, 1984). This interaction specifies that the environmental effects are different for some genotypes relative to others. In other words, the trait value of any individual can be considered as the result of three distinct sets of factors: (1) the genetic factors that confer heritable differences within an environment, (2) the environmental factors that modulate the effects of the genotypes, and (3) the inherent variability of the trait given that genetic and environmental factors are identical.

The first member of *Equation 1*, the total genetic variance (V_G), can be further broken down into:

$$V_G = V_A + V_D + V_I$$

where V_A represents the additive genetic variance, V_D the genetic variance due to dominant effects and V_I the genetic variance explained by epistatic interactions. Each of these terms will be discussed more in depth in the next sections of this Introduction.

1.3.1 Heritability and additive genetic variance

The first set of factors of *Equation 1* refers to the total genetic variance. This category includes additive genetic effects as well as intra-genomic (non-additive) interactions (i.e. dominance, pleiotropy, epistasis). Additive genetic variance reflects the variance in a trait that is due to the effects of each individual allele being added together, without any interactions with other alleles or genes.

In the context of this thesis, we will consider two more equations. They both address the heritable components of a trait, which is of prime importance for this work. The first one describes broad-sense heritability, or the widest estimate of the role of genes in influencing the total phenotypic variability, as follows:

$$H^2 = \frac{V_G}{V_P}$$

This equation reflects all of the genetic contributions to the phenotypic variance, including non-additive variance. From a breeding perspective, this way of estimating heritability is not the most relevant. Selective pressure can only influence V_A , which is the genetic component of variance responsible for parent-offspring resemblance. It is described in the breeder's equation, describing narrow-sense heritability (Wray & Visscher, 2008):

$$h^2 = \frac{V_A}{V_P}$$

In the breeding world, most studies are focusing solely on the additive effects of loci on phenotypic variation. However, accurately mapping genotypes to phenotypes also requires to incorporate non-additive genetic interactions and understand how these vary with environment.

1.3.2 Gene pleiotropy: the Swiss army genes

Genetic correlation is a non-additive phenomenon referring to the proportion of variance that two traits share due to shared genetic causes, such as linkage (Falconer, 1960). It arises when a particular allele has an effect on more than one phenotype. Also referred to as pleiotropy, it reflects the capacity of a gene to be responsible for stable genetic correlations between quantitative traits (Figure 1.2).

Figure 1.2: Pleiotropy between phenotypic modules in *Drosophila* spp. Grey lines connect modules with a significant overlap of greater than 4 genes. Source: Ayroles et al., 2009.

It is of fundamental importance to account for pleiotropy when predicting correlated responses to artificial selection (Hill & Robertson, 1966). Usually, positive correlations occur between traits that share a common biological process (Mitchell-Olds et al., 2007) and negative genetic correlations are often found between components of fitness (Roff &

Fairbairn, 2007). Genetic correlations are useful in plant and animal breeding as they allow the substitution of more easily measured but highly genetically correlated traits (Crabbe et al., 1990). By taking into account that selecting for/against a certain trait will also increase/decrease all traits that are positively/negatively correlated, more accurate breeding values can be predicted (Crossa et al., 2010), and unexpected artificial selection consequences can be avoided.

1.3.3 Epistasis and other context-dependent effects

Pleiotropic effects can themselves be genetically variable when differences in epistatic interactions occur between loci that affect multiple traits (Wang et al., 2010). In quantitative traits, epistasis describes the situation in which the expression of one gene is affected by the expression of one or more independently inherited genes. For instance, if the expression of gene *A* depends on the expression of gene *B*, but gene *B* becomes inactive, then the expression of gene *A* will not occur. Epistatic interactions are common, for instance between nuclear and mitochondrial genes (Hoekstra et al., 2013; Kasashima & Endo, 2015), but these remain difficult to study because of methodological limitations. The significance threshold to detect these interactions lowers when the number of pairwise marker/marker interaction to test increases, therefore requiring a larger number of samples from mapping populations. Epistasis explains part of the “missing heritability” problem mentioned earlier. Geneticists used to assume that the proportion of the phenotypic variance explained by all variants affecting the trait (h^2_{all}) was equal to the estimate of apparent heritability (h^2_{pop}). This was based on the assumption that the trait was only composed of additive effects, i.e. not taking into account V_I , V_D , V_E or $V_{G \times E}$. For traits with genetic interactions, h^2_{pop} may significantly exceed h^2_{all} : even when all the variants have been discovered, the estimate of the missing heritability will not converge to zero with increasing sample size. Instead, it will converge to $1 - (h^2_{\text{all}} / h^2_{\text{pop}})$. With larger analyses, the proportion of heritability explained has grown to 20–30% in some well-studied cases and > 50% in a few, but, for most traits, the majority of the heritability remains unexplained (Weedon et al., 2008). Genetic interactions may inflate the apparent heritability and are hard to distinguish with standard methods (Zuk et al., 2012). For instance Howard et al. (2014) compared estimates of accuracy among 10 parametric and 4 non-parametric methods. They simulated a factorial design including different progeny types (F_2 and backcross), heritability estimates (0.30 and 0.70) and types of genetic architecture (additive

and epistatic). For additive genetic architectures, there were no significant differences of estimated accuracies among the methods. However, non-parametric methods produced more accurate predictions than parametric ones for genetic architectures including epistatic effects. To answer the need for more accurate predictions to achieve genetic gains, complex trait studies are now commonly linking statistical estimates of epistasis to their biological meaning (Cooper et al., 2020). This can be achieved by plotting two-locus genotype–phenotype patterns for epistatic pairs in non-parametric models connected to experimentally determined gene-interaction (Carlborg & Haley, 2004) or by using gene regulatory networks to infer the genetic regulatory structures, such as positive- or negative-feedback loops (Gjuvsland et al., 2007).

1.3.4 QTL mapping

Quantitative Trait Loci (QTL) are regions of the genome that have a statistical association of varying strength with a phenotypic trait, and these traits can be of discrete or continuous nature. With the democratisation of high-throughput sequencing methods (e.g. Illumina multiplex sequencing-by-synthesis platforms), QTLs been relatively precisely identified using thousands of SNPs. When a sufficient number of genome-wide markers have been genotyped in a mapping population, a detailed record of associated phenotyping data is used to test for statistical associations. The four widely used methods for identifying QTLs are 1) single-marker analysis, 2) interval mapping by maximum likelihood, 3) interval mapping by regression, and 4) composite interval mapping. In most cases, the first step towards defining a QTL is to group together all of the linked markers. A linkage group is established when two or more SNPs are located in close physical proximity to each other on the genome and are inherited together. The level of correlation between the markers is measured as linkage disequilibrium (LD), which is determined by the spatial distance between them and the amount of recombination events in that area of the genome. A QTL can be defined when a marker within a linkage group shows a statistically significant correlation to a phenotype. One way to identify a QTL is to use the logarithm of the odds (LOD) score, which compares the likelihood of obtaining the phenotype if the two loci are linked, to the likelihood of observing the same phenotype purely by chance. A positive LOD score indicates the presence of linkage, whereas negative LOD scores suggest that the probability of linkage is less likely.

Over the last 40 years, QTL mapping has enabled the discovery of a vast number of variants in disease resistance in plants (Concibido et al., 2004; Shi et al., 2018), genetic variation of natural populations (Schielzeth & Husby, 2014; Slate, 2005), traits of commercial relevance for livestock (Legarra et al., 2015; Yassumoto et al., 2020) or even human diseases (Liu, 2018; Mell et al., 2019) to name only a few. However, a known issue of this type of analysis is that only the allelic diversity that segregates between the parents of the mapping population can be assayed. Therefore, a bottleneck remains the number of individuals needed to achieve sufficient power (Ashton et al., 2017), and QTLs found in association with a trait remain population specific, rather than commonly implicated in the phenotype investigated.

1.3.5 Genome-wide association studies

More recently, genome-wide association studies (GWAS) have emerged as a complementary tool to QTL mapping. This technique also identifies associations between markers and phenotypes based on LD (Meuwissen et al., 2001), but does not require *a priori* pedigree information. It has therefore been widely employed to detect QTLs both in captive (Palaikostas et al., 2018a) and natural populations (Santure & Garant, 2018). GWAS can be conducted on genetically diverse, unrelated individuals and are particularly advantageous when controlled crossing or generation of large mapping populations is difficult. However, the power of a GWAS relies on the number of markers in relation to the extent of LD in the population (Korte & Farlow, 2013). When applied to natural populations or collections of outcrossing individuals characterised by a rapid LD decay, tens of thousands of markers are required to obtain an adequate level of resolution, although this number varies depending on genome sizes. Another disadvantage of GWAS is its small power to detect rare allelic variants, whose frequency can be increased in controlled crosses and being then captured with QTL mapping (Figure 1.3). Finally, because of intrinsic limitations of statistic methods, there is also a higher frequency of false discovery of loci in GWAS than in linkage mapping (Hayes, 2013).

Figure 1.3: Relationship between effect size and allele frequency. Rare genetic variants with large effect sizes are often identified in family-based QTL mapping, while common genetic variants with small effect sizes are easier to detect using GWAS on a whole population.

GWAS has proved useful for fine mapping of loci associated with numerous complex traits (Li et al., 2018; Palaiokostas et al., 2018). Combining GWAS, which detects common variants, and QTL mapping, which has the power to detect rare variants, provides more reliable results than using one of these methods alone (He et al., 2013; Sonah et al., 2015).

1.4 Environmental influence on traits

Accounting for additive and non-additive genetic effects is important to explain phenotypic variation. However, phenotypic changes can be due to non-genetic changes (Lin et al., 2019; Marine et al., 2020; Réale et al., 2003). These are explained by the 2nd and 3rd terms of *Equation 1* on phenotypic variance. The environment in which species live is not constant. Variations occur in time and space and organisms cope with environmental fluctuations through different strategies (e.g. regulation of physiological systems, reallocation of energy towards defensive mechanisms, and/or behavioural change to cope

with or avoid changes). For ectotherm species such as fish, which cannot regulate their internal temperature, thermal fluctuations are particularly important and they have to acclimate constantly in order to keep homeostasis.

1.4.1 Phenotypic plasticity

Phenotypic plasticity is defined in multiple ways: the ability of a single genotype to produce more than one alternative form of morphology, physiological state, and/or behaviour in response to environmental conditions (West-Eberhard, 1989); more broadly the potential for an organism to produce a range of different, relatively fit phenotypes in multiple environments (DeWitt et al., 1998) or merely an environment-dependent phenotype expression (Dewitt & Scheiner, 2004). A common feature here is that alternative alleles, or their products, react differently to the environment. This universal coping mechanism has been reported across all taxa, from humans (Feinberg, 2007), to animals (Laurila & Kujasalo, 1999; Miranda, 2017), plants (Sultan, 2000), and fungi (Slepecky & Starmer, 2009). It occurs in various aspects of an individual's life, including ontogeny (Gunter et al., 2013; Sultan, 2000), lifespan (Borges, 2009), response to biotic (Agrawal, 2001; Leal et al., 2017) or abiotic factors (Dwivedi et al., 2019; Tollrian & Heibl, 2004), pathogens or disease susceptibility (Agrawal, 1999; Feinberg, 2007), and animal behaviour (Laurila & Kujasalo, 1999; Miranda, 2017).

Phenotypic plasticity is found at different integrative levels. At the population level, it refers to the overall phenotypic response of a species to different environments. The genetic basis of those responses is mainly explained by transcriptional (Aubin-Horth & Renn, 2009; Schlichting & Pigliucci, 1993), post-transcriptional (Albertin et al., 2013), and/or epigenetic (Jonsson & Jonsson, 2019) regulations. At the individual level, phenotypic response patterns can be described using reaction norms (Hutchings et al., 2007; Scheiner, 1993). Consider two distinct families with different genotypes. A flat reaction norms will denote an absence of phenotypic plasticity in both families. The genetic differences between them are consistent in different environments (Figure 1.4 A).

Figure 1.4: Diversity of reaction norms. Mean phenotype and variance in two environments for a genotype are denoted with circles, and those for another genotype are denoted with triangles. Genotype means across environments are connected with a line, the reaction norm. Boldface terms adjacent to the panes indicate significant genetic (G), environmental (E) or gene-environment interaction ($G \times E$) variance, respectively.

If the reaction norms appear sloped and parallel, it indicates phenotypic plasticity, as well as additive genetic variation: both genotypes are similarly plastic (Figure 1.4 B). If the reaction norms are oppositely sloped, then this is a sign that different genotypes react differently to the environment: genetic variation for plasticity (i.e. $G \times E$) (Figure 1.4 C). Non-parallel reaction norms are commonly observed within the same species (Rashwan, 2016; Sae-Lim et al., 2015; Slepecky & Starmer, 2009). They are determined by allelic variations that have different effects according to the environment.

At a genomic level, phenotypic plasticity can be investigated by QTL mapping or GWAS carried out in various environmental conditions. QTL mapping has commonly been used for fungi (Wei & Zhang, 2017); and a combination of both strategies is routinely used in animals (Parks et al., 2013; Vieira et al., 2000) and plants (Yang et al., 2016). However, although the number of studies looking at variants interacting with the environment are increasing (e.g. Campitelli et al., 2016; Peltier et al., 2018; Wellenreuther et al., 2019), the experimental set up and therefore the identification of the genetic basis of phenotypic plasticity is not trivial. Other methods, complementary to QTL mapping and GWAS, can be used, one of them is presented hereafter.

1.4.2 Gene expression, the intermediate phenotype

The expression of genes into transcripts is the most fundamental level at which a genotype gives rise to a phenotype, and represents in itself a quantitative characteristic (Rockman & Kruglyak, 2006). Because they represent the phenotypes most immediately connected to DNA sequence variation, transcripts are ideally suited to serve as a bridge in studies investigating the relationship between genotype and macro-phenotypes. Essentially, the study of this intermediate layer brings an understanding of the molecular networks and physiological pathways that lead to changes in systemic phenotypes (Kitano, 2002). Changes in gene activation and transcription often occur during periods of stress and are part of the mechanisms involved in phenotype plasticity (Hodgins-Davis & Townsend, 2009). Therefore, one approach to understanding the ability of organisms to tolerate environmental stress is to identify stress-induced changes in transcript abundance (Huang & Agrawal, 2016) (note: numerous other molecular mechanisms accomplish plasticity, including translation, enzyme, transcription factor binding sites and hormonal regulation, ultimately producing local or systemic responses). These intermediate molecular phenotypes can vary genetically in populations and their location differs from one cell type or tissue to another, making it a site-specific investigation. Another interesting feature of gene expression studies is that it allows to examine plastic responses for very large sets of traits (Swindell et al., 2007; Wellenreuther et al., 2019) compared with classic macro-phenotypes. Although relatively unbiased, the significant increase in the number of traits accessible raises new challenges for the analysis and interpretation of data. These have become a central challenge for genome-wide genetic mapping of gene expression and require the development of new statistical techniques as well as the improvement of existing methods, some of which are discussed in the next section.

1.4.3 RNA-sequencing and statistical frameworks

To quantify genome-wide expression levels, messenger RNA (mRNA) is isolated from a sample of a cell type or tissue, fragmented at random position and reverse transcribed into complementary DNA (cDNA). Fragments meeting a certain size are amplified by Polymerase Chain Reaction (PCR). cDNA is then sequenced using NGS and the resulting reads are aligned to a reference genome, if available. High-throughput RNA sequencing (RNA-seq) (Wang et al., 2009) is now preferred over micro-arrays because it

allows the discovery of new variants without any pre-requisite knowledge of the genomic content. This method is based on read counts that are absolutely or probabilistically assigned to transcripts (Oshlack et al., 2010). RNA-seq data therefore consist of integer counts, unlike microarrays which yield intensities that are essentially continuous measurements. A number of early publications applied statistical methods developed for microarrays to analyse RNA-seq read counts. These methods included Generalized Linear Models with over-dispersion (Lu et al., 2005), Poisson distributions (Oshlack et al., 2010), Poisson distributions with moderated test statistic approach (Smyth, 2004) or log-count analyses after normalisation by sequencing depth, such as included in the limma package (Cloonan et al., 2008). Estimates were applied to the average gene expression level, forgetting that the dynamic ranges in RNA-seq data are discrete and larger than in continuous microarray data. This led the probability of detecting highly expressed genes as differentially expressed to be greater than that for low-count genes (Oshlack & Wakefield, 2009). RNA-seq data requires to be analysed by statistical methods designed specifically for discrete counts. A distributional assumption is needed as we estimate the probability of extreme events (large fold changes appearing by chance) from limited replicates. Much interest has focused on the Negative Binomial distribution (NB) as a model for read counts. The NB is a generalisation of the Poisson distribution. It allows for additional variances (called over dispersion), beyond the variance expected from randomly sampling from a pool of molecules that are characteristic of RNA-seq data. The advantage of the NB resides in its two parameters: (1) the observed data at the gene level uses counts or estimated counts of fragments for each feature. The spread of values among biological replicates arises as a continuous mixture of Poisson distributions (i.e. a compound probability distribution); and (2) the mixing distribution of the Poisson rate is a gamma distribution, which is a better representation than a simpler, one parameter Poisson distribution.

Another obstacle to applying normal-based statistical methods to read counts is that the counts inevitably have unequal variabilities, even after log-transformation. Differential expression analysis requires that gene expression values are compared among samples. Large counts have a much larger standard deviation than small counts. An important step is therefore to normalise sequencing depth, which is the most important factor when comparing samples. This can be done directly or by accounting for the number of transcripts, which can differ significantly between samples (Conesa et al., 2016). While a logarithmic transformation counteracts this, it overdoes the adjustment somewhat, so that

large log-counts will have smaller standard deviations than small log-counts (Bullard et al., 2010). Several pipelines relying on different normalisation approaches are available for researchers to use. On one hand, count-based methods such as edgeR (Robinson et al., 2010), DESeq (Trapnell et al., 2013), DESeq2 (Love et al., 2014), baySeq (Hardcastle & Kelly, 2010), TSPM (Auer & Doerge, 2011), PoissonSeq (Li & Li, 2012) and DSS (Wu et al., 2013). The latter aims to specify the exact probabilistic distribution of counts based on NB or over-dispersed Poisson distributions. On the other hand, limma-trend and limma-voom (Law et al., 2014), which are derived from microarray models. Simulations have shown that a key advantage of the limma pipelines is that they seem to provide accurate type I error rate control when the number of RNA-seq samples is small (Law et al., 2014). Ultimately, the choice of the tool used should be made in function of the power to control error rate, which will be dictated by the experimental design (number of samples, number of replicates, library preparation, sequencing depth, inherent biological variability).

1.5 Growth is a complex trait

In vertebrates, somatic growth is a complex quantitative and highly polygenic trait (De-Santis & Jerry, 2007; Posbergh & Huson, 2021; Yuan et al., 2018). It is sensitive to a range of intrinsic (within individual), natural and extrinsic factors (Enberg et al., 2012; Morrongiello & Thresher, 2015). At the physiological level, the endocrine regulation of growth is mediated by the somatotrophic axis (Cryer & Cherring, 2001), through the pituitary growth hormone (Gh) (Rousseau & Dufour, 2007), growth hormone receptors (Ghr), insulin-like growth factor-I (Igf-I), IGF-I receptors (Igfr), and IGF binding proteins (Igfbp) (Reinecke, 2010). Also highly affected by nutritional and environmental factors (Pérez-Sánchez et al., 2018), the somatotrophic axis is conserved throughout vertebrates (Cowan et al., 2017). For the purpose of this thesis, the next section briefly describes its mechanism in teleost fish.

1.5.1 Endocrine regulation of somatic growth

The somatotrophic axis comprises the hypothalamus, the pituitary gland (referred to as pituitary hereafter), and the liver. The central nervous system perceives environmental cues through the hypothalamus, and communicates them to the endocrine system,

represented by the pituitary (Löhr & Hammerschmidt, 2011). The latter consists of two lobes: the neurohypophysis, which contains nerve bundles originating from the hypothalamus, and the adenohypophysis, which secretes hormones, such as Gh, to the rest of the body (Machluf et al., 2011). Gh functions as an important regulator of metabolism and somatic growth in fish species (Canosa et al., 2007). It binds to its receptor Ghr present at the surface of the cells of target tissues (Fuentes et al., 2013), such as the liver. There, it stimulates the production of Igf-I and -II (Le Roith et al., 2001, Figure 1.5). Igf are involved in processes such as the stimulation of protein synthesis, proliferation, differentiation, skeletal muscle elongation, cell migration, and survival (Duan et al., 2010; Reinecke, 2010). In fish, both Gh and Igf-I promote growth, although Igf-I seems to be ultimately responsible for body growth (Picha et al., 2008).

Figure 1.5: Schematic representation of the endocrine regulation of growth in fish and the hypothalamic and pituitary hormones involved in such regulation that have been related to a role in the control of food intake and/or energy balance (modified from Blanco, 2020).

1.5.2 The impact of temperature

As cold-blooded animals, fishes are heavily influenced by water temperature. Thermal fluctuations are perceived by the brain, in turn influencing the gene expression and function of various proteins, including growth hormones, to regulate endocrine functions. Average daily levels of Gh and Igf-I have been shown to present seasonal variations in fishes, which are tightly correlated with annual variations in water temperature, although photoperiod length also has a strong influence (Cowan et al., 2017; Marchant & Peter, 1986; Triantaphyllopoulos et al., 2020). Peak in Gh production has been shown to coincide with the highest water temperature, whereas annual peaks of Igf-I present a delay of some months after the Gh peak (Deane & Woo, 2009). Interestingly, the highest annual growth rates coincide with peaks in Igf-I rather than Gh, reinforcing the idea that Igf-I is ultimately responsible for growth in fishes and not Gh like in other vertebrates (Deane & Woo, 2009).

Taken to the organismal level, large differences in sizes and growth rates have been shown to be induced by small differences in average temperature (Sebens, 1987). The simplest way to describe the effects of temperature on an organism's growth is to generate a thermal performance curve (TPC) (Schulte et al., 2011). Many studies have documented the shapes of TPCs across levels of biological organisation and taxa. TPCs tend to be unimodal and left skewed with three distinct regions: (1) a rising phase as temperature increases; (2) a plateau phase that encompasses the thermal optimum (T_{opt}) for the trait; and (3) a steep falling phase at higher temperatures (Figure 1.6).

Although species share those common three regions, the overall shape of the TPC varies among taxa. This is in part the result of adaptive differences, but can also be caused by neutral variation and various types of plasticity, including gene expression, epigenetic effects, developmental plasticity and acclimation (Schulte et al., 2011). Due to phenotypic plasticity, the breadth of the thermal range within a species also varies among populations, with age, or within a life stage (Bowler & Terblanche, 2008).

Figure 1.6: A thermal performance curve (TPC) summarising the effects of temperature on the rates of biological processes. Critical features of a TPC are the slope of the increase in rate with temperature, the thermal optimum for the process (T_{opt}) and the thermal breadth, which is the range of temperatures over which the rate of the process is maximised (modified from Schulte et al., 2011).

Human-induced climate change is altering patterns of mean and extreme temperatures across the globe (IPCC et al., 2018). Ectotherm species must find ways to cope with changes in their environment, and depending on the success of these strategies, different outcomes are possible: (1) extinction or extirpation, (2) migration or shift in current distribution range, (3) adaptation through changes in the genetic composition of the population, or (4) acclimation through phenotypic plasticity. Shifts in behaviour like locomotion or foraging activity (Abram et al., 2017; Huey & Kingsolver, 1989), as well as in geographic distributions and abundance of species (Brun et al., 2020; Parmesan, 2006; Woodin et al., 2013) have started to be observed. Already, studies are finding that increased temperatures are causing fish growth to decline (Schaal & Lotterhos, 2021), which is of particular concern, especially from an aquaculture point of view.

1.5.3 Genomic insights to mitigate the effects of temperature variations

Genomic information can be used by breeding programmes to produce phenotypes that compensate for growth losses in condition of sub-optimal temperatures. This can be done through direct selection for faster growth (so that the breadth of the optimal growth temperature is increased); or through selection for horizontal shifts in TPCs (so that the optimal growth temperature of a species falls into the range of the area's thermal profile). Historically, most breeding programmes have selected for faster growth rates under a range of grow-out environments (Gjedrem, 2005). The focus on increasing growth alters both growth rate and total mass at the end of a grow-out cycle. Gains obtained can reach beyond the maximum change in annual growth rates and total annual growth (Klinger et al., 2017). For example, selection for growth in Atlantic salmon (*Salmo salar*) increased harvest weight by 12% per generation. The cumulative genetic gains over multiple generations of selection is then around 200% (Janssen et al., 2017). In the gilthead seabream (*Sparus aurata*), genomics selection led to an increase in harvest weight of 10–15% per generation, resulting in a cumulative genetic gain over multiple generations of 100% (Janssen et al., 2017). These values are greater than the 25–41% total increase in annual growth required to offset the most extreme climate-induced decreases (Klinger et al., 2017), indicating that selection-based methods could compensate for the potential losses from climate change. In contrast to selecting for faster growth across all temperatures, few breeding programmes have explicitly selected for shifts in the optimal temperature for growth. *GxE* interactions can potentially result in a re-ranking of individuals within a breeding programme, such that a selected genotype performs well at one temperature but poorly at another (Sae-Lim et al., 2016). This risk seems to be population specific. A review found that the environment explained only a small portion of the variation in growth rates (e.g. 1–5%) as fishes of different genotypes were exposed to a range of grow-out conditions (Gjedrem & Baranski, 2009). This would indicate little horizontal shift in TPCs. Another review, however, found strong evidence of re-ranking in rainbow trout (*Oncorhynchus mykiss*) and gilthead seabream genotypes exposed to different temperatures, indicating strong *GxE* interactions between growth and temperature, and potential for horizontally shifting TPCs (Sae-Lim et al., 2016). Finally, a study on the Atlantic silverside (*Menidia menidia*) found that cold and warm adapted populations of the same species undergo both absolute changes in the growth rates (e.g. cold adapted populations with shorter growing seasons grow faster at all temperatures) and horizontal shifts in TPCs for growth (Yamahira & Conover, 2002).

1.6 Wider relevance of this research

1.6.1 Aquaculture under climate change

Aquaculture is the fastest growing food-producing sector, and accounts for over 50% of global fish production (FAO, 2018, Figure 1.7), promoting the sector as an important part of the solution for meeting the growing food demands of this century (Béné et al., 2016). Currently, 694 aquatic species are farmed worldwide, supporting millions of people through the provision of nutrition, food security and livelihoods, as well as through the alleviation of poverty (Galappaththi et al., 2020). According to the Global Aquaculture Alliance, 62% of food fish will come from aquaculture by 2030 (GAA, 2021). The recent Intergovernmental Panel on Climate Change 1.5°C report (which assesses the effects of climate change and its potential future risks) identified the aquaculture sector as one of the key areas that will be impacted (IPCC et al., 2018).

Figure 1.7: World capture fisheries and aquaculture production. Source: FAO, 2018.

Despite its critical role in global food security, the vast majority of aquaculture's breeding programmes are based on stocks without genomics-informed selective breeding (Gjedrem & Baranski, 2009; Janssen et al., 2017). Aquatic breeding schemes tend to lag behind their terrestrial livestock counterparts in terms of the uptake of genomic

technologies, and for many species, molecular tools are only applied for pedigree reconstruction and subsequent family selection (Chavanne, 2016). However, aquaculture is rapidly diversifying and more genomic information are being generated: genetic parameters are estimated for commercially important traits (e.g. growth, feed conversion) to help improve conventional selective breeding; and new technologies (e.g. marker-assisted selection, genomic selection and genome editing) are integrated into selective breeding programmes to accelerate gains (Shen & Yue, 2019). There is a very high potential for cumulative and permanent improvement of production traits, as shown by the typically high genetic gains reported in programmes mentioned earlier.

As aquaculture develops and grows in response to demand, so is its need for space, which is driving the relocation of cultivation systems from in-land to offshore. In open-ocean, control over the environment is minimal and fish growth and production are strongly constrained by the physical state of the ocean, including ambient seawater temperatures (Klinger et al., 2017). The increase of extreme weather events due to climate change are likely to happen over the lifespan of a commercial operation, which leaves breeding programmes with at least three choices:

- 1) Move fish to a more favourable environment, although this is economically costly, logistically difficult and politically challenging (Lovatelli et al., 2013).
- 2) Grow a different species whose optimal growth range is more in alignment with the new environmental conditions. However, changing species requires developing new expertise among staff, additional legal permits and fees, as well as access to broodstock (Leung et al., 2008).
- 3) Use selective breeding to alter the thermal optimum for growth of the currently bred species, to overcome impacts of climate change in the operational area (Klinger et al., 2017).

1.6.2 The role of aquaculture in New Zealand

Island countries surrounded by water, such as Aotearoa New Zealand, have a strong interest in developing and using their coastline, while protecting ecosystems from degradation by the introduction of exotic species. The waters around Aotearoa support significant commercial fisheries of both inshore and deep-water species (Fisheries New

Zealand, 2019). However, these fisheries are fragile, and most of the quotas are regularly exceeded (Fisheries New Zealand, 2019). Aquaculture has been recognised as an alternative to substitute captured fisheries and help alleviate natural stock depletions. Although the fastest growing sector of agricultural production in the country, programmes rely almost exclusively on the farming of the three species: Greenshell™ mussels (*Perna canaliculus*); Pacific oysters (*Crassostrea gigas*); and only one finfish species, chinook/king salmon (*Oncorhynchus tshawytscha*), which was introduced from North America (Camara & Symonds, 2014; Davies et al., 2019). The paucity of species means that there is a need to add resilience to the sector by diversifying the range of species farmed. Furthermore, the aquaculture industry has set an ambitious target to increase its revenues to \$1 billion by the year 2025 through a combination of expanding production of the existing species, and adding new, high-value native species to the list of farmed species (Camara & Symonds, 2014).

1.6.3 Trevally (*Pseudocaranx georgianus*) – araara

The study species for this thesis research was the silver trevally (*Pseudocaranx georgianus*, Cuvier 1833) – referred to as “trevally”. In Aotearoa, its Māori name is araara. Indigenous Māori people have a strong cultural connection to trevally, where it is considered as taonga (i.e. has value, or is treasured). Trevally is a shoaling pelagic species found throughout the coastal waters of southern Australia and around New Zealand (Gomon et al., 2008; Smith-Vaniz & Jelks, 2006). It is most common at depths of approximately 80 m, although its range is thought to be 10–238 m (Mundy, 2005) and can reach over 40 years of age (Fisheries New Zealand, 2019). In many regions of NZ, trevally is a major component of recreational and commercial fisheries (MPI, 2021). The importance of inshore fin fish such as trevally is reflected in the National Fisheries Plan, which states that “They contribute to our cultural and social traditions, to our nation’s economy and to our sense of overall well-being”. Trevally is both targeted and caught as bycatch within inshore bottom trawls and purse seine fisheries. In May 2018, it had been reported that 85% of available trevally Annual Catch Entitlements (ACE) for the 2017/18 fishing year had already been caught. Given that approximately 30% of landings of the two previous fishing years occurred between June and the end of the year, the landings were

again predicted to exceed the available ACE for the 2017/18 period (Fisheries New Zealand, 2019).

The *Pseudocaranx* genus belongs to the carangid family, which contains 146 species within 30 genera, distributed in tropical waters in the Atlantic, Indian, and Pacific oceans (Roberts et al., 2015). This family includes species with well-established breeding programmes around the world, like for the yellowtail kingfish (*Seriola lalandi*) (Orellana et al., 2014; Poortenaar et al., 2001). Species of the *Pseudocaranx* genus appear to do well in farm-like conditions, as demonstrated by long-lasting successful cultures of striped jack (*P. dentex*) in Japan where it has been bred since the 1990s (Fukusho, 1995). However, compared with its sister species who have faster growth rates (e.g. Abbink et al., 2012; Fernández-Montero et al., 2018; Skaramuca et al., 2001), trevally is much slower, which means that its growth rate will likely need to be improved. Finally, although commercially valuable, there has been limited genetic research reported on this species. Recently, a few studies have been carried out using genetic tools to elucidate the *Pseudocaranx* phylogeny (Bearham et al., 2020) and investigate the genetic structure of wild populations around the coast of New Zealand (Kemp, 2020), however, no genomic or QTL mapping studies have been carried out on trevally to date.

1.7 Aims and thesis structure

This PhD research was undertaken using the trevally population held at The New Zealand Institute for Plant and Food Research Limited (Plant & Food Research) in Nelson, New Zealand. The research was funded by the Ministry of Business, Innovation, and Employment (MBIE) programme “Accelerated breeding for enhanced seafood production” (#C11X1603). The first general aim of this PhD research was to investigate the pedigree structure and genetic diversity of a new trevally population, the genetic basis of growth traits, and how they are influenced by the environment. A second general aim of the thesis was to produce a range of genetic resources that could contribute to ongoing genome investigations in this and related fish species.

1.7.1 Specific objectives

- Determine the pedigree structure and inbreeding levels of the existing population.
- Describe growth patterns in trevally.
- Calculate key genetic parameters of 10 growth traits.
- Develop a high-density linkage map, used to support the genome assembly and position the markers for QTL mapping.
- Combine QTL mapping and GWAS to scan for loci affecting growth rate.
- Assess the influence of temperature on the phenotypic plasticity of growth.
- Annotate the trevally transcriptome and find differentially expressed genes in two of the major organs involved in somatic growth in fish.
- Identify major key physiological pathways involved in the modulation of phenotypic plasticity of growth.
- Discuss the implications of the genetic results for the breeding programme.

1.7.2 Thesis structure

Each chapter of this thesis (except Chapters 1 and 5) has its own specific objectives and is divided into sections including introduction, methods, results, and discussion. Chapters 2, 3, and 4 have been prepared as manuscripts for publication and are in varying stages of being submitted.

Chapters 1 and 5 present an introduction and final discussion to tie together the results of all the other chapters. Specifically they discuss the broad implications of the data chapters and how the results can be applied to the trevally breeding and management.

Chapter 2 is the first data chapter. This chapter provides the groundwork for the following chapters by conducting a review of the pedigree structure of the study population. It describes in detail the Whole Genome Sequencing (WGS) and Genotyping-by-Sequencing (GBS) methods used to generate the genotype datasets that were also used in Chapter 3. It also investigates the seasonal growth patterns of trevally as well as heritability and genetic parameters for these growth traits.

Chapter 3 is the second data chapter. This chapter investigates the genetic architecture of growth traits and constructs a linkage map. It reports the first high-density linkage map computed for trevally. It also demonstrated that QTL mapping and GWAS used in parallel are a powerful approach for the identification and validation of loci controlling complex traits.

Chapter 4 is the third data chapter. This chapter investigates the plasticity in gene expression patterns and growth in response to different temperatures. To do so, morphological phenotypic differences in growth were measured, and pituitary gland and liver gene expression patterns determined.

Chapter 5 is a general discussion of the major findings of this PhD research, the implications for the implications for New Zealand aquaculture industry breeding programmes, and some future research directions.

2 Evaluating new species for aquaculture: a genomic dissection of growth in New Zealand silver trevally (*Pseudocaranx georgianus*)

Authors

Noemie Valenza-Troubat, Elena Hilario, Sara Montanari, Peter Morrison-Whittle, David Ashton, Peter Ritchie, Maren Wellenreuther.

Author Contributions

Conceived and designed the experiments: NVT, PR, and MW; Data collection: NVT; Laboratory work: NVT; Data analysis: NVT; Wrote the primary manuscript: NVT; Prepared final manuscript: NVT, PR, and MW; Advised laboratory or data analysis: EH, SM, PMW and DA.

Publication status

This chapter has been accepted for publication in Evolutionary Applications.

2.1 Abstract

Aquaculture is the fastest growing food production sector worldwide, yet industry has been slow to implement genomic techniques as routine tools. Applying genomics to new breeding programmes can provide important information about pedigree structure and genetic diversity; key parameters for a long-term success. It can also provide insights on potential gains for commercially important, yet complex, quantitative traits such as growth rate. Here we investigated a population of 1100 captive-bred F₁ silver trevally (*Pseudocaranx georgianus*), a promising new species for New Zealand aquaculture. We used whole-genome information, coupled with image-based phenotypic data collected over two years, to build the pedigree of the population, assess its genetic diversity, quantify 10 growth traits and estimate their genetic parameters. Successful parentage assignment of 664 F₁ individuals showed that the pedigree consisted of a complex mixture of full- and half-sib individuals, with skewed reproductive success among parents, especially in females. Growth patterns showed seasonal fluctuations (average increase across all traits of 27.3% in summer, and only 7% in winter) and strong inter-family differences. Heritability estimates for growth traits ranged from 0.27 ± 0.07 to 0.76 ± 0.06 . Genetic and phenotypic correlations between traits were high and positive, ranging from 0.53 ± 0.11 to 1.00 ± 0.00 and 0.50 ± 0.00 to 1.00 ± 0.00 , respectively. The implications of these findings are threefold: first, the best on-growing conditions are in warmer months, where highest growth peaks can be achieved; second, size- and family-based selection can be used as early selection criterion if pedigree structure and inbreeding levels are closely monitored; third, selection for body length results in concomitant increases in height and weight, traits of paramount importance for aquaculture. It is concluded that there is substantial potential for genetic improvement of economically important traits, suggesting that silver trevally is a promising species for selective breeding for enhanced growth.

2.2 Introduction

Aquaculture is the fastest growing food production sector worldwide (FAO, 2018). Compared with terrestrial breeding programmes, the industry has, however, been slow to implement the use of genomic information to inform selective breeding decisions. This is despite the proven utility of genomics to provide insights into important commercial traits,

such as faster growth and disease resistance (Gjedrem et al., 2012). In New Zealand, aquaculture production relies almost exclusively on the farming of three species: GreenshellTM mussels (*Perna canaliculus*); Pacific oysters (*Crassostrea gigas*); and only one finfish, chinook/king salmon (*Oncorhynchus tshawytscha*), introduced from North America (Camara & Symonds, 2014; Davies et al., 2019). The paucity of species means that there is a strong need to add resilience to the sector by diversifying the range of species farmed.

One of the most important and well-studied traits in breeding programmes is growth rate. This is because it is easily quantified and often highly heritable, and has a huge impact on commercial returns. Improvements in growth rates reduce the overall time that is needed to raise individuals to market size, thereby speeding up the turnover of production stock and decreasing costs (Gjedrem, 2005; Ye et al., 2017). Usually, growth rate in animals is measured using either body weight or body length as a proxy, but as phenotyping methods improve, other traits such as width, girth, or height are starting to be used to provide additional insights (Zenger et al., 2017). However, these traits are often quantitative and have a polygenic origin (Wellenreuther & Hansson, 2016), thus limiting the gains obtained by traditional breeding based on phenotypic selection of individuals (Gjedrem et al., 2012). Genome-wide markers have been used to identify loci associated with traits of importance and to guide selective breeding to efficiently increase trait gains, but the application of genomic approaches has long been limited to model species or species of high economic value. Recently, the decrease in DNA sequencing and genotyping costs has meant that large numbers (e.g. thousands) of genome-wide markers can now be easily generated for almost any species at a cost that can be afforded by research groups. Reduced representations of the genome such as in genotyping by sequencing (GBS) (Elshire et al., 2011) offer a quick and cost-effective way of genotyping a large number of individuals, while Whole Genome Sequencing (WGS) can be used to generate an accurate reference genome and in-depth genomic information for a population. These technologies have opened-up opportunities for high-throughput and high-density genotyping in a wide range of non-model species.

Genomic insights can inform a number of breeding decisions. First, a potential barrier for new breeding programmes is the lack of prior knowledge on key parameters such as the relatedness among individuals and number of contributing parents (Falconer & Mackay, 1996). This information is particularly challenging to gather for aquatic species where the founding individuals might be caught from the wild and/or exhibit mass

spawning behaviours. If not carefully monitored, poor management can lead to inbreeding pressure and early loss of genetic diversity, which must be avoided for a successful long-term breeding programme (Wang et al., 2002). Second, a useful parameter to assess is the selection potential of a trait. Estimates of heritability provide insights into the relative contribution of genetic components compared with environmental factors. Knowing if the heritability of a trait is considered as high (~ 0.9) or low (~ 0.1) (Wray & Visscher, 2008) will indicate how well the trait will respond to selection. Broad sense heritability is defined as the ratio of the total genetic variance on the total phenotypic variance, and includes effects such as dominance and epistasis, which do not respond to selection. Narrow sense heritability reflects the ratio of the additive genetic variance on the phenotypic variance and represents the part that will respond to selection (Wray & Visscher, 2008). Third, another application of genomic tools for predicting the response to selection is estimating genetic correlations between two or more traits (Lynch, 1999). Because the genes that contribute to traits can be highly genetically correlated and co-inherited, it is particularly important to know the magnitude of these correlations when predicting breeding values. The improvement of multiple traits simultaneously may seem efficient but, because of unexpected correlations, it could result in the co-selection of undesired phenotypes (Falconer & Mackay, 1996).

In this study, we applied for the first time genomics tools to a newly formed population of silver trevally (*Pseudocaranx georgianus*, Cuvier 1833) (referred as trevally hereafter), a shoaling pelagic species found throughout the coastal waters of southern Australia and around New Zealand (Gomon et al., 2008; Smith-Vaniz & Jelks, 2006). In Aotearoa New Zealand, indigenous Māori people have a strong cultural connection to trevally, where it is considered as taonga (i.e. has value, or is treasured) and is referred to as araara. In many regions, trevally is a major component of recreational and commercial fisheries (MPI, 2021). *Pseudocaranx* is also a genus that appears to do well in farm-like conditions, as demonstrated by the fact that other species in this genus have already been successfully developed for large-scale commercial farming in Asia (FAO, 2018). Furthermore, this genus belongs to the Carangidae family, which contains other species with well-established breeding programmes around the world, e.g. for the yellowtail kingfish (*Seriola lalandi*). The goal of this study was to use genome-wide data coupled with extensive phenotypic data to provide the first quantitative assessment of growth traits in a population of 1100 captive-bred F₁ trevally. Specifically, 1) we used a mixture of WGS

and GBS data to reconstruct the molecular pedigree of the population, 2) calculated inbreeding values within each generation, 3) used image-based phenotyping data collected over a period of two years to describe the growth patterns, and finally 4) estimated the heritability and genetic correlations of ten growth traits to determine their potential for enhanced growth performance through a selective breeding programme.

2.3 Methods

2.3.1 Populations studied and holding conditions

The trevally population studied consisted of an F_0 wild broodstock ($n = 22$) and a captive reared F_1 population ($n = 1100$). F_0 individuals were originally captured during two net tows in February 2012 in the North Taranaki Bight (Lat. 3845267 – Long. 17420626 and Lat. 3851887 – Long. 17419780). Live fish arrived two days later at the Maitai Finfish Facility operated by The New Zealand Institute for Plant and Food Research Limited (PFR) in Nelson, New Zealand, where all research was subsequently carried out. The Finfish Facility receives ambient seawater from an underground bore, which is filtered using mesh filters and UV treatment. The F_1 generation was produced using hormone-induced mass spawning in December 2015. Induced spawning of F_0 individuals was achieved subsequent to an intramuscular injection of human chorionic gonadotropin (hCG, Choloron[®]) at a target dose of 600 IU/kg of bodyweight. Leading up to spawning, surviving parents ($n = 19$) were fed a specialised diet containing fresh fish and oil supplements. Following injection, two individuals became egg bound and died, most likely before spawning. Spawning occurred 48 hours post-injection of hCG. Fifty grams of eggs were collected each day from the tank outlet over three consecutive days and placed in 450L hatchery tanks provided with a 12-hour light cycle, gentle aeration and water flow. At seven days post-hatching, larvae were combined into a single 5000L tank. The larvae were fed a combination of live rotifers and artemia. At 1 month old, the juveniles were then fed a combination of artemia, O.range (NRD) dry crumbs, and a wet diet consisting of minced fresh fish. At 6 months of age, all fish were transferred into a single 5000L tank with natural lighting and fed a combination of dry commercial pellets (Skrettings Nutra RC 1.2 mm and 1.8 mm) until 1 year old, and then Ridley pellets (2 mm, 3 mm, 4 mm and 6 mm) and a wet diet (fish mince or portions of fresh fish). In November 2017, at 2.1 years old, 1100 F_1 were

randomly selected and transferred to a single 13,000L tank. During this time, all fish were tagged by inserting a Passive Integrated Transponder (PIT) tag (GPT12, Biomark[®]) into the body cavity. To minimise the risk of overcrowding and stunting of growth, F₁ were moved for a final time four months later (March 2018) to a single 50,000L tank and were maintained under the same ambient temperatures and photoperiod for the remainder of the experiment.

2.3.2 Tissue sampling, DNA extraction and library preparations

Thirteen surviving F₀ were tagged and fin clipped in January 2017. Fin clips were placed directly into chilled 96% ethanol, heated to 80°C for 5 min within 1 h of collection, and then stored at –20°C until needed. Total DNA was extracted as described in Ashton et al. (2019) with the following modifications: proteinase K digestion time was increased to 1.5 h and the 80°C inactivation step was omitted; the RNA removal was performed after the salting-out step; and the DNA was quantified by fluorescence using the Qubit High Sensitivity dsDNA kit (Thermo Fisher Scientific, Waltham, USA) used in accordance with the manufacturer's instructions. DNA quality was assessed by agarose gel electrophoresis (average fragment size ~40 kbp) and using spectrophotometry (absorbance ratios at 260/280 nm and 260/230 nm). Illumina shotgun fragment libraries with an insert size of at least 125 bp were generated for each of the 13 individuals and sequenced (paired-end, 125 bp reads) over three lanes of the HiSeq 2500 platform at the Australian Genome Research Facility (AGRF, Melbourne, Australia). Samples of fin tissue from 1100 F₁ individuals were collected during the first phenotyping round (November 2017) and stored as described above. Total genomic DNA was extracted by SlipStream Automation (Palmerston North, New Zealand) using the same protocol as for the F₀. Genotyping for the F₁ was carried out using a modified GBS approach (Elshire et al., 2011; Hilario, 2015). DNA integrity was checked by capillary electrophoresis (High Sensitivity genomic DNA kit), Fragment Analyzer (Advanced Analytical). One microgram of total genomic DNA was used for digestion with restriction enzymes. A double digestion was performed with *Pst* I and *Msp* I by incubation at 37°C for 3 h, the adaptor ligation step omitted drying out the DNA/adaptor mixture. The barcoded adaptors were associated with the *Pst* I cut sites and designed by Deena Bioinformatics (Wageningen, Netherlands). Adaptors were annealed according to Ko et al. (2003). A high-fidelity enzyme was used for amplifications

(AccuPrime Taq DNA polymerase High Fidelity, Life Technologies). Amplification, quality check and clean up were done separately before pooling samples. A total of 12 pools of 96 samples each were prepared and sent to AGRF for sequencing on a HiSeq 2500 platform (single-end, 100 bp reads).

2.3.3 Genotyping data quality checking and processing

Sequencing data quality for both F_0 and F_1 generations were checked using FastQC version 0.11.7 (Andrews, 2010). As the F_0 and F_1 sequence data were generated using different sequencing technologies, different filtering parameters were used.

F_0 filtering/pre-processing: raw reads from the F_0 were trimmed using trimmomatic version 0.36 (Bolger et al., 2014) (using the parameters HEADCROP: 9, TRAILING: 10, SLIDINGWINDOW: 5:20, MINLEN: 75). Read groups were added and bam files were sorted and indexed using Picard toolkit (Toolkit, 2015). The trevally reference genome developed by PFR (Ruigrok et al., 2021) was indexed using Burrows-Wheeler Aligner (BWA) version 0.7.17 (H. Li & Durbin, 2009). Reads were aligned to the reference using BWA-mem and the variant calling was done using samtools version 1.9 and BCFtools version 1.9 (Li, 2011). The samples were combined using mpileup and the call was run using the multiallelic caller option. A first round of filtering was then done using VCFtools version 0.1.14 (Danecek et al., 2011). Briefly, indels were removed from the parental call, which was subsequently filtered for high missing rates per individuals (threshold set at 0% missingness) and minimum SNP quality and depth ($Q > 10$, $DP > 9$). A further filter was applied for missing data per SNPs (set at 0% missingness). Finally, SNPs were filtered for maximum depth ($\text{maxDP} = \text{average DP} + 3 \text{ stand deviations} = 445$).

F_1 filtering/pre-processing: the F_1 samples were de-multiplexed from the 12 sequencing libraries using the process_radtags module available in the STACKs version 2.1 pipeline (Catchen et al., 2013) and the reads were trimmed using Fastq-mcf in ea-utils version 1.1.2-806 (minimum sequence length = 50, quality threshold causing base removal = 33) (Aronesty, 2013). Like for the parents, read groups were also added and the bam files were sorted, indexed, aligned to the reference using BWA-mem. The variant calling was also done using samtools and BCFtools. Using VCFtools, indels and individuals with over 50%

missing data were removed, the maximum depth was set at 8,000 and SNPs with over 20% missing data were filtered.

Recombining datasets: After indexing, a list of common SNPs between F_0 and F_1 was obtained using BCFtools isec and the two sets were merged using VCFtools vcf-merge. Finally, the resulting F_0 and F_1 dataset was filtered to keep only SNPs in common using VCFtools -positions.

2.3.4 Pedigree reconstruction and F_0 sex prediction

Sequoia version 2.0.7 (Huisman, 2017) was used in the R statistical environment version 3.2.3 (R. Core Team, 2013) to iteratively reconstruct a maximum-likelihood pedigree. To prepare the common SNP dataset for pedigree reconstruction, PLINK version 1.9 (Purcell et al., 2007) was first used to test for and discard loci in linkage disequilibrium (LD) with the --indep function, evaluating 50 SNP windows, five SNPs at a time, with a variance inflation factor (VIF) cut-off = 1.5 and a minor allele frequency (MAF) of 0.4 was set for the population. The initial parentage assignment was accomplished with this genotype file and a life history file, using the parameters MaxSibIter = 0, Err = 0.05, MaxMismatch = 10, MaxSibshipSize = 900 and Tassign = 0.5. This allowed to scan the pedigree for obvious errors, as well as for duplicates that were accidentally retained. To construct the full pedigree, the parameter data frame (= Specs) was then altered to use the initial parentage assignment as prior information and MaxSibIter = 3, MaxSibshipSize = 900, Err = 0.2, Tfilter = -2, Tassign = 0.5. All other parameters were kept as default. We assessed the accuracy of the reconstructed pedigree and the ability of the SNP dataset to correctly identify familial relationships by checking for Mendelian errors using PLINK. The pedigree was visualised using Circos version 1.2.1 (Krzywinski et al., 2009)

Broodstock sex prediction based on the molecular pedigree was confirmed *a posteriori* by collecting a gonadal biopsy from each of the 13 F_0 individuals in December 2018. In brief, broodstock in the advanced stages of reproductive development were mass-sedated in tank (25 ppm Aqui-S; Aqui-S New Zealand Ltd, Lower Hutt, New Zealand). A gonadal biopsy was taken by inserting a glass cannula (Natelson tube, 3 mm outside diameter) connected to a plastic tubing into the gonopore of the fish and applying gentle aspiration by syringe. A portion of biopsy sample was placed in Ringer's solution (180 mM

NaCl; 4 mM KCl; 1.5 mM CaCl₂; 1.2 mM MgSO₄; 3 mM NaH₂PO₄; 12.5 NaHCO₃ – pH 7.5) and from this a wet mount slide was prepared and examined under a compound microscope for the presence of oocytes or sperm.

2.3.5 Inbreeding calculations

Using the same set of filtered SNPs used for the parentage assignment, inbreeding values for each individual found with both parents in the pedigree were calculated with a method-of-moments F coefficient (F_H) using PLINK. This statistic is equal to Nei's F_{IS} statistic, but is calculated using a different formula:

$$F_h = \frac{O(Hom_i) - E(Hom)}{m - E(Hom)}$$

where $O(Hom_i)$ is the observed number of homozygous loci for the i^{th} individual, and $E(Hom)$ is the Hardy–Weinberg expected mean number of homozygous genotypes across m loci (Kardos et al., 2015). The distribution of inbreeding values was then visualised using the ggplot2 library in R. A Welch two-sample t-test was used to compare mean inbreeding values between F_0 and F_1 . To compare mean inbreeding values between families, a linear mixed model was fitted with family as a random effect to run an ANOVA. A *post hoc* Tukey correction for multiple comparisons was used to find out which family values were significantly different from the others.

2.3.6 Phenotyping, trait estimations and phenotypic correlations

In total, eight sets of external images were taken for each F_1 individual, corresponding to a measurement roughly every four months during a period of two years. Images were taken on a custom-built imaging rig using fitted Panasonic Lumix DMC-GH4 cameras. Images were analysed using PFR's Morphometric Software™ (<https://www.plantandfood.co.nz/page/morphometric-software-home/>). The software extracts the outline of each individual fish from images, locates the XY coordinates of morphometric features on the outline (e.g. upper lip and narrowest cross-section of the tail),

and then uses those coordinates to make measurements. The measurements were converted from pixels to mm using the length of rulers also present in the images.

Four growth traits were directly extracted from the pictures for each individual fish. Peduncle length (PL) was measured by locating the upper lip and narrowest cross-section of the tail and then measuring the distance between these points. Height was measured at three positions along the fish – 25%, 50%, and 75% of the way from the upper lip to the narrowest cross-section of the tail (H25, H50 and H75, respectively, Figure 2.1). To do this, the software first located a starting position along the peduncle length and then measured the distance between the top and bottom edge of the fish at a 90° angle to the peduncle length.

Figure 2.1: Morphometric traits (PL: peduncle length, H25: height at 25% of PL, H50: height at 50% of PL, H75: height at 75% of PL, EW: estimated weight, $\Delta H25$: net gain in H25, $\Delta H50$: net gain in H50, $\Delta H75$: net gain in H75, ΔPL : net gain in PL, ΔEW : net gain in EW), measured in the New Zealand silver trevally *Pseudocaranx georgianus*. Peduncle length, and the three height measurements were obtained automatically using a custom image analysis script. Height measurements were measured relative to length. Weight was estimated using a Bayesian hierarchical approach (Froese et al., 2013). Net traits were calculated using the first measurements as initial point.

In addition, six other traits were estimated: weight (EW), and net gains in height at – 25%, 50%, and 75% of PL (ΔH_{25} , ΔH_{50} and ΔH_{75} , respectively), in peduncle length (ΔPL) and in weight (ΔEW). The weight estimations were done following Froese et al. (2014), using a Bayesian hierarchical approach. Briefly, a set of manually recorded measurements from a subgroup of 143 F₁ trevally was used to generate the parameters used to predict weight (W) from length (L) using the length-weight relationship:

$$W = aL^b$$

where parameter b indicates growth in body proportions as the slope of a regression over log-transformed weight-at-length data, and a , the parameter describing body shape, as the intercept of a regression line over log-transformed weight-at-length data. The accuracy of the prediction model was estimated by calculating its R-square score. The net gain in each trait for each time point was calculated as the difference between the initial measurement in November 2017 and the measure of that month.

The phenotypic correlations between individual traits were measured using a Pearson's correlation matrix, which was constructed using all phenotypic measurements in Python version 2.7, using the Numpy library (McKinney, 2010).

2.3.7 Trait heritability and genetic correlations

Variance and covariance components were estimated using linear mixed animal models and restricted maximum likelihood methods with ASREML version 4.0 (Gilmour et al., 2015) in R. Narrow-sense heritability of each trait was estimated using a univariate analysis, modelled as follow:

$$y_i = \mu + a_i + e_i$$

where μ is the population mean, a_i is the breeding value and e_i is a residual term (Galwey, 2014). The heritability models were run separately for each time measure, with the target trait predicted using as fixed intercept effect.

The genetic covariances were estimated in a series of bivariate analyses. A bivariate model was fitted for each trait combinations to estimate genetic correlations and their standard errors, using the equation:

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \mathbf{Z}\mathbf{u} + \mathbf{e}$$

where $\mathbf{X}$ and $\mathbf{Z}$ are matrices and $\mathbf{y}$, $\mathbf{u}$ and $\mathbf{e}$ are vectors (Thompson et al., 1995).

2.3.8 Ethics statement

All research carried out in this study was approved by the animal ethics committee of Victoria University of Wellington, application number 25976. All data used in this study including the genome assembly, WGS and GBS sequencing libraries, phenotype data and supplemental material is deposited in an open data repository, which is accessible via www.genomics-aotearoa.org.nz/data.

2.4 Results

2.4.1 Filtering allowed to increase quality of calls tenfold in offspring

A total of 1.23 billion DNA sequence reads were generated for the F_0 generation, resulting in 13x coverage for each of the 13 individuals; a total of 3.05 billion reads were produced for all 12 pooled libraries of F_1 , with approximately 3 million sequence reads for each individual library, resulting in 0.42x genome coverage per individual fish (Supplementary Table 2.1). Quality was high across the full length of the reads for all F_1 plates, except towards the end of plate 5, where it was slightly reduced quality (Supplementary Figure 2. 1). Twenty-three F_1 sample extractions did not yield enough DNA for sequencing. A further 29 out of 1077 offspring libraries failed to be sequenced, probably due to low quality DNA.

The initial variant calling yielded 20.8 and 2.1 million markers for the F_0 and F_1 , respectively (Supplementary Figure 2. 2 A–D). After removing indels, 17.8 million sites were kept for the F_0 and 1.8 million for the F_1 . Forty-seven offspring were subsequently removed based on missing data (Supplementary Figure 2. 2 E, F). The average SNP read depths were 181.99 for the F_0 and 5,126.33 for the F_1 . After filtering for quality and depth,

17.7 and 1.1 million SNPs were kept for the F_0 and F_1 , respectively. The missing rate per SNP filtering resulted in 17.1 million and 214,700 sites for the parents and the offspring, respectively (Supplementary Figure 2. 2 G, H). The last filter for maximum depth in the parental call retained 16.9 million SNPs. These filtering steps resulted in a total of 171,923 SNPs shared between the F_0 and F_1 generations. Filtering enhanced largely the F_1 dataset, going from 68% missing data to 0.06%. The parental dataset went from 0.01% to 0% (Supplementary Table 2.1).

2.4.2 Reconstruction of the pedigree showed skewed parental contributions

A subset of 1,525 SNPs was used for the parentage assignment. Pedigree reconstruction allowed to determine sex for all F_0 individuals, which was further confirmed by biopsy. Visualisation of the pedigree showed that out of 13 sequenced F_0 , 10 individuals participated in the spawning (3 females and 7 males), generating 21 families (Figure 2.2 A). Both parents were identified for 63% (664) of the individuals in the F_1 population. The remaining F_1 had either one (31.4%) or both parents (3.6%) not genotyped. This is explained by mortalities that occurred in the F_0 broodstock before the start of this study. Genotyping information is therefore missing for some parents. The contributions were skewed among both sexes, but particularly so in females, with one of them contributing to up to 60.2% of the total F_1 population (Figure 2.2 B).

Figure 2.2: Molecular pedigree of the trevally population. (A) Circos representation of the pedigree structure of the population of trevally. 21 families were identified. Numbers in the outer in the outer ring indicate the total number of offspring per parent. Numbers in the ribbons indicate the number of offspring in the family. (B) Number of offspring produced by each parent in the F₀ generation divided into females (red) and males (blue).

2.4.3 Unbalanced inbreeding levels detected within families

The inbreeding F statistic (F_H) was calculated for 13 F_0 and 664 F_1 individuals with both parents known. The values ranged from a minimum of -0.28 to a maximum of 0.06 with a median of -0.06 for the F_0 and from -0.56 to 0.18 with a median of -0.06 for the F_1 . Variation in the inbreeding values did not significantly differ between the wild-caught F_0 and the F_1 generation (-0.08 to -0.06 respectively, p -value = 2.65^{-05}) (Figure 2.3 A). Most of per family average inbreeding values ranged from -0.02 to -0.11 except for families having female 8 as mother, which had higher average F_H values, ranging from 0.06 to 0.14 (Figure 2.3 B). The mean inbreeding values for fam11_5, 12_2, 12_4 and 12_5 were significantly lower (p -value < 0.01) than those of other families (Supplementary Table 2). Families with female 8 as a mother had a sample size too low ($n = 1$ to $n = 5$) to conclude if the means were statistically significantly different from those of the other groups.

Figure 2.3: Inbreeding scores for trevally individuals based on (A) generation, and (B) family. Visualised are the 1st, 2nd (median), and 3rd quartile and whiskers extending 1.5 times the interquartile range from the median (95% confidence interval). The coefficient of inbreeding used is Nei's F_{IS} and significant differences between groups are shown in Table S2. No significant differences (p -value > 0.01) were found between the F_0 and F_1 generations. Significant differences (p -value < 0.01) were found between different families.

2.4.4 Growth pattern in trevally is seasonally influenced and is under family-effects

For 2 years, phenotyping of the F_1 trevally cohort occurred roughly every 4 months, from the age of 2.1 to 4.1 years old. Ten growth traits that could either be directly extracted from the images (H25, H50, H75 and PL), indirectly measured as in the case of weight, or

calculated *a posteriori* as in the case of the net gains traits were recorded across eight time points. The *a* and *b* parameters used in the Bayesian hierarchical approach (Froese et al., 2014) were 0.02 and 2.99, respectively. They predicted weight with an accuracy of 0.92. The number of measurements per time point differed as a result of natural mortality and access to the individuals within the tanks. Between the first and second time points the most significant drop was observed from 1093 to 748 individuals, because of a high number of PIT tags being rejected from the gut cavity. The increase in sample size in the last measurement is explained by the fact that all individuals could be retrieved from the tank.

High variation was observed in all traits (Supplementary Figure 2.3). Between the first and the last measurements, standard deviation less than doubled for all traits but increased four-fold for EW and Δ EW (Supplementary Table 2.3). Coefficients of variation (CV) of height, length and weight traits remained constant throughout the experiment showing that variability in relation to the mean of the population was stable. The CV of net gain traits were inflated in the first measure (March 2018) and decreased over time (November 2019) (Supplementary Figure 2.4). Growth increase was not linear throughout the duration of the experiment. In all traits recorded, a seasonal pattern was detected (Figure 2.4): higher gains in all traits were concomitant with warmer ambient water temperatures. During summer (December to February), growth increased by an average of 27.3% across all traits (min: 22.2%, max: 31.1%), whereas during winter, growth only increased 7% on average (min: 2.3%, max: 12.3%).

When subdividing the F_1 by families, full-sibs with higher initial measures for traits H25, H50, H75, PL and EW remained higher throughout the experiment (Supplementary Figure 2.5 A - E). For instance, families 11_10 and 11_3 had the highest average measures of H25 in November 2017 with 61.28 mm and 64.00 mm respectively, and 106.88 mm and 110.83 mm respectively, in November 2019. However, gains in net growth traits (Δ H25, Δ H50, Δ H75, Δ PL and Δ EW) did not follow the same trend: compared with family 11_10 scoring the highest values for the measured traits across time, family 11_5 had a higher final net gain in all traits but Δ EW (Supplementary Figure 2.5 F - J).

Figure 2.4: Influence of seasonal temperature on the growth rate of trevally over two years for peduncle length (PL), height at 25% of PL (H25), height at 50% of PL (H50), height at 75% of PL (H75), estimated weight (EW), net gain in H25 ($\Delta H25$), net gain in H50 ($\Delta H50$), net gain in H75 ($\Delta H75$), net gain in PL (ΔPL), and net gain in EW (ΔEW). The area under the curve represents the relative percentage of gain measured in November 2017 (Nov17), March 2018 (Mar18), June 2018 (Jun18), October (Oct18) 2018, January 2019 (Jan19), April 2019 (Apr19), August 2019 (Aug19) and November 2019 (Nov19). The bottom band indicates the ambient temperature recorded in the tank.

2.4.5 Genetic correlations, phenotypic correlation and trait heritability

The estimates of heritability, variances, covariances, and phenotypic correlations between traits are reported in Supplementary Table 2.4. Based on Pearson's correlation coefficients, strong phenotypic correlations were observed between all height, length and weight traits throughout the experiment (0.85 ± 0.02 to 1.00 ± 0.00) and moderate to strong correlations were found for the net growth traits (0.50 ± 0.00 to 0.98 ± 0.01) (Figure 2.5).

Narrow-sense heritability was estimated for 10 phenotypic traits. Heritability estimates remained consistent throughout the experiment. The heritability range was moderate to high (0.67 ± 0.05 to 0.76 ± 0.06) for the measured traits (H25, H50, H75, PL and EW) and moderate (ranging from 0.28 ± 0.07 to 0.68 ± 0.07) for the net gain traits (Δ H25, Δ H50, Δ H75, Δ PL and Δ EW).

Strong positive genetic correlations were found between all measured height, length and weight traits throughout the experiment (close to unity, 0.94 ± 0.00 to 1.00 ± 0.00), and between all net gain traits (0.84 ± 0.07 to 1.00 ± 0.03), indicating that the alleles associated have a pleiotropic effect and influence multiple traits simultaneously. Correlation between measurements of height, length, weight and net gain traits were genetically moderate to strong (0.59 ± 0.10 to 0.99 ± 0.00).

The genetic variance of each trait increased over time, especially for length and weight traits (PL: 149.87 in November 17 to 524.09 in November 19; EW: 458.59 to 9178.11 in November 19; Δ PL: 98.70 in March 18 to 388.12 in November 19; Δ EW: 477.98 in March 18 to 7194.48 in November 18), showing that variation between individuals increases as fishes get bigger.

Figure 2.5: Heat maps of Pearson's phenotypic correlations (above the diagonal) and genetic correlations (below the diagonal) in (A) November 2017, (B) March 2018, (C) June 2018, (D) October 2018, (E) January 2019, (F) April 2019, (G) August 2019 and (H) November 2019, between height at 25% (H25), height at 50% (H50), height at 75% (H75) of the peduncle length (PL), estimated weight (EW), and net gain in height 25% ($\Delta H25$), height 50% ($\Delta H50$), height 75% ($\Delta H75$), peduncle length (ΔPL) and estimated weight (ΔEW). Positive correlations are displayed in blue and negative correlations in red colour. Colour intensity and the size of the circle are proportional to the correlation coefficients.

2.5 Discussion

Here, we applied for the first time a genomics-informed approach to study a captive trevally population. Data from different genotyping methods were pooled to reconstruct a two generation pedigree of 13 F₀ and 1100 F₁ and investigate the inbreeding levels in each generation and within families. Phenotyping data were added to assess the quantitative genetic architecture of 10 growth traits over 2 years. The results of this study will support breeding efforts in trevally, as well as inform more generally genomic work on other teleost species.

Marker-based pedigree reconstruction enabled to determine the sex of the broodstock and showed that most individuals contributed to F₁ offspring. However, reproductive contributions were skewed, particularly among females. Skewed contributions have been reported from a wide range of captive fish populations, including closely related species such as the yellowtail kingfish (Dettleff et al., 2020) and geographically co-occurring species such as the Australasian snapper (*Chrysophrys auratus*) (Ashton, et al., 2019). Because this study was carried out on F₁ obtained via hormone-induced spawning, different explanations could account for this biased distribution. First, physiological states of hCG-injected individuals (e.g. different stages of the reproductive cycle) may have influenced their ability to spawn, or limited gamete availability. Second, different survival rates of progeny could have skewed family representations (note: DNA sampling was conducted at 2.1 years of age). Third, courtship and mating behaviour might have been at play. This has been documented in yellowtail kingfish (Dettleff et al., 2020; Moran et al., 2007), where only one female and one male mated at any given time. Any of these factors, or a combination of them, could explain the skewed parental representation.

The initial level of genetic variation entering a breeding programme is set by the amount of heterozygosity of the founding population. It is crucial to monitor its subsequent loss in the next generations, as this will help to avoid inbreeding depression and a diversity bottleneck for a breeding programme. In our study, the wild F₀ parents represented the baseline for inbreeding statistics. No statistical differences were observed between the two generations, which were both slightly outbred (–0.08 and –0.06 on average for F₀ and F₁, respectively). Values were similar to the ones found in other wild-caught marine species such as orange clown fish (*Amphiprion percula*) (0.018) (Salles et al., 2016), Australasian

snapper (0.055) (Ashton, et al., 2019), or pacu fish (*Piaractus mesopotamicus*) (0.054 to 0.247) (del Pazo et al., 2021). Although the average inbreeding rates did not vary between generations, some slight differences could be observed between families in the F₁ generation.

During our 2-year study, we recorded 10 growth traits across eight time-points to describe the growth patterns of trevally. All fish were maintained under the same rearing conditions throughout the duration of the experiment. Despite this, some external parameters, such as food availability, might have varied slightly for some individuals because of the potential for hierarchical social behaviour in confined pools. A certain level of trait variability is evident in the data because not all individual fish could be extracted and sampled at each sampling point for logistical reason, meaning that for some data points, some very large or small fish were not included. We found that families which had initially higher values in growth traits remained larger throughout the experiment. However, some smaller families showed higher net gains compared with the larger families, demonstrating that the bigger fish were not necessarily the fastest growing proportionally. Finally, growth showed a pattern of being strongly influenced by seasonal temperatures. An increase in growth rate was observed during the warmer months of the year ($T^{\circ}\sim 21^{\circ}\text{C}$) compared with the colder months ($T^{\circ}\sim 11^{\circ}\text{C}$). This can be explained by an increase in metabolism during summer as water temperature rises and day length increases (Pauly, 1980). Similar results have been found in other species such as in chinook/king salmon (Perry et al., 2015) where the optimal growing temperature is 19.0°C ; or in yellowtail kingfish, where optimal rearing temperatures were found to be around 26.5°C (Abbink et al., 2012). However, too high temperatures can also have a negative effect on growth rates once it exceeds a level of tolerance, as shown in varied species of coral reef fishes (Munday et al., 2008).

Genomic-based pedigree allowed the estimation of narrow-sense heritability for measured growth traits, which were consistent over time and higher than those reported in other studies; for example, heritability estimates for growth traits (weight or length) were 0.26 (Whatmore et al., 2013) to 0.42 (Premachandra et al., 2017) in yellowtail kingfish; 0.30 to 0.34 in Asian seabass (*Lates calcarifer*) (Ye et al., 2017); 0.09 to 0.30 in Australasian snapper (Ashton et al., 2019); and 0.42 to 0.72 in Atlantic salmon (*Salmo salar*) (Thorland et al., 2020). These estimates are population specific and can be inflated if traits are also influenced by non-additive genetic effects (such as epistasis and dominance) (Visscher et al., 2008; Wray & Visscher, 2008). It is also to be noted that

estimates are often biased upwards due to difficulties to separate environmental and non-additive genetic effects common to full-sibs from additive genetic effects when full-sib families are reared in separate tanks until tagging (Kause et al., 2005). In this study, the high estimates observed in directly measured traits (H25, H50, H75, PL and EW) could be explained by the holding conditions of the fish. Although all fish were kept in the same conditions, early rearing effects (e.g. smaller tank until March 2018) might have been confounded with non-additive genetic variation. Heritability estimates of the net growth traits were more comparable with values previously reported for teleost species mentioned above (0.27 to 0.68). It is likely that correcting for the growth period that happened in a restricting environment, from hatch to November 2017, helped remove some of the early rearing effects influencing the heritability estimates on the main phenotypes. Individuals were substantially smaller in November 2017, their environment — a 13,000L tank where density was higher, oxygen levels likely lower, and competition might have occurred for space and food — could have had more of an impact on growth during juvenile stages (where there is less variance) compared with ~6 months later when growth had increased. Thus, the differences in growth observed in November 2017 could more likely be due to limiting environmental resources stunting their growth. From March 2018 onwards, the environment was less likely to be a limiting factor and the genetic component better explained the observed differences.

2.5.1 Future directions and management implications

This study represents the first in-depth genetic investigation of reproductive success and growth rates for a captive trevally population in New Zealand. Genome-wide marker sets combined with reference genomes will be useful for a wide range of future applications. Investigating the basic genetic structure of a founding population yields fundamental insights into the biology of a species, and is of primary importance when establishing a long-term breeding programme. Furthermore, understanding underlying genetic mechanisms of growth can support informed decisions about how to selectively breed species to fast track gains, while at the same time ensuring the long-term viability of the breeding programme as a whole (e.g. avoiding inbreeding).

In trevally, like in other teleost, seasonal effects have a significant influence on the realised growth rate. This finding has important implications for the selection of

aquaculture locations and monitoring of optimal rearing temperatures; holding fish in a warmer environment could increase the growth rate and reduce the time taken to reach harvesting size. Our study showed that families exhibiting larger measurements initially remained the largest individuals in the population throughout the duration of the experiment. Early removal of small individuals can thus be a useful hatchery-management method to maximise the realised growth potential overall. However, if there is no accompanying genetic management plan, it can result in a genetic bottleneck as often only a few families are selected, which increases the risk of inbreeding depression over several generations (Kincaid, 1983). The implementation of a genomic information will enable selection decisions to be made on earlier measurements (e.g. March 2018 measure) while confidently preserving genetic diversity in the programme. Early rearing effects are common in land-based facilities, where populations are kept in separate or small tanks, and this can influence the estimates of heritability of commercially important traits. We found that the inclusion of the net growth for each trait can improve the estimate of heritability, particularly if environmental differences had an impact on early performance. Focusing selection based on the net gain traits could help reduce the impact of environmental effect on the heritability estimates that would otherwise be difficult to separate out.

The high heritability estimates found in our study highlight that loci have a strong effect on the traits. Hence there is potential for making strong genetic improvements via selective breeding. Phenotypic and genotypic correlations between traits were all positive and moderate to strong. A breeding objective for increased fish length is thus expected to produce a positive response on the other growth traits measured in this study, provided that they have the same allometric relationship later in life. Future work on this species should focus on a single trait such as length, because it can be easily measured using high-throughput phenotyping methods. Taken together, our results suggest that trevally is a suitable future candidate for selective breeding for enhanced growth.

3 Unravelling the complex genetic basis of growth in New Zealand silver trevally (*Pseudocaranx georgianus*)

Authors

Noemie Valenza-Troubat, Sara Montanari, Peter Ritchie, Maren Wellenreuther

Author Contributions

Conceived and designed the experiments: NVT, PR, and MW; Data collection: NVT; Data analysis: NVT, SM; Wrote the primary manuscript: NVT; Prepared final manuscript: NVT, SM, PR, and MW.

Publication status

This chapter has been prepared for publication but will be published after submitting my thesis.

3.1 Abstract

Growth directly influences production rate and therefore is one of the most important and well-studied trait in animal breeding. However, understanding the genetic basis of growth has been hindered by its typically complex polygenic architecture. Here, we performed quantitative trait locus (QTL) mapping and genome-wide association studies (GWAS) for 10 growth traits that were observed over two years in 1100 F₁ captive-bred trevally (*Pseudocaranx georgianus*). We constructed the first high-density linkage map for trevally, which included 19,861 single nucleotide polymorphism (SNP) markers, and discovered eight significant QTLs associated with height, length and weight on linkage groups 3, 14 and 18. Using GWAS, we further identified 113 SNPs associated with 9 traits, uncovering 29 genetic hot spots involved in growth. Two of the markers found in the GWAS co-located with the QTLs previously mentioned, demonstrating that combining QTL mapping and GWAS represents a powerful approach for the identification and validation of loci controlling complex traits. This is the first study of its kind for trevally. Our findings provide important insights into the genetic architecture of growth in this species and supply a basis for fine mapping QTLs, marker-assisted selection, and further detailed functional analysis of the genes underlying growth in trevally.

3.2 Introduction

Growth facilitates essential functions such as reproduction or the ability to adapt to environments. Although there are exceptions, increase in body size is usually positively correlated with numerous fitness traits such as higher mating success and fecundity, increased offspring quality and lengthened longevity (Dmitriew, 2011; Gebhardt-Henrich & Richner, 1998). Most vertebrates exhibit a finite amount of growth. In fish, however, the process of growing is indeterminate, or indefinite, and continues throughout their life (Weatherley, 1972), although the rate tends to decline as body size increases (Pedersen & Jobling, 1989). This highly complex process is the result of interactions between environmental effects and genetic differences. Factors including sex (e.g. Deniel, 1990; Imsland & Jonassen, 2003), age (e.g. Deniel, 1990; Essington et al., 2011) or food availability (Jones, 1986) influence growth rate, as well as abiotic factors changing seasonally such as temperature (e.g. (Imsland et al., 2007; Karås & Klingsheim, 1997),

photoperiod (e.g. Imsland & Jonassen, 2003) or oxygen levels (Brett & Groves, 1979; Mallya, 2007). The genetic basis of growth traits is typically highly polygenic (De-Santis & Jerry, 2007; Wellenreuther & Hansson, 2016). Functional relationships between genetic variations and physiological parameters of growth have been described in commercially relevant species such as cod (*Gadus morhua*) (see review by Imsland & Jónsdóttir, 2002), Atlantic salmon (*Salmo salar*) (Tsai et al., 2015) or tilapia (*Tilapia mossambica*) (Liu et al., 2014), but are still poorly understood in many non-model species.

With the development of next generation sequencing technologies (NGS) the cost of high-density genotyping has drastically decreased, therefore enabling the more widespread application of quantitative trait locus (QTL) mapping experiments. Many variants associated with complex phenotypes have since been found in a variety of non-model species (Shi et al., 2020; Wang et al., 2019). However, QTL mapping approaches only allow the identification of genetic regions that are polymorphic between two parents, and relevant in a particular environment, therefore potentially missing some common variants associated with the traits. For this reason the location and effects of detected QTLs can vary between mapping populations. For example, in the Atlantic salmon, QTLs involved in growth have been found on different linkage groups according to different studies (Baranski et al., 2010; Besnier et al., 2020; Tsai et al., 2015).

Recently, genome-wide association studies (GWAS) have been widely employed to detect QTLs both in captive (Palaïokostas et al., 2018b) and natural populations (Santure & Garant, 2018). This technique also identifies associations between markers and phenotypes based on linkage disequilibrium (LD) (Meuwissen et al., 2001). Compared with QTL mapping, GWAS can be conducted on genetically diverse, unrelated individuals and is particularly advantageous when controlled crossing and generation of large segregating populations is difficult. GWAS has proved useful for the identification of loci associated with numerous growth traits in fish (Li et al., 2018; Yang et al., 2020). However, the power of a GWAS relies on the number of markers in relation to the extent of LD in the population (Newell et al., 2011). When applied to natural populations or collections of outcrossing individuals, characterised by a rapid LD decay, tens of thousands of markers are often required to obtain an adequate level of resolution. Because of the intrinsic limitations of statistical methods comparing thousands of tests, GWAS has a higher frequency of false discovery of loci than QTL mapping (Hayes, 2013). A final disadvantage of GWAS is its low power for detecting rare allelic variants. However, their frequency can be increased in

controlled crosses and therefore captured with QTL mapping. Hence, combining GWAS and QTL mapping often gives more complete and reliable results than using one of the methods alone (Fraslin et al., 2020; Su et al., 2018).

Aquaculture is the fastest growing food-production sector in Aotearoa New Zealand. Currently, the local industry relies almost exclusively on the farming of three species: GreenshellTM mussels (*Perna canaliculus*), Pacific oysters (*Crassostrea gigas*) and only one finfish, king/chinook salmon (*Oncorhynchus tshawytscha*), which is an introduced species (Camara & Symonds, 2014; Gentry et al., 2017). Hence, there is a strong interest in diversifying the range of species available for aquaculture. The native finfish trevally has been identified as a suitable candidate for New Zealand aquaculture; however, its rate of growth needs to be improved, which can be done using selective breeding. In this study, we investigated the genetic architecture of 10 growth traits in a new population of trevally. More specifically, we genotyped and phenotyped 1100 F₁ trevally to 1) generate a high-density linkage map, 2) detected rare QTLs associated with growth using linkage mapping in a sub-family comprising 89 individuals, 3) identified common SNPs associated with growth using a GWAS on the whole population, and 4) overlapped the results and annotated supported regions.

3.3 Methods

3.3.1 Study population

The trevally population used in this study was generated as part of a breeding programme started at the Plant and Food Research (PFR) finfish research facility in Nelson, New Zealand, in 2015. The full description of holding conditions and pedigree are described in Chapter 2. Briefly, the population comprised 13 wild caught F₀ broodstock and 1100 F₁ captive-bred offspring. Induced mass spawning of the F₀ generation in a single tank was used to produce the offspring F₁ generation. This resulted in a complex pedigree, including a combination of unrelated, full- and half-siblings in the F₁ generation. F₁ offspring were held in a single tank from hatch, in 2015, receiving the same feeding regime, light, water flow and aeration until the end of this experiment. The seawater tanks in the facility are located on the seaward side of Port Nelson and receive ambient seawater from an underground bore which is filtered using mesh filters and UV treatment.

3.3.2 Phenotyping and trait estimation

10 growth traits were used in the current study, namely peduncle length (PL), height at 25 (H25), 50 (H50) and 75% (H75) of the PL, estimated weight (EW) and related net gain traits (Δ PL, Δ H25, Δ H50, Δ H75 and Δ EW respectively). These measurements were recorded on three occasions throughout the experiment, at the beginning (November 2017), in the middle (October 2018), and at the end (November 2019), when the fishes were a little over two, three and four years old, respectively. Using the Morphometric SoftwareTM (<https://www.plantandfood.co.nz/page/morphometric-software-home/>), the outline of each fish was extracted from images and morphometric were located, and then used to make measurements. PL was measured by assessing the distance between the upper lip and narrowest cross-section of the tail. Height was measured at three positions along the fish: 25, 50, and 75% of the PL. The weight estimations (EW) were done following a Bayesian hierarchical approach as described in Froese et al. (2014). The net gain for each trait was calculated as the difference from the initial (November 2017) with subsequent measurements. The number of individuals measured differed between time points because of natural mortality that occurred during the study. The normality of the data was assessed visually using Quantile-Quantile (QQ) plots generated in the R statistical environment version 3.2.3 (R. Core Team, 2013).

3.3.3 Genotyping and variant calling

DNA extractions, genotyping and filtering were carried out as described in Chapter 2. Sequencing of the 13 F₀ bloodstock was done using Whole-Genome Sequencing (WGS) and genotyping-by-sequencing (GBS) was used for the 1100 F₁ offspring. However, for this analysis, both F₀ and F₁ were mapped simultaneously to the reference genome (Ruigrok et al., 2021) using the Burrows-Wheeler Aligner (BWA) version 0.7.17 (Li & Durbin, 2009) and variants were called jointly with the parallel module of freebayes version 1.3.1 (Garrison & Marth, 2012), with minimum of five observations and minimum mapping quality of 10. The SNPs with single-sample sequence coverage (sequencing depth) < 3 were removed to reduce the number of putatively erroneous genetic variants, and missing data and minor allele frequency (MAF) were set to < 20% and > 0.05, respectively.

3.3.4 Linkage map construction and QTL identification

The parents of each F_1 individual in the dataset were identified with Sequoia version 2.0.7 (Huisman, 2017) as reported in Chapter 2. The full SNP dataset (i.e. before the stringent filtering performed for the parentage analysis) was filtered for Mendelian errors ($> 5\%$), and checked for distorted segregation using a chi-square test with $\alpha=0.05$. The linkage map was constructed in LEPMAP version 3.0 (Rastas, 2017), using the largest family. Markers were separated into linkage groups (LG) with the SeparateChromosomes module (logarithm of odds (LOD) limit = 14, minimum markers per LG = 50). The order of the markers was computed with the OrderMarkers module. Single markers at the end of each LG were removed if they were more than 3 cM apart from the next closest marker. The accuracy of the final linkage map was investigated by plotting the marker genetic position (cM) versus their physical position (bp) on the available reference genome mega-scaffolds (number of mega-scaffolds = 24) for each LG (Marey plots, data not shown). MapChart version 2.32 (Voorrips, 2002) was used to visualise the genetic map. As the F_0 were assumed to be outbred, the linkage map and the genotypes of the mapping family were input as a 4-way cross in the R package R/qtl version 1.47-9 (Broman et al., 2003) for interval mapping. Standard interval mapping was performed and a genome-wide permutation test (Doerge & Churchill, 1996) with 1000 permutations was used to determine the LOD significance thresholds (p -value = 0.05).

3.3.5 Genome-Wide Association Study

GWAS was carried out on the entire genotyped F_1 population ($n=1100$). SNPs were removed if the call rate was smaller than 0.8, $MAF < 0.01$, if Mendel error rate $> 5\%$ (based on trios identified with the parentage analysis carried out above), and they were LD-pruned using an $r^2 > 0.80$ in a 50kb sliding window with five variants. Association analysis were performed using the Fixed and random model Circulating Probability Unification (FarmCPU) method (Liu et al., 2016) implemented in GAPIT3 version 3.1.0 (Wang & Zhang, 2020). A Bayesian information criterion (BIC)-based model selection was used to find the optimal number of principal components (PCs) for each time measure, to account for family and population structure. The cut-off for significant association was a False Discovery Rate (FDR) adjusted p -value = 0.05 (Benjamini & Hochberg, 1995), to control for multiple testing. To assess how well the model used in GWAS accounted for population

structure and family relatedness, results of the GWAS were visualised with QQ plots implemented in GAPIT3, which depicted the distribution of the actual p -values compared with the theoretical ones. Manhattan plots were used to visualise the SNPs associated with the different phenotypes, using the physical position of the markers on the reference genome.

3.3.6 Ethics statement

All research carried out in this study was reviewed and approved by the animal ethics committee of Victoria University of Wellington in New Zealand (Application number 25976).

3.4 Results

3.4.1 Phenotypic values of growth traits

All traits exhibited large levels of phenotypic variation, as discussed in Chapter 2. A summary of the mean values and standard deviations of the 10 growth phenotypes is shown in Table 3.1.

In both the family used for QTL mapping and in the entire population used for GWAS, a normal distribution of the residuals was observed for the ten traits investigated (Figure 3.1).

Table 3.1: Summary of the phenotypes height at 25%, 50% and 75% of the body (H25, H50 and H75 respectively), Peduncle Length (PL), estimated weight (EW) and net gains traits associated (Δ H25, Δ H50, Δ H74, Δ PL and Δ EW respectively) across the whole F_1 population and for the largest family. Included are the number offspring phenotyped (n), mean, and standard deviation (SD).

	All						Family					
	Nov 2017 (n=1,094)			Oct 18 (n=719)			Nov 2017 (n=87)			Oct 2018 (n=68)		
	Mean	SD		Mean	SD		Mean	SD		Mean	SD	
H25	49.71	9.02		66.94	13.51	90.1	52.21	7.95	73.71	11.32	96.52	12.64
H50	59.45	11.02		79.99	16.39	105.53	63.1	9.29	86.87	13.41	111.9	14.77
H75	48.58	9.91		61.26	12.69	85.63	52.85	8.99	66.72	9.75	92.36	12.39
LEN	156.79	23.92		200.64	35.36	264.09	167.59	21.32	218.19	30.04	280.78	33.07
EW	89.9	41.85		191.93	99.89	426.34	107.43	37.85	238.01	89.61	498.52	159.82
Δ H25				15.85	7.8	39.08			20.43	6.53	42.78	9.21
Δ H50				18.75	8.42	44.34			22.46	6.92	46.84	9.89
Δ H75				11.08	6.77	35.54			12.88	5.54	37.79	8.33
Δ PL				40.03	18	103.59			47.91	15.91	109.31	23.15
Δ EW				95.83	64.45	330.78			125.99	60.54	384.63	135

Figure 3.1: Distribution of the peduncle length measurements in November 2017 ($n = 1,094$), October 2018 ($n = 719$) and November 2019 ($n = 694$).

3.4.2 QTL mapping revealed rare variants associated with growth

13 F_0 and 1094 F_1 were successfully genotyped, and a common dataset of 171,923 SNPs was generated from variant calling and basic quality filtering. The largest family (including 87 offspring and 2 parents, $n=89$) was used to assemble the sex-averaged linkage map. A total of 21,665 SNPs were polymorphic in this family and passed the chi-square test, and 19,861 were successfully mapped to 24 LGs, and 1,830 after removal of the co-mapping loci (Figure 3.2).

Figure 3.2: Visualisation of the sex-averaged map built from the largest segregating family ($n = 89$). The 24 linkage groups represent the expected 24 *Pseudocaranx georgianus* chromosomes

LG numbers were randomly assigned, as no previous reference had been published. The genetic map spanned 1,335.46 cM, with an average marker distance of 0.73 cM. The largest gap was on LG14 and was 4.10 cM long. The longest and shorter LGs were 17 (82.89 cM) and 8 (48.14 cM), respectively (Table 3.2).

The genome-wide significant thresholds for QTL mapping ranged between 4.60 and 4.81 for the ten phenotypic traits (Table 3.3).

Table 3.2: Marker statistics of the linkage map constructed from the largest trevally family (n=89).

LG	Number of markers	Length (cM)	Average marker distance (cM) of individual linkage groups
1	85	61.27	0.72
2	80	58.41	0.73
3	81	57.26	0.71
4	68	49.34	0.73
5	81	56.69	0.70
6	65	51.72	0.80
7	87	60.09	0.69
8	70	48.14	0.69
9	78	54.43	0.70
10	84	57.22	0.68
11	82	61.88	0.75
12	71	49.30	0.69
13	79	59.00	0.75
14	64	53.51	0.84
15	67	52.84	0.79
16	67	56.27	0.84
17	108	82.89	0.77
18	82	54.95	0.67
19	80	53.82	0.67
20	71	55.04	0.78
21	65	52.83	0.81
22	73	49.29	0.68
23	63	45.39	0.72
24	79	53.88	0.68
Total	1830	1335.46	0.73

Table 3.3: List of significant Quantitative Trait Loci (QTL) detected for height at 25% and 75% of the peduncle length (H25 and H75 respectively), Peduncle Length (PL), Estimated Weight (EW) and net gain for H75 (Δ H75) and PL (Δ PL) in November 2017 (Nov17) and November 2019 (Nov2019), using a high-density linkage map.

Time	Traits	QTL ID	LOD threshold alpha=0.05	LOD	p-value	Significance ¹	LG ²	Peak position	PVE ³
Nov 2017	H25	trevally000114_4454829	4.71	4.73	0.05	*	3	23.2	22.14
	H75	trevally000114_23355470	4.64	5.5	0.01	**	3	52.13	25.26
	PL	trevally000114_4498123	4.71	4.83	0.04	*	3	24.9	22.55
	EW	trevally001200_2242455	4.71	4.88	0.04	*	18	0	22.78
Nov 2019		trevally000114_4498123	4.6	5.39	0.01	**	3	24.9	24.83
		trevally001200_2242455	4.6	4.82	0.03	*	18	0	22.53
	Δ H75	trevally001025_25180484	4.69	7.36	0	*****	14	17.59	39.71
	Δ PL	trevally001025_25205374	4.81	5.41	0.01	**	14	17.02	31.04

¹: Genome wide significance: * p < 0.05, ** p < 0.01, *** p < 0.005, **** p < 0.001

²: Linkage group

³: Percentage of Variance Explained, calculated as $PVE = 1 - 10^{(-2 \times LOD) / n}$

In total, eight significant QTLs were identified across three LGs (Figure 3.3), of which six were for traits recorded during the first measurement (November 2017), and two for the last measurement (November 2019).

Figure 3.3: Eight significant Quantitative Trait Loci (QTL) associated with growth traits were found in trevally, across three linkage groups (LG3, LG14 and LG18). In November 2017, two QTLs were found associated with peduncle length (PL_1_2017 and PL_2_2017, on LG3 and LG18 respectively), which were also associated with estimated weight (EW_1_2017 and EW_2_2017). In the same year, two more QTLs were found associated with height at 25% (H25_2017) and 75% (H75_2017) of the peduncle length, both on LG3. Finally, two QTLs were found in November 2019, one for net gain in peduncle length (ΔPL_2019) and one for net gain in height at 75% of the peduncle length (ΔH75_2019), both on LG14.

In particular, in the first measurement, one QTL was detected on LG3 for both H25 and H75, with an R^2 of 22.1 and 25.3, respectively, and two QTLs were found on LG3 and 18 for both PL and BW, explaining 22.6, 22.8, 24.8 and 22.5 of the phenotypic variation, respectively. In the last measurement two QTLs were found on LG14 for both Δ H75 and Δ PL, with R^2 of 39.7 and 31.1, respectively (Table 3.3). No QTLs were found in the middle measurement (October 2018).

3.4.3 Genome-wide association found strongly associated common SNPs

A total of 1,024 F_1 individuals had both genotypic and phenotypic data for all measurements. After filtering based on Mendelian errors, MAF and LD pruning, 107,067 SNPs were left and used in the GWAS. Model selection in GAPIT resulted in no PCs to be used as co-variates in any of the traits (Figure 3.4 A). QQ plot showed that FarmCPU adequately accounted for the confounding effects of family and population structure (Figure 3.4 B). A total of 113 SNPs were significantly associated ($-\log_{10}(p) > 7.03$) with at least one of the ten measured traits (Figure 3.4 C, Supplementary Table 3.1). Among these, four, seven and three SNPs were associated with more than one trait at the first, second and third measurement, respectively. No genetic regions were found to be associated with Δ H75 in October 2018 and November 2019. Significant associations were in some cases found for SNPs less than 0.5 Mb apart, highlighting hot spots on chromosomal regions. A total of 29 hot spots (containing two or more SNPs in a 2kb region) were identified. Of those hot spots, five regions contained 4 or more SNPs: 4 markers were located between 18.45 and 20.38 Mb on LG1, 4 others between 0.57 and 1.24 Mb on LG2; 8 markers were found between 0.05 and 1.80 Mb on LG5; 6 markers were located between 6.00 and 9.95 Mb on LG12; and 4 SNPs between 18.51 and 22.23 Mb on LG22 (Supplementary Table 3.1).

Figure 3.4: GWAS results for the net gain in Peduncle length (ΔPL) in November 2019. (A) Summary for Bayesian Information criterion (BIC) for the optimal number of Principal Components (PCs) to use in the model. (B) Quantile-Quantile (QQ)-plot of p -values. On the Y-axis is the observed negative base 10 logarithm of the p -values, and on the X-axis the expected observed negative base 10 logarithm of the p -values under the assumption that the p -values follow a uniform [0 to 1] distribution. The grey area shows the 95% confidence interval for the QQ-plot under the null hypothesis of no association between the SNP and the trait. (C) Manhattan plot of the results of the GWAS. On the X-axis is the physical position of the SNPs on the genome divided by chromosomes, and on the Y-axis is the negative log base 10 of the p -values. The horizontal green line represents the significance threshold.

3.5 Discussion

To the best of our knowledge, this is the first study that has identified the genetic basis of a quantitative trait in trevally, which represents the ground work for any further genetic study in this species. We constructed the first genetic map for trevally, which spanned 24 LGs and confirmed the 24 mega-scaffolds of the trevally reference genome (Ruigrok et al., 2021). A linkage map built for yellowtail kingfish (*Seriola lalandi*), the

closest sister species to trevally, was also resolved into 24 LGs (Nguyen et al., 2018), consistent with our findings.

Eight significant QTLs related to growth were mapped to three trevally linkage groups. Other studies in teleost have found multiple QTLs affecting growth such as in Atlantic salmon (Bentsbn, 1994; Besnier et al., 2020; Tsai et al., 2015), Australasian snapper (*Chrysophrys auratus*) (Ashton, Ritchie, et al., 2019), spotted sea bass (*Lateolabrax maculatus*) (Liu et al., 2020) or yellowtail kingfish (Nguyen et al., 2018), supporting the hypothesis of a polygenic regulation. No significant QTLs were identified with QTL mapping for the October 2018 traits. In the segregating family used for QTL mapping, the number of offspring decreased from 87 in 2017 to 68 in 2018 and 67 in 2019 as the result of natural mortality. Although some statistically significant QTLs were found in the measurements made in the latter period of the experiments, the smaller sample sizes in the two last measurements reduced the power of the study, which could explain the absence of QTLs detected in October 2018. Indeed, sample size is known to influence the power of a study to detect QTLs (Hong & Park, 2012), and is regularly discussed as one of the most important concerns when designing a mapping experiment (Ashton et al., 2017). Interestingly, two traits (Peduncle Length and Estimated Weight) shared the same QTLs on two LGs in the first measurement period (November 2017). This is consistent with the level of high genetic and phenotypic correlations reported for body length and body weight in Chapter 2. These findings suggest that selection applied on easily measurable traits such as length will result in the concomitant improvement of more difficult to assess, yet valuable, growth traits (such as weight) in the breeding of trevally.

GWAS identified 113 SNPs associated with growth, spanning 22 LGs, further supporting the hypothesis of a polygenic basis of growth-related traits in trevally. Growth is considered as a complex trait and has been found to be polygenic across the tree of life, in very diverse taxa from plants like in the model species *Arabidopsis thaliana* (Wieters et al., 2021), to vertebrates like humans (Sinnott-Armstrong et al., 2021), pigs (Zhang et al., 2019), and other fish species (e.g. Gutierrez et al., 2015; Li et al., 2018; Yang et al., 2020). In our study, a lower number of loci were found with the QTL mapping experiment than with the GWAS, which can be explained by the smaller amount of genetic variation represented in the single F₁ family compared with the overall breeding population, which was derived from thirteen parental individuals. Four QTLs and twenty markers with genome-wide significance were found to have a significant association with more than one

trait. This was expected, as the different phenotypes were all deduced from the Peduncle Length and they all then measured the same process underlying growth.

For each trait, the markers found in association were different from a year to another, both in QTL mapping and in GWAS. These differences could be due either to significant loci changing over time, by switching on or off, or to environmental variation, as it was observed in other QTL mapping studies that investigated traits highly affected by the environment (e.g. Ashton et al., 2019; Sun et al., 2017).

By comparing the relative physical positions of the SNPs, two regions found with QTL mapping appeared to be in close proximity with two significant SNPs identified by GWAS. In particular, the QTLs for H25, PL, and EW measured in 2017 spanned the 4.45-4.50 Mb region on chromosome 3 (corresponding to 23.20-24.90 cM on LG3) and SNP trevally000114_4956408, located at 4.96 Mb on chromosome 3, was found to be significant for PL in 2019 with GWAS; and the QTLs for $\Delta H75$ and ΔPL in 2019, encompassing the 25.18-25.21 Mb region on chromosome 14 (17.02-17.59 on LG14), are close to SNP trevally001025_25939455, located at 25.94 Mb on chromosome 14 and significant for $\Delta H50$ and ΔPL in 2019 (Figure 3.5, Table 3.3 and Supplementary Table 3.1). Being identified with two different statistical analyses and for multiple traits across two years, these two regions are then of particular interest for the understanding of the genetic determinism of growth in trevally. A BLAST search of the 100kb regions flanking those markers against the NCBI nucleotide database did not return any sequence similarities, indicating that they are located in non-coding or in non-annotated regions. Intergenic regions can still have a functional role in gene expression and regulation (Wyrick & Young, 2002). These SNPs could be located in an intron that acts as a regulatory region (promoter, enhancer, silencer or insulator).

Figure 3.5: Comparison of Quantitative Trait Loci (left) and SNP associated (right) with the net gain in peduncle length (ΔPL) in November 2019.

Compared with the heritability estimates found in Chapter 2, the present analysis still seems slightly underpowered. The heritability estimates range found was moderate to high (0.67 ± 0.05 to 0.76 ± 0.06) for the measured traits (H25, H50, H75, PL and EW), moderate (ranging from 0.28 ± 0.07 to 0.68 ± 0.07) for the net gain traits ($\Delta H25$, $\Delta H50$, $\Delta H75$, ΔPL and ΔEW) and remained consistent throughout the experiment. In the present study, the percentage of phenotypic variance explained (PVE) by the QTLs ranged from 22.1% to 47.36% and from 4.74% to 46.81% with GWAS, indicating that neither techniques were able to capture all of the genetic components of growth traits (Table 3.4).

Table 3.4: Comparison of heritability estimates found in Chapter 2 and the Percentage of Variance Explained (PVE) by Quantitative Trait Loci (QTL) found in interval mapping and markers from the Genome Wide Association Study (GWAS) in the present Chapter, in November 2017 (Nov17), October 2018 (Oct18) and November 2019 (Nov19).

Time	Traits	Heritability	PVE QTL	PVE GWAS
Nov 2017	H25	0.67±0.05	22.14	18.36
	H50	0.75±0.05		24.20
	H75	0.73±0.05	25.26	38.48
	PL	0.74±0.05	45.33	28.79
	EW	0.75±0.05	47.36	42.29
Oct 2018	H25	0.74±0.06		38.59
	H50	0.75±0.06		24.15
	H75	0.68±0.07		4.74
	PL	0.7±0.06		40.64
	EW	0.7±0.06		5.30
	ΔH25	0.46±0.08		10.94
	ΔH50	0.52±0.08		43.47
	ΔH75	0.28±0.07		
	ΔPL	0.47±0.08		46.81
Nov 2019	ΔEW	0.63±0.07		18.03
	H25	0.73±0.06		20.29
	H50	0.72±0.06		41.58
	H75	0.68±0.07		31.16
	PL	0.69±0.07		26.35
	EW	0.72±0.06		21.40
	ΔH25	0.56±0.08		16.13
	ΔH50	0.54±0.08		15.17
	ΔH75	0.43±0.08	39.71	
	ΔPL	0.51±0.08	31.04	21.74
	ΔEW	0.68±0.07		11.85

Ashton et al. (2017) showed that a moderate negative correlation ($r = -0.48$) was observed between sample size and PVE of QTLs, indicating that with more samples the PVE of QTLs declined. In this study, the PVE by the SNPs found in the GWAS were calculated by adding the PVE of each QTL. Additional genetic interactions, other than additive effects, should be considered. While heritability is a key feature of a trait indicating its potential for improvement via selection, polygenic traits are often influenced by non-additive genetic effects such as dominance or epistasis (Glover et al., 2017). Understanding the genetic mechanisms that underlie a trait is an important part of explaining phenotypic

diversity. This is particularly relevant when looking at traits related to fitness (e.g. growth, shyness, foraging or predator awareness) in populations that are undergoing domestication but still occasionally interbreed with wild conspecifics (e.g. when new broodstock is caught and added).

3.5.1 Future directions

In this study, the combination of QTL mapping and GWAS enabled the identification of genomic regions that control growth in a large captive trevally breeding population. From a farming perspective, parameters such as stocking density (Irwin et al., 1999) or feed availability and quality (e.g. Holm et al., 1990; Lee et al., 2018) can be carefully managed to accelerate fish growth in a land-based aquaculture facility. However, when aiming to develop new species for aquaculture, it is fundamental to understand the genetic architecture of commercially important traits that can potentially be enhanced through selective breeding. The findings of this study provide a useful framework for determining the genetic basis for growth traits in trevally. The identification of multiple QTLs through QTL mapping and genetic markers commonly involved in growth-related traits via GWAS, represents an essential step for the establishment of marker-assisted selection (MAS) breeding programme for trevally. Fine mapping, confirmation and annotation of relevant regions will bring a deeper understanding of the genetic architecture of growth. Specifically, further studies should aim to look at LD blocks to find possible associations with candidate genes.

4 Differential expression analyses reveal extensive transcriptional plasticity induced by temperature in New Zealand silver trevally (*Pseudocaranx georgianus*)

Authors

Noemie Valenza-Troubat, Marcus Davy, Roy Storey, Peter Ritchie, Maren Wellenreuther.

Author Contributions

Conceived and designed the experiments: NVT, PR, and MW; Data collection: NVT; Laboratory work: NVT; Data analysis: NVT, MD; Wrote the primary manuscript: NVT; Prepared final manuscript: NVT, PR, and MW; Advised data analysis: RS.

Publication status

This chapter has been prepared for publication, but will be published after submitting my thesis.

4.1 Abstract

Ectotherm species, such as marine fishes, depend on environmental temperature to regulate their vital functions. In finfish aquaculture production, being able to predict physiological responses in growth and other economic traits to temperature is crucial to address challenges inherent in the selection of grow out locations. This will become an even more significant issue under the various predicted future climate change scenarios. In this study, we used the marine teleost silver trevally (*Pseudocaranx georgianus*), a species currently being explored as a candidate for aquaculture in New Zealand, as a model to study plasticity in gene expression patterns and growth in response to different temperatures. Using a captive study population, temperature conditions were experimentally manipulated for 1 month to mimic seasonal extremes. Phenotypic differences in growth were measured in 400 individuals, and gene expression patterns of pituitary gland and liver were determined in a subset of 100 individuals. Results showed that growth increased 50% in the warmer compared with the colder condition, suggesting that temperature has a large impact on metabolic activities associated with growth. A total of 265,116,678 single-end RNA sequence reads were aligned to the trevally genome, and 28,416 transcript models were developed (27,887 of these had Genbank accessions, and 17,980 unique gene symbols). Further filtering reduced this set to 8,597 gene models. 39 and 238 differentially expressed genes (DEGs) were found in the pituitary gland and the liver, respectively ($|\log_2FC| > 0.26$, $p\text{-value} < 0.05$). Of these, 6 DEGs showed a common expression pattern between both tissues, all involved in housekeeping functions. Temperature-modulated growth responses were linked to major pathways affecting metabolism, cell regulation and signalling, previously shown to be important for temperature tolerance in other fish species. An interesting finding of this study was that genes linked to the reproductive system were up-regulated in both tissues in the high treatment, indicating the onset of sexual maturation. Few studies have investigated the thermal plasticity of the gene expression in the main organs of the somatotrophic axis simultaneously. Our findings indicate that trevally exhibit substantial growth differences and predictable plastic regulatory responses to different temperature conditions. We identified a set of genes that provide a list of candidates for further investigations for selective breeding objectives and how populations may adapt to increasing temperatures.

4.2 Introduction

Ectotherm species, such as most marine fishes, depend on temperature to maintain homeostasis, which makes them particularly sensitive to temperature fluctuations in their environment (Ficke et al., 2007). This is because the metabolic demand of these species is in part determined by the thermal conditions (Brett & Groves, 1979), which in turn influences physiological processes such as growth and reproduction (Fry, 1971; Gobler et al., 2018; Hochachka & Somero, 2002). Gene expression studies of the whole genome offer an opportunity to gain insights into the molecular machinery underlying these responses.

Like all organisms, teleost fish have adapted to thrive within a certain range of temperatures. These ranges vary between species and stages of life of the individual (Dahlke et al., 2020; Pörtner & Farrell, 2008). When exposed to temperatures outside of that range, individuals become stressed and suffer a reduction in performance. When temperatures rise above the optimal limit, metabolic costs rise (Sebens, 1987), and oxygen transport to tissues becomes limited (Ern et al., 2016). Below the optimal range, metabolic costs are lower but general activities, such as prey capture and digestion, decrease, which can negatively affect the ability of fishes to accumulate sustenance and energy. A lack of energy means a species is unable to sustain growth and other critical functions (Sebens, 1987). For example, cold exposure has been shown to decrease motility performances between 10% and 55% in juvenile silver perch (*Bidyanus bidyanus*) (Parisi et al., 2020).

In fish, growth is mainly regulated by the somatotropic axis (hypothalamus-pituitary-liver) (Reinecke, 2010). Some of the major hormones involved in this axis are somatostatin (Ss) and growth hormone releasing hormone (Ghrh), which are secreted by the hypothalamus. Growth hormone (Gh) is secreted by the pituitary gland (referred as pituitary hereafter) and binds to its receptor (Ghr) present at the surface of target organs such as the liver. This stimulates the release of insulin-like growth factor-I (Igf-I) (Fu et al., 2016; Norbeck et al., 2007) and activates the metabolism of fats, carbohydrates and proteins (Norbeck et al., 2007). Although the function of these hormones has been well studied in some model fish species (Duan et al., 2010), few studies have investigated the thermal plasticity of their gene expression in the main organs of the somatotropic axis simultaneously.

Phenotypic plasticity, or the capacity of a genotype to produce different phenotypes when exposed to different environments, enables organisms to adjust their physiology in

response to different thermal conditions. This coping mechanism can happen at different stages of life (Fu et al., 2019; Sandblom et al., 2014): developmental plasticity is often reversible whereas plastic responses later in life, often referred to as acclimation, can be reversible if exposure to stress is ended (Beaman et al., 2016). One of the critical intermediate steps of this mechanism is the induction of changes in gene expression levels (Hodgins-Davis & Townsend, 2009). This allows molecular and physiological pathways to be regulated to adjust to a new environment and continue to maintain a high level of fitness (Beaman et al., 2016). Studying expression plasticity thus provides insights into how different phenotypes are generated by the interactions between genotype and the environment (Huang & Agrawal, 2016). Moreover, the study of the expression levels of genes allows to examine plastic responses for a large set of traits (transcripts levels) that are relatively unbiased compared with traditional phenotypic traits, which can be skewed because of either preconceived notions of their importance or ease of measurement. Temperature-mediated expression changes have been observed in several fish species (Metzger & Schulte, 2018; Smith et al., 2013; Veilleux et al., 2015). Genetic variability has been showed to affect the plastic response within populations such as in the Australasian snapper (*Chrysophrys auratus*) (Wellenreuther et al., 2019) or in the three-spined stickleback (*Gasterosteus aculeatus*) (Metzger & Schulte, 2018), as well as between species (e.g. Haugen & Vøllestad, 2000; Jensen et al., 2008; Baumann & Conover, 2011; Hutchings, 2011). Consequently, each new species investigated has a potentially unique plastic response to temperature changes, which can itself vary among populations.

In this study, we used the New Zealand silver trevally as an aquaculture model to investigate transcriptomic responses to two temperature treatments. Specifically, using a replicated tank array, we experimentally measured temperature-dependent growth of 400 juvenile, sexually immature trevally over 1 month, and used high-throughput RNA-sequencing (RNA-seq) to compare differential gene expression responses in the pituitary and the liver in a subset of 100 individuals. The specific objectives of this study were to (1) measure growth responses following exposure to warm/cold temperatures; (2) quantify the expression changes of all genes in the pituitary and liver at different temperatures; and (3) annotate the differentially expressed genes in pituitary and liver to understand the potential molecular mechanisms regulating growth under different temperature.

4.3 Methods

4.3.1 Fish holding procedures, experimental setup and trait phenotyping

The New Zealand Institute for Plant and Food Research Limited (PFR) has been holding trevally since 2015, and in 2016, a selective breeding programme was started to select for enhanced growth (Valenza-Troubat et al., 2021). As part of this, an F₁ generation was produced in December 2018 at the Nelson Research Centre in New Zealand, in December 2018. A subset of this generation were randomly selected for this experiment. Prior to experimentation, the F₁ cohort was kept together in a 30,000L tank and provided with flow-through filtered seawater at ambient temperature and light conditions. At the start of the experiment, in May 2019, 5-month-old fish were anaesthetised (10 mg/L AQUI-S®, AQUIS NZ Ltd, Lower Hutt, New Zealand) and weighed. Four-hundred individuals were randomly assigned to a high (20°C) or a low (13°C) treatment (n = 200 per treatment), and divided into five replicate tanks per treatment (n = 40 per replicate). The tank allocation was randomised to avoid any spatial effects within the bay. Fish were acclimatised at ambient temperature (16°C ± 0.9) for a period of 10 days following their tank assignment. The temperature was then increased or decreased 1°C per day over a period of 5 days, until treatment temperatures of either 20.0°C or 13.0°C were reached, henceforth referred to as high and low temperature treatments, respectively. These temperatures mimic those found in the Tasman Sea during winter and summer. Temperature loggers (HOBO, Onset Computer Corporation, MA, USA) were used to monitor each tank and individual heaters were used to adjust the temperatures if needed. The low treatment had a mean temperature of 13.01°C and the high treatment had a mean temperature of 20.02°C, with minimal variation across the experiment (absolute maximum differences in the low and high treatments were ± 1.7°C and ± 0.5°C, respectively, Figure 4.1 A). The fish were maintained in 800 L tanks supplied with 1 µm filtered, UV filtered, recirculating aerated seawater (5 L/min flow), and fed daily with commercial pellets (3 mm, Nova ME; Skretting, Cambridge, Tas., Australia), at a ration equivalent to 2% body-mass, spread across four feeds per day. Dissolved oxygen levels were checked four times per day to ensure levels were kept at > 90% and tanks were cleaned every 3–4 days. The temperature conditions were then maintained for 32 days. Upon completion of the trial, a total of 100 fish (50 per treatment, 10 per replicate) were anaesthetised (25 ppm AQUI-S®) then netted from their tank and subsequently sacrificed using an overdose of anaesthetic (60 ppm AQUI-S®).

Immediately following euthanasia, fish were weighed again and dissected immediately to snap-freeze pituitary and liver tissues.

4.3.2 RNA extraction, Illumina library preparation and sequencing

Variations in temperature have been shown to differentially induce expression changes in different tissue types (Metzger & Schulte, 2018; Wellenreuther et al., 2019). We selected the pituitary as the endocrine gland of the somatotrophic axis, which regulates growth in teleost (Pérez-Sánchez et al., 2018) and the liver as its main target organ. Total RNA of both tissues were extracted separately for each individual using the NucleoSpin® RNA kit (Macherey-Nagel, Düren, Germany), in accordance with the manufacturer's instructions, which included an on-column DNase digestion. Due to size differences, total RNA was eluted in 40 µl of TE buffer for the pituitary samples and 50 µl for the liver samples. RNA integrity was checked by capillary electrophoresis with the Fragment Analyzer and the HS total RNA kit (Agilent Technologies, Palo Alto, CA, USA). Normalised starting quantities of total RNA (164 and 227 ng for ng for the pituitary and the liver, respectively) were used to prepare 200 Illumina sequencing libraries with the QuantSeq™ 3' mRNA-seq library kit plus Dual PCR add on extra barcodes (Lexogen, Vienna, Austria). Library preparations were performed as per the manufacturer's instructions. Sequencing was performed by the Australian Genome Research Facility (AGRF, Melbourne, Australia), 100 bp single-end reads on three lanes of NovaSeq (Illumina).

4.3.3 Genome annotation and completion rate

Assembly of the trevally genome is described in Ruigrok et al. (2021). Briefly, it was outsourced to DoveTail Genomics (Santa Cruz, California, USA), who provided a Meraculous (Chapman et al., 2011) *de novo* assembly from Chicago and Dovetail Hi-C data, by scaffolding first with the Chicago data plus HiRise. The resulting assembly was used as input into the HiRise pipeline along with the Dovetail Hi-C data. For this analysis, *de novo* repeats were identified using RepeatModeler version 1.0.11 (Smit et al., 2015) with the search NCBI engine. Repeats were classified by RepeatModeler into simple, tandem and interspersed repeats. A total of 12.8% of the genome was masked for repeats using

RepeatMasker version 4.0.5 (Smit, 2004). Automated gene models were predicted with the BRAKER2 pipeline version 2.1.0 (Hoff et al., 2018), using this experiment's trevally RNA sequences and the trevally genome as input. Gene completion levels were evaluated in the genome assembly using BUSCO version 3.0.2 (Simão et al., 2015; Waterhouse et al., 2018) with the lineage set vertebrata_odb9 to assess completeness of single-copy orthologues. E-utilities version 11.4 (<https://ftp.ncbi.nlm.nih.gov/entrez/entrezdirect>) was used to download protein sequences for zebrafish (*Danio rerio*) (88,504 sequences) and yellowtail kingfish (*Seriola lalandi*) (39,513 sequences) from NCBI (<https://www.ncbi.nlm.nih.gov/>) on 7 September 2020. BLASTX version 2.2.26 (Altschul et al., 1997) was used to search downloaded protein sequences using a translated nucleotide query for each transcriptome gene locus model. These results were merged with species specific Genome wide annotation for zebrafish provided in the package org.Dr.eG.db version 3.11.4 (Carlson, 2019), using Entrez stable gene identifiers (Maglott et al., 2010) and Genbank accessions to annotate BLASTX alignments of gene locus models. Common Gene Locus (gene model g1 ... g28416) from blast reports were also used to link transcripts to zebrafish, and yellowtail kingfish accession and description information. The Bioconductor (Huber et al., 2015) packages GO.db, and org.Dr.eG.db (Carlson, 2019) provided annotation maps describing the entire Gene Ontology (<http://geneontology.org>).

4.3.4 Sequencing data processing

The quality of the RNA-seq raw sequences from pituitary and liver tissues was checked using FastQC version 0.11.7 (Andrews, 2017) and MultiQC version 0.11.7 (Ewels et al., 2016). Reads were aligned to the trevally reference genome using the splice site aware aligner STAR version 2.6.1 (Dobin et al., 2013), with the parameter `--clip5pNbases` (clipping 12 bases from the 5' end of each mate) and the aligned reads were indexed and sorted using samtools version 1.7 (Li et al., 2009). The number of reads aligning to each transcript for each sample was counted using HTSeq version 0.9.1 (Anders et al., 2015). Reads were validated to belong to the correct tissue type by aligning them to NCBI downloaded sequences of genes unique to pituitary (Fsh- β and Lh) or liver (hemopexin b and apolipoprotein A-IV-like) using the Burrows-Wheeler Aligner (BWA) version 0.7.17, with the `bwa-mem` option.

4.3.5 Differential expression analysis

Filtering, normalisation and analysis were done using the Bioconductor package edgeR version 3.14.0 (Robinson et al., 2010). Duplicated genes and those with unknown annotation information were filtered out prior to normalisation. After graphically comparing the trimmed mean of M-values (TMM) (Robinson & Oshlack, 2010) with the TMM with singleton pairing (TMMwsp) normalisation methods (Supplementary Figure 4.1), the normalisation factors were estimated using TMM to adjust for different library sizes. This step did not require correction for individual gene lengths as QuantSeq library preparations preferentially target the 3' end of individual transcripts. Genewise dispersions were estimated by conditional maximum likelihood, conditioning on the total count for each gene (Smyth & Verbyla, 1996) utilising information between genes (Robinson & Smyth, 2007). Within a tissue, differential expression between temperature treatments was assessed for each gene using a Negative Binomial Generalized Linear Model with the function “glmTREAT” (McCarthy et al., 2012), which contains an in-built test for differential expression relative to a minimum required fold-change threshold. A gene was considered to be differentially expressed (DEG) between both treatments when the $|\log_2FC| > 0.26$ (e.g. 1.2 difference between treatments) and FDR-adjusted p -value < 0.05 (Benjamini & Hochberg, 1995).

4.3.6 Gene ontology

Gene Ontology (GO) analysis tested for over-representation of GO terms using the edgeR function goana (Young et al., 2010). The enrichment was ran on the DEGs with the parameter species = "Dr" mapping to the zebrafish GO terms. GO-terms with p -value < 0.01 were considered significant and were visualised with the online tool REVIGO (Supek et al., 2011). The same set of genes was also input into the Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa & Goto, 2000) using the edgeR function kegga, with the same parameter as mentioned above.

4.3.7 Ethics statement

All rearing, holding and sampling procedures were performed using standard hatchery practices in accordance with New Zealand's Animal Welfare Act (1999) and the Victoria University of Wellington code of ethics, application number 25976.

4.4 Results

4.4.1 Gain in weight increased in the high treatment

The average weights at the beginning of the experiment were 38.34 ± 7.73 and 37.31 ± 7.62 in the high and low treatment, respectively. At the end of the experiment, average weights were 62.28 ± 13.09 in the high and 41.74 ± 8.15 in the low treatments. While no significant differences were observed between the fish at the beginning of the experiment (t-test, p -value = 0.1813), growth was significantly different upon termination (t-test, p -value $< 2.2 \cdot 10^{-16}$), where fish increased 50% more in weight in the warmer compared with the colder condition (Figure 4.1 B).

4.4.2 Raw sequencing data and quality statistics

The three lanes of Illumina HiSeq2000 produced close to 1.22 billion single-end reads (Supplementary Table 4.1). On average, 60% of the reads were successfully aligned, leaving over 772.5 million reads for downstream analysis. The final number of reads per individual ranged from 1.33 million to 9.19 million (mean = 3.86 million $\pm$ 1.78 million). The number of reads aligned in each treatment group was balanced with 403.3 million in the high group (20°C) and 369.2 million in the low group (13°C).

Figure 4.1: Experimental temperatures and physiological data. (A) Temperature of high (red) and low (blue) treatments recorded in the replicate tanks throughout the experiment. (B) Average weights in the high (red line) and low (blue line) treatments recorded at the beginning and at the end of the experiment. Each dot represents the average weight in a replicate tank. The asterisk shows a statistically significant difference between means (p -value < 0.01).

4.4.3 Functional annotation and filtering

A total of 28,416 gene models were predicted using BRAKER. A summary of the number of transcripts kept after each filtering steps is given in (Table 4.1).

Table 4.1: Summary of filtering steps of the transcripts.

Step	Genes	
	Keep	Exclude
Before filtering	28,416	
GenBank accession known	27,887	529
Unique gene symbol	17,980	9907
Filter by expression	8597	9383
After filtering	8597	

27,887 meaningful unigenes (98.14% of the total gene models) were annotated against the genome-wide annotation for zebrafish databases using BLASTX. No threshold for the *E*-value cut-off was used so that all annotation matches were included. However, the exploratory plot (Supplementary Figure 4.2) illustrated that longer matches (with fewer mismatches) had smaller *E*-values, indicating that when *E*-value < 0.01, *p*-values and *E*-value are nearly identical. The highest correlation was observed between *E*-value and bitscore (vertical small *E*-value, with a low bitscore; large *E*-value not significant). A bimodal distribution (two clusters of reads) was observed when comparing the percentage of identity to bitscore, length and mismatch. Gene models seem to split in a mass of two matches, one of small length and another one of larger length. 17,980 gene models were left after filtering for duplicated symbols and 8597 genes from the expression set mapped to the filtered gene models after filtering for a minimum count of 0 (Supplementary Figure 4.3).

4.4.4 Differential gene expression analysis revealed extensive transcriptional plasticity

Differential gene expression patterns in pituitary and liver in response to elevated temperature were determined after 32 days. Multidimensional scaling of the samples showed that most of the variation between the samples was explained by the tissue type (Supplementary Figure 4.4 A), rather than the treatment (Supplementary Figure 4.4 B) (Note: the clustering was made using the 2000 first genes models of the 8597 gene model list). Normalisation of the data showed that the pituitary samples generally needed higher size factors than the liver samples (Supplementary Figure 4.5 A). Overall, the normalisation factors were normally distributed (Supplementary Figure 4.5 B) and more biological variation was observed within the pituitary samples (Supplementary Figure 4.6 A) compared with the liver samples (Supplementary Figure 4.6B). 39 DEGs were detected in the pituitary, accounting for 0.46% of the gene models (Figure 4.2 A, Supplementary Table 4.2), and 238 DEGs were found in the liver, accounting for 2.72% of the transcriptome (Figure 4.2 B, Supplementary Table 4.3).

In the pituitary, 15 and 24 DEGs were up- and down-regulated respectively, in the high versus low treatment. In the liver, 109 DEGs were up- and 129 DEGs were down-regulated (Figure 4.3 A). In both tissues, many of these DEGs corresponded to genes usually considered as “housekeeping genes”, i.e. expressed in most cells and involved in the maintenance of basal cellular functions. These included for instance genes coding for RNA splicing proteins, translation factors, or enzymes involved in metabolism. The growth hormone receptor gene *Ghrb* was found to be upregulated in the liver. Interestingly, genes involved in the production of the sexual hormones Fsh and Lh were up-regulated in the high treatment in the pituitary. In the liver, two genes coding for zona pellucida proteins (the acellular membrane surrounding the egg), were found to be down-regulated. Comparison of gene expression between treatments and tissue types identified five common down-regulated transcripts in pituitary and liver (Figure 4.3 B), which were *tet2* (promoting DNA demethylation); *iscub* (which codes for the scaffold protein involved in the *de novo* synthesis of iron-sulphur clusters within mitochondria); *thyn1* (involved in the induction of apoptosis); butyrophilin-like protein 2 gene (associated with the immune system and the major histocompatibility complex); and *gstr* (involved in detoxification processes). One gene was found up-regulated in both tissues, coding for a peptidylprolyl isomerase, which is involved the cellular response to oestrogen stimulus.

Figure 4.2: Differential expression of genes (DEG) between high and low treatment in the pituitary (A) and the liver (B). The MA plots show the average log of Count per Million (CPM) (X-axis) and log₂ Fold Changes (Y-axis). DEGs up- and down-regulated in the high versus low treatment are coloured in red and blue respectively (p -adjusted < 0.05), $|\log_2FC| > 0.26$. Those shown in black were unigenes that did not show significant expression.

Figure 4.3: Differential gene expression (DEG) within the tissues. (A) Barplot of the number of DEGs in each tissue between high and low temperature treatment. (B) Venn diagram showing the number of DEGs up- and down-regulated in the pituitary (PIT) and in the liver (LIV). There were only six DEGs shared between pituitary and liver, five of which were down-regulated, and one up-regulated.

4.4.5 Gene ontology to reveal molecular pathways

The most significantly enriched GO-term in each tissue depended on the temperature treatment: in the pituitary, genes involved in the female gonad development (GO:0008585), thyroid hormone metabolic process (GO:0042403) and the negative regulation of cellular component organisation (GO:0051129) were up-regulated (Figure 4.4 A, Supplementary Figure 4.7 A), whereas genes involved in myeloid cell homeostasis (GO:002262) were down-regulated (Figure 4.4 B, Supplementary Figure 4.7 B, Supplementary Table 4.4). In the liver, genes involved in translation (GO:0006412), protein activation cascade (GO:0072376) as well as cellular amide metabolic processes (GO:0043603) were up-regulated (Figure 4.4 C, Supplementary Figure 4.7 C), and genes involved in catabolic processes (GO:0009056) were down-regulated (Figure 4.4 D, Supplementary Figure 4.7 D, Supplementary Table 4.5).

The distribution of DEGs among KEGG terms further confirmed that in the pituitary, the most enriched up-regulated term was GnRh-signalling pathway (ko4912) and down-regulated was lysosome (ko04142) (Supplementary Table 4.6). In the liver, terms that were part of the ribosome (ko03010) were up-regulated, and terms that were part of the proteasome (ko03050) were down-regulated (Supplementary Table 4.7).

Figure 4.4: Network plots of Gene Ontology (GO) enriched terms for biological processes which were up (A) down (B) regulated in the pituitary and up (C) or down (D) regulated in the liver. Bubble colour indicates the user-provided p -value; bubble size indicates the frequency of the GO term in the underlying GOA database. Highly similar GO terms are linked by edges in the graph, where the line width indicates the degree of similarity. The initial placement of the nodes is determined by a ‘force-directed’ layout algorithm that aims to keep the more similar nodes closer together.

4.5 Discussion

Plastic responses to temperature are essential among ectothermic organisms, as all aspects of their physiology are directly dependent on their thermal environment. Therefore, understanding how thermal variation modulates phenotypic plasticity and its effect on major molecular pathways involved in the endocrine regulation of somatic growth is important. In the present study, transcriptomes of pituitary and liver tissues from 100 trevally were sequenced to investigate their physiological regulation mechanisms under different temperatures. Two-hundred RNA-seq libraries were constructed and sequenced, and 28,416 transcripts (N50 = 1,980 bp) were identified after assembly. A total of 17,980 genes with unique gene symbols were annotated successfully, which accounted for about 63.3% of all predicted gene models. To the best of our knowledge, this study represents the largest investigation of the thermal phenotypic plasticity of any carangid to date in terms of number of samples, and one of the largest for any teleost species. Our results show that somatic growth of trevally increased 50% more in the high (20°C) compared with the low (13°C) treatment. These findings are in agreement with other studies on other teleost. In the Carangid family specifically, optimal temperature ranges for metabolic rate have been found to range from 20–25°C for yellowtail kingfish (Brown et al., 2011; Pirozzi & Booth, 2009) and 22–26°C in greater amberjack fingerlings (*Seriola dumerili*) (Fernández-Montero et al., 2020; Fernández-Montero et al., 2018). Differences in growth rate show that temperature has an effect on the phenotype of trevally, which is in line with the seasonal influence described in Chapter 2.

We found that 39 and 238 genes were differentially regulated in the high versus low treatments in the pituitary and liver, respectively, demonstrating that temperature has an effect at the gene expression level. Many house-keeping genes (Watson, 1970), usually ubiquitously expressed across tissue type and developmental or cell cycle stage and required for the maintenance of functions essential for a cell's existence, were among the DEGs. Differential expression of housekeeping genes has been linked to stress (Torres-Contreras et al., 2018; Wang et al., 2013) and used as a marker for diseases (Byun et al., 2009). Important genes and pathways linked to the somatotrophic and reproductive axes were also highlighted. The growth hormone receptor gene (*Ghrb*), which is the transmembrane receptor for growth hormone, was found to be up-regulated in the liver. This class of receptor is a critical regulator of growth and metabolism in vertebrates,

however, growth hormone, its ligand, was not found to be significantly differentially expressed in the pituitary. Interestingly, the term growth (GO:0040007) was enriched in the pituitary in the low compared with the high treatment. Several hypotheses could explain this unexpected result: first, this pathway could need to be inactivated in order for growth to happen; second, at the time of the termination of the experiment, the fishes in the low treatment might have been over expressing genes involved in the growth pathway in order to counter-balance their slow metabolism. Our results also suggest that the expression of genes associated with female gonad development was increased in the pituitary in the high temperature treatment. Fsh and Lh are gonadotropin hormones synthesised in the pituitary and modulate gonadal steroids (Nett et al., 2002). Zona pellucida proteins (ZPs) were also found to be down-regulated in the liver. Studies have found ZPs to be lowly expressed in the liver of sexually mature fish like in the gilthead seabream (*Sparus aurata*) (Modig et al., 2006). This result highlights that either high temperatures onset early sexual maturation of immature juvenile fish, or that cold temperature delayed reproductive development in trevally. Overall, warmer temperature appeared to stimulate protein synthesis, cellular differentiation, lipid metabolism as well as regulate transcription of reproductive hormones. These processes are all important for myogenesis, somatic growth and sexual maturation in teleost. Colder temperatures had a positive effect on the expression of genes involved in blood cell differentiation and tissue injury, inflammation, and immunity (lysosome).

To the best of our knowledge, very few studies have simultaneously investigated the gene expression of both pituitary and liver, two of the main organs involved in the endocrine regulation of somatic growth. Two studies used comparative transcriptome analysis between pituitary and liver in teleost, of which only one investigated the effects of temperature on these two vital organs. Fu et al. (2019) looked at hypothalamo-pituitary (HP) and liver transcriptomes of bighead carp (*Hypophthalmichthys nobilis*) with high or low body weight and found genes enriched for the GnRH signalling pathway in the HP organs, which the KEGG analysis in our study also highlighted. GnRH is part of the reproductive feedback loop, where it is responsible for the release of Fsh and Lh by the pituitary. It indicated that fish in the high treatment have begun to mature sexually, which is a proxy for general sufficient growth as well as fitness. This implies that the juvenile immature fish grew to a sufficient stage to begin the process of sexual maturation in the warmer treatment. A study on golden pompano (*Trachinotus ovatus*), a species from the

same family as trevally, also compared the effects of different temperature treatments on pituitary and liver gene expression (Zhou et al. 2020). Interestingly, they found more genes to be differentially expressed in the pituitary than in the liver (458 and 205, respectively). Our study showed the inverse relationship, with only 39 DEGs in the pituitary and 238 DEGs in the liver, showing that metabolic processes were by far the largest in the liver, which we would expect given the important involvement of this tissue in the regulation of growth, as well as in other physiological functions. It is to be noted that the low treatment in the pompano study corresponded to the high treatment of this study (20°C), which could explain some of the differences observed. Similar to their results, we found enriched terms linked to transporter activity, enzyme activity, and transcription factor activity in the pituitary and immune regulation, digestion, and protein metabolism pathways in the liver.

Another interesting comparison to make is with other teleost that share a similar geographic distribution to trevally. Using a similar set-up, Wellenreuther et al. (2019) investigated the effects of akin temperature treatments on the gene expression of the Australasian snapper, another native species of New Zealand, also considered as a potential aquaculture species. They identified that temperature had a major effect on gene expression in both domesticated and wild strains of snapper, highlighting changes in protein synthesis and cellular multiplication. Our study also found enriched terms for ribosomal activity (GO:0005840) in the liver, and nucleosome (GO:0000786) and mitotic sister chromatid cohesion (GO:0007064) in the pituitary, which are involved in cell division. Interestingly, differences in growth between domesticated and wild snapper were found to occur through an interaction of the immune response and anabolic growth pathway modulation, highlighting that in the wild F₁ strain, immune-related activity was being prioritised over growth-related functions. Immune system responses were also found to be influenced by temperature in trevally; however, our findings show an enrichment in these pathways in the high compared with the low treatment. This could demonstrate that despite geographical similarities, these two species have differences in thermal performance curves. Interspecific adaptive differences can be caused by gene expression plasticity but mechanisms, including, epigenetic effects, developmental plasticity and neutral genetic variation should not be excluded (Schulte et al., 2011).

4.5.1 Future directions

In marine organisms, temperature is one of the most important environmental factors, and affects the majority of physiological activities, such as metabolism, growth, and development (Brierley & Kingsford, 2009; Seibel & Drazen, 2007). Our study shows that key responses to temperature stress depends on the activation of complex molecular networks involved in metabolic, immune and stress signal transduction pathways. We also highlighted the onset of sexual maturation in juvenile trevally. Our workflow demonstrates the utility of RNA-seq analysis for transcriptome profiling and shows that it can be used as a powerful strategy to assess a species' responses to environmental fluctuations. Given that the Intergovernmental Panel on Climate Change (IPCC) projects a global increase in water temperature, under all emission scenarios, of 1.5°C to 3°C by the end of this century (IPCC et al., 2018), the physiological constraints imposed on fishes are likely to cause significant challenge for aquaculture programmes worldwide. Climate change has already started influencing the intensity and frequency of extreme weather events globally (Pachauri et al., 2014) so predicting the fish's responses to ambient water fluctuations is becoming an increasing concern for sea-pen farming. The economic impact for aquaculture could be profound if plasticity mechanisms are not well understood. Uncontrolled phenotypic plasticity mechanisms have been shown to influence the genetic gain calculation in many livestock breeding programmes (Nguyen et al., 2017), and are an important parameter to assess for newly domesticated species (Zhang et al., 2016). Temperature fluctuations induce shifts in a variety of phenotypic traits, mostly related to reproduction and life history. Onset of early sexual maturation will be of prime importance for breeding programmes as it reduces growth rates due to the reallocation of energetic resources to reproductive organs (Gjerde et al., 1994; Oppedal et al., 1997). Moreover, organisms that are able to quickly adjust their physiology to environmental changes, i.e. exhibiting high phenotypic plasticity, may simply be better equipped to survive thermal fluctuations (Fu et al., 2016; Sandblom et al., 2014). Although all fish species exhibit unimodal thermal performance curves, each has its own specific parameters (i.e. thermal tolerance range, optimal growth temperature), and regulatory mechanisms can therefore vary. Selection for faster growth can offset the potential losses from climate change (Janssen et al., 2017). However, the genetic variability that differentiates plastic responses among families should also be harnessed to mitigate the effects of temperature through selection for shifted optimal temperature for growth (Sae-Lim et al., 2016). Adaptive potential found through standing

genetic and epigenetic variation, and newly derived mutations will be a way for selective breeding programmes to produce phenotypes that can compensate for growth losses in sub-optimal temperatures.

5 General Discussion

5.1 Overview

Within a species or even a population, there are numerous potential sources of phenotypic variation. Each of these sources signifies a distinctive underlying origin and decides whether a trait has an evolutionary potential, holds an ability to respond to natural/artificial selection, and whether it can respond to environmental variations. This thesis aimed to investigate the significance of both genetic and environmental factors in regulating the phenotypic expression of growth in the silver trevally *Pseudocaranx georgianus* (family: Carangidae). More specifically, this work investigated the genetic parameters (Chapter 2) and architecture (Chapter 3) of growth, as well as the proportion of variance explained by the environment (Chapter 2 and 4). The information collected provide a baseline for future studies about the selection potential for growth, and more broadly its genetic architecture and how trevally respond to different temperatures. This discussion chapter reviews the major findings and considers the future research directions as well as some potential applications of this research.

5.2 Summary of key findings

5.2.1 Data quality is of utmost importance

Filtering is a crucial step for any type of analysis in genomics. Depending on the sequencing technique used and the quality of the data, adjusting the filtering parameters is a fundamental step. Ensuring that erroneous data is removed (as best as possible) will provide higher quality, representative data for subsequent analyses. However, the diversity of techniques, questions and organisms studied means that there are no universal filtering guidelines or cut-off levels. Choosing filtering parameters therefore requires an in-depth understanding of their downstream effects, including what steps are required to reduce false positives and false negatives. In this thesis, different filtering criteria of the F_0 dataset did not have a large impact on the output (Chapter 2), because of the adequate sequencing depth that Whole Genome Sequencing (WGS) provided (coverage $\sim 30\times$). However, applying the same stringency thresholds of filtering used on the WGS data to the F_1 data resulted in a loss of information. When coverage (i.e. read depth per locus per individual) is insufficient, certain alleles may not be detected. For reduced representation approaches such as Restriction site Associated DNA (RAD) sequencing or Genotyping-By-Sequencing (GBS), filtering decisions can hence drastically influence the output. Consequently, F_0 and F_1 datasets were treated separately. The cut-off values for missing data in the GBS dataset were iteratively increased (separately per locus and per individual) to compensate for the much shallower sequencing depth. Low coverage was potentially caused by one or a combination of the following factors:

- Severe bias across enzyme cutting sites, i.e. the enzymes chosen were cutting too frequently.
- The multiplexing rate was too high. Sequencing a large number of samples (1100 F_1 individuals) brought statistical power to our study. However, a high level of multiplexing increases the amount of missing data (Torkamaneh et al., 2020). This is related to the fact that for any restriction site, only 30–50% of the restriction sites are actually sampled, although standard GBS does preferentially target sites more so than random tag GBS (Beissinger et al., 2013).

These issues highlight the need for planning when using reduced representation approaches. The choice of restriction enzyme(s) used, their cut frequency in the genome,

the level of multiplexing and the sequencing depth, are all important factors that will affect the quality of data downstream. WGS is now considered the best approach, especially as costs continue to decrease.

Although the initial filtering of the WGS and GBS datasets were the same in Chapter 2 and 3, the variants used in each chapter were called differently. This was the result of a long period of trial and error when running the interval mapping analysis presented in Chapter 3. Appendix B explains in detail the iteration process that led to the final result. Briefly, the initial significance thresholds computed after permutations in R/qtl were abnormally high. After several unsuccessful attempts that included checking the format conversion, marker distortion, and different parameters for R/qtl and LepMap, it appeared that the issue was not coming from either the map or the QTL mapping itself. An entirely new SNP variant call was made; however, the same issue occurred again, at least eliminating the possibility of an error at this step. After a few more attempts, the issue was traced back to a misunderstanding of the way LepMap encodes the genotypes in the map. Once these were recoded correctly, the issue was resolved. The rest of the analysis was conducted with the most recent iteration of the map, which had been obtained with the new SNP variant calls..

5.2.2 Reconstruction of the study pedigree and evaluating inbreeding level

One of the first results in this thesis involves the reconstruction of the trevally pedigree and the associated levels of inbreeding (Chapter 2). This type of analysis is particularly important for future breeding programmes, as it captures the population's structure, which any decision-making relies on. Prior to this work, the founding broodstock had been collected from the wild and the F₁ generation was produced by tank-based mass spawning. After hormone injection, all reproductively ready broodstock individuals had an opportunity to contribute to the breeding event. Pedigree reconstruction using the genotyping data identified parents for 63% of the F₁. It was discovered that reproductive contributions within the broodstock were highly uneven, particularly among females. This phenomenon has been reported in a wide range of captive fish populations, including in the closely related yellowtail kingfish (*Seriola lalandi*) (Dettleff et al., 2020) and in the geographically co-occurring species Australasian snapper (*Chrysophrys auratus*) (Ashton et al., 2019). Unequal contributions can lead to inbreeding and loss of genetic diversity. It

is therefore crucial to control inbreeding rates in subsequent generations. The initial level of genetic variation entering a breeding programme is set by the amount of heterozygosity of the founding population. In this study, the 13 wild F_0 parents represented the baseline for inbreeding statistics. Although the average inbreeding levels did not vary between generations, some slight differences could be observed between families in the F_1 generation. The number of breeding individuals in a population (i.e. effective population size – N_e) has a significant impact on the maintenance of genetic diversity and on the level of inbreeding depression over time in small populations (Frankham et al., 2014; Jamieson & Allendorf, 2012). Studies have suggested that to limit inbreeding depression to 10% over five generations, a $N_e > 100$ individuals is required for wild populations (Frankham et al., 2014). These numbers could be applied to the present trevally population as it is an outbreeding diploid species with uncontrolled spawning. A more detailed discussion about broodstock management and further recommendations for the trevally breeding programme are discussed in the “Future research and applications” section of this chapter.

5.2.3 Trait heritability is a key determinant in selective breeding programmes

Heritability provides a broad view of the relative importance of genetic and environmental factors in producing a specific trait (Visscher et al., 2008). In its narrow sense, it is defined as the proportion of variation in a trait attributable to additive genetic variance (Visscher et al., 2008). It is typically one of the most important statistics generated when investigating the genetic basis of a trait, as it directly informs on the potential to achieve selection gains. Since growth rate was identified as one of the key limiting factors of the aquaculture potential for trevally (which is currently lower than those reported in already cultivated sister species such as the yellowtail kingfish *Seriola lalandi* and the California yellowtail *Seriola dorsalis*), we calculated the heritability estimates of 10 traits related to growth (Chapter 2). All estimates were high from the first measurement, which is encouraging: the most noticeable selection of individuals for the breeding programme will most likely occur during their first year of life (e.g. from 100,000 to 4000 individuals). Our results indicate higher values than those reported in other studies. For example, heritability estimates for growth traits (weight or length) were 0.26 (Whatmore et al., 2013) to 0.42 (Premachandra et al., 2017) in yellowtail kingfish; 0.30 to 0.34 in Asian seabass (*Lates calcarifer*) (Ye et al., 2017) and 0.42 to 0.72 in Atlantic salmon (*Salmo salar*)

(Thorland et al., 2020). This is likely caused by two factors: first, the study population was kept in the same conditions from hatching, essentially fixing the environmental variance component as constant and giving our study increased power to detect the additive genetic variance participating in the traits. Second, it could also be that the estimates are inflated because of some other fixed effects that this study was not able to capture, such as variance from the day of spawning. Since heritability estimates are population specific, we would expect them to slightly lower if a second cohort was added to this analysis. Third, estimates can also be inflated if traits are under the influenced of non-additive genetic effects (Visscher et al., 2008; Wray & Visscher, 2008), which we investigated in Chapter 3.

5.2.4 Linkage maps and QTL mapping allow to understand the genetic architecture of traits

The trevally linkage map was a new and important genetic resource to investigate the architecture of growth (Chapter 3). It is one of the densest linkage maps reported to date in carangids. A total of 19,861 SNPs were arranged into 24 linkage groups (LGs), which is akin to the number of chromosomes for this species. A linkage map built for yellowtail kingfish, was also resolved to 24 LGs (Nguyen et al., 2018), consistent with our finding. Fine-scale linkage maps are among the key resources for genetic research, acting as road-maps for the genome. Although high-quality genome assemblies are now becoming the new “gold standard” for QTL and candidate gene applications (Braasch et al., 2015), linkage maps constitute the primary genome mapping tool for non-model species in the absence of complete genomes. In this thesis, the linkage map was used to inform the reference genome for trevally, arrange SNP markers in the QTL analysis and find potential candidate gene positions using integrated marker and genome scaffold information.

Interval mapping revealed eight significant QTLs related to growth, localised on three linkage groups. Interestingly, two traits (Peduncle Length and Estimated Weight) shared the same QTLs on two LGs in the first measurement period (November 2017), which is consistent with the level of high phenotypic correlation reported for body length and body weight in Chapter 2. However, the small number of individuals in the mapping population ($n = 89$) significantly reduced the power of detection of QTLs, only capturing rare variants involved in the traits. Sample size is known to influence the power to detect QTLs (Hong & Park, 2012), and is regularly discussed as one of the most important

concerns when designing a mapping experiment (Ashton et al., 2017). In breeding programmes, QTL studies are often performed on relatively small sample sizes due to financial constraints. Because of these limitations, many studies (including this one) still detect few QTLs with moderate to large effect sizes (e.g. Liu et al., 2019; Zhu et al., 2019). This could corroborate the idea that some quantitative traits have a relatively simple genetic architecture. However, a large-scale simulation experiment performed by Beavis (1994) showed that when the sample size is small (e.g. $n < 100$), the statistical power of detecting a small QTL is as low as 3% and the estimated effects are typically inflated 10-fold. This phenomenon is now known as the “Beavis effect”: the number of individuals needed to map small effect QTLs increases as the ratio of the average difference in phenotype between marker allele genotypes (δ) on the phenotypic standard deviation of the trait within marker genotype classes (σ_W) decreases and as allele frequencies depart from 0.5 (Figure 5.1).

Figure 5.1: Numbers of individuals (log10 scale) required to detect Quantitative Trait Loci for a range of effect sizes in association mapping populations in which the minor allele frequency is 0.5 (blue), 0.25 (red) and 0.1 (green), modified from Mackay et al. (2009).

Moreover, closely linked QTLs with effects in the same direction tend to give the appearance of a single large effect QTL (Mackay et al., 2009). Putative effects are generally detected only at the peak (the point within the region where linkage is strongest), even though, in some cases, QTLs may lie within the identified region but not at the exact point of the peak (Crosses, 2001). In a similar way, closely linked QTLs with opposite effects tended to be missed, as there are too few recombinants. That is, not only do studies tend to report only estimates that are statistically significant, but, within regions, only the largest of all significant results are reported (Allison et al., 2002). Historically, in animal breeding, linkage analysis mapped QTLs to large intervals of 20 cM or more (Georges et al., 1995; Van Laere et al., 2003), which is too long to identify the underlying causal mutations. Goddard & Hayes (2009) estimated that no less than 300,000 SNPs would be needed for studies across cattle breeds as markers need to be spaced less than 10 kb apart to show consistent linkage disequilibrium (LD) phase.

A crucial parameter in this type of analysis is the percentage of phenotypic variance explained (PVE) by a SNP (Kent, 1983). This parameter combines the allele frequency with the mean difference between the SNP genotypes. An approximate idea of the number of individuals needed for a mapping study can be easily calculated and reduced, if necessary, by using animals that have been progeny tested so that the mean of their progeny can be used instead of their own phenotypic value (Goddard & Hayes, 2009). This is advantageous as long as the mean phenotype of the progeny is more highly correlated with the animal's breeding value (BV) than is its own phenotype. Although limited correlation was found between the number of genetic markers and QTLs reported in studies (Ashton et al., 2017), higher marker densities are necessary to identify multiple closely linked regions that were probably individually too small to have been detected otherwise. Albeit using a high-density linkage map, the number of QTLs found through interval mapping in our study was still low and the PVE high, which reflects a power issue due to the number of individuals being a limiting factor.

5.2.5 Applying GWAS analyses to fine map QTLs

Using QTL mapping, rare variants were detected only within the largest family of the trevally population. In addition, a GWAS was used to detect common alleles using the overall F₁ population, and thus increasing the total sample size. Results showed that 113

SNPs spread over 22 chromosomes were linked to growth, which is more in accordance with the hypothesis of a polygenic basis of growth-related traits. Compared with the heritability estimates found in Chapter 2, the results in Chapter 3 still seem slightly underpowered, which indicates that neither techniques were able to capture all the genetic variance involved in the traits. As the number of individuals in a study population increases, the PVE for identified SNPs decreases, as it is observed in Chapter 3 between GWAS in comparison with QTL mapping. This pattern was first described in Fisher's infinitesimal theory (1930), suggesting that traits are determined by an infinite number of unlinked and non-epistatic loci. With the assumption of normality and linearity, each locus has an infinitesimal effect. Using a random mutation model, Fisher showed that very small mutations have a 50% chance of being favoured by selection, but that this probability falls rapidly for progressively larger-effect mutations (Figure 5.2, blue). Yet, a number of "powerful" studies, with high sample sizes, are still finding genomic regions for quantitative traits explaining a substantial fraction of phenotypic variance, which cannot be explained by the Beavis effect. Another interpretation of these empirical results is the exponential model developed by Orr (1998). Orr's model represents the distribution of factors during an adaptive walk, and uses a simple exponential form. At the beginning of the adaptive "walk" towards the optimum, a lot of evolutionary space is available, in which any mutations, even those with large effects, can have a chance to be favoured by selection. Later, when the organism is close to its optimum, the space is reduced so successfully fixed mutations must have smaller effects (Figure 5.2, green). In this model, quantitative variation is determined by a small number of QTLs of relatively large effect followed by an increasing number of genes with progressively smaller effects. Hence, strong QTLs are expected to have a higher chance early on during the adaptive walk (e.g. early in the selection process).

The PVE by the SNPs found with GWAS were calculated by adding the PVE of each individual marker and ranged from 4.74% to 46.81% per trait. The missing percentages indicate that they are not just additive effects, but also gene-by-gene interactions to be considered. A more detailed discussion about additional potential investigations on non-additive genetic effects is provided in the "Future research and applications" section of Chapter 5.

Figure 5.2: Representation of a two-dimensional Fisher–Orr adaptation model. Each dimension represents the fitness conferred by a quantitative trait; mutations are pleiotropic as they simultaneously influence both traits. The present phenotypic position f for the population is shown by the outer black circle, the red central point represents the optimal phenotype and the grey dashed lines show intermediate states. Green arrows represent random mutations of different sizes and illustrate Fisher’s view that large mutations are usually disadvantageous; the largest mutation would be disadvantageous whatever the angle of displacement. Blue arrows show an idealised adaptation process under Orr’s model with four sequential mutations and show how early mutations (1 and 2) tend to have larger effects than later mutations (3 and 4). Source: Farrall (2004).

5.2.6 Growth is strongly influenced by temperature

The description of growth patterns and their seasonal variation in Chapter 2 provides important insights for the breeding programme. During 2 years of data collection, 10 different traits related to growth were recorded, across eight time-points. A steep increase in growth rate was observed during the warmer months of the year ($T^{\circ}\sim 21^{\circ}\text{C}$) compared

with the colder months ($T^{\circ}\sim 11^{\circ}\text{C}$), strongly hinting at temperature-related phenotypic plasticity. The pedigree reconstruction highlighted an interesting additional insight: although full-sibs with higher initial directly measured traits remained higher throughout the experiment, net growth traits did not follow the same trend. For example, compared with family 11_10, which scored the highest values for the directly measured traits, family 11_5 had a higher final net gain in all traits but one. This points to a potential $G \times E$ interaction, where different genotypes respond differently to the same environmental fluctuation.

To further investigate the involvement of V_E in the plastic growth of trevally, we analysed the gene expression of the pituitary and liver exposed to summer- and winter-like temperatures in Chapter 4. These two organs are part of the somatotrophic axis and have major roles in the physiological regulation of growth. It was found that after 1 month, somatic growth had increased 50% more in the high (20°C) compared with the low (13°C) treatment. These findings corroborate the results from Chapter 2 and are in agreement with other studies on teleost. Particularly, in the carangid family where species showed high optimal temperature ranges for metabolism, e.g. $20\text{--}25^{\circ}\text{C}$ for yellowtail kingfish (Brown et al., 2011; Pirozzi & Booth, 2009) and $22\text{--}26^{\circ}\text{C}$ in greater amberjack fingerlings (*Seriola dumerili*) (Fernández-Montero et al., 2018). Genes influenced by temperature included the growth hormone receptor gene (Ghr). This class of transmembrane receptor is a critical regulator of growth and metabolism. However, its ligand, Gh, was not found to be significantly differentially expressed in the pituitary. Other DEGs up-regulated were Fsh and Lh, coding for the follicle-stimulating and the luteinizing hormone, respectively. Both of these gonadotrophins act in synergy in the reproductive pathway (Schally et al., 1971). Sexual maturity is usually triggered once a sufficient size or age is attained, making it a good proxy for growth and fitness in general (Chattopadhyay & Chattoraj, 2017; Devlin & Nagahama, 2002; Thorpe, 1994). Not only do the results of Chapter 4 confirm that temperature regulates somatic growth through the somatotrophic axis, but they also highlight pleiotropic interactions between the genes involved in both growth and reproduction.

5.3 Future research and recommendations

5.3.1 The complexity of decision making and resulting biases

Recent advances in genomic technologies have been tremendous. More sequencing options as well as an increase in the number of bioinformatics tools available mean that there are more opportunities to tailor experimental designs and analyses to the particular biological questions. Each experiment is unique because of its aim, the study species, the sequencing technique used, the packages and software selected for the analysis, and in the selection of their parameters. The growing number of options has, however, also increased the potential for bias and error, and made comparisons among different studies more challenging. Each choice influences the result obtained downstream. As seen in Chapter 2, GBS introduces bias through shallow depth of sequencing. Furthermore, different variant callers yield distinct results, and more or less stringent parameters will change the set of SNPs used, among other things. Given the sheer amount of data used in experiments, genomics now requires comprehensive skills in bioinformatics and, consequently, a deep and full understanding of the tools used and their key parameters. During the course of this thesis research, large amounts of time were spent formatting files, reading manuals and deciphering obscure parameter nomenclatures, rather than analysing actual data. Software and programmes are unforgiving mentors, but ultimately, I am grateful for those hours. Once the analytical frameworks I used became clear and I familiarised myself with the jargon of each new tool, I learnt some of my most useful coding skills, which in turn gave me a deeper understanding of the theory applied through them. Most of the pitfalls I encountered can be avoided by systematically:

- Subsetting large datasets and rigorously testing various parameters to plot and compare their outputs.
- Simulating data for which we have prior knowledge of the ‘truth’, and choose software or parameters giving results closest to the ‘truth’. It cannot be stress enough how simulating data should be standard practice.

The way forward will be to develop standardised, automated and robust pipelines. Although much more efficient in the long term, this task is initially time-consuming. Unfortunately, building workflows is generally not recognised by the wider scientific community, which still mostly values a researcher through their publications. GitHub has

been a brilliant platform for version control and more generally for sharing code. The resources found on the open source community are central to the success of contemporary research projects. Nowadays, most studies include half a dozen researchers working together. Biologists are collaborating with bioinformaticians and routinely using artificial intelligence and neural networks to cope with the ever-growing quantity of data as only an integrative, holistic view can hope to understand the immense intricacy of biological systems. We can simply not reduce life and its diversity to standalone fields anymore. A multidisciplinary approach will be the way forward to piece together biological systems in their wider context.

5.3.2 Aquaculture breeding programmes

One of the primary goals of this thesis was to generate genomic resources and information for the trevally breeding programme. Although these results are highly informative, incorporating them into the decision making is almost as complicated as collecting them. Many factors must be considered and each one of them will have complex interactions affecting outcomes.

For instance, the results of this research indicated that trevally have distorted parental contributions, which may be a consequence of uncontrolled tank-based spawning or subsequent family-specific mortality. A cohort mating strategy could be used to increase the number of parents contributing and even out the mating differences. Tank-based spawning using genotype data could help select less closely related individuals when constructing the broodstock for the next generation. This approach, also called walk-back selection, can be mixed with additional steps that reduce contribution distortions, such as holding the broodstock in multiple tanks and standardising the volumes of fertilized eggs from each tank. However, these techniques introduce environmental variation, which would need to be accounted for in downstream analyses. If walk-back selection were to be implemented, further research would also be needed to determine its effectiveness in controlling inbreeding over multiple generations. It was mentioned earlier that simulating genomic data is important to test bioinformatics tools with regards to the specific experimental design and type of data dealt with. In the same manner, simulations should be constructed with the specific configurations of the current facility's capacities, the biological limitations of the species, the target selective breeding gains, the target

inbreeding limits, and the relevant research such as the results from this thesis provide. This will allow to model genetic diversity versus selective pressures, compute manual versus tank-based spawning and to determine if the current broodstock contains sufficient genetic diversity to balance selective pressure in future generations.

The results from Chapters 2 and 3 could be implemented in the breeding programme by attributing BVs to future genotyped individuals to predict their likelihood of showing a certain phenotype. This predicted information could enable better, earlier identification of growth traits, which would lead to better time management in hatcheries and save costs. The greater the BV, the greater the genetic merit of an individual will be for a given trait. Across a population, this translates to a bell curve of the distribution of BVs (Figure 5.3).

Figure 5.3: Distribution of breeding values across a population. The families with the best breeding values are shown in the yellow area.

Although long-term gains can be significant, this approach requires a high initial investment: fish handling, fin clipping of a high number of individuals, as well as extensive phenotyping. This last point is essential as BVs have been shown to be more accurate as the number of traits recorded increases (Crouch & Bodmer, 2020).

This leads use to another key decision for any breeding programme: which traits to record, how many and what tools to use. While the results of this thesis are informative and relevant, future research could make significant advances by developing and implementing high-throughput phenomic methods. Phenomics is the full set of phenotypes of an individual and is seen as the counterpart to genomics (Houle et al., 2010). Many genetic studies (including this thesis) still use relatively simplistic methods; for example, a small number of simple phenotypes recorded during discrete phenotyping events. One of the primary limiting factors for phenomic methods is that they are often significantly more complicated to put into place. Phenotypic data collection technology is often not capable of collecting sufficient volumes of data. Unlike genomics, which expanded with the development of high-throughput sequencing, phenotype data collection is often still manually collected. The development of high-throughput phenotyping technology will be central to more widespread incorporation of phenomic approaches into genomic studies.

Ultimately, deciding to use genomic instead of phenotypic selection for a breeding programme should be informed by results such as those gathered in this thesis, the literature, and according to the capacities of the facility and staff availability. These questions require ongoing discussions between researchers, facility managers and support staff to be able to put together a realistic plan for implementing such decisions.

5.3.3 Enhancing our understanding of the genetic architecture of growth

The combination of GWAS and QTL mapping used in Chapter 3 enabled the identification of common and rare genomic regions that control growth in trevally. This molecular information not only has utility in developing trevally as a new species for aquaculture, but also in more fundamental research. To the best of our knowledge, this is the first study that has identified the genetic basis of a quantitative trait and investigated non-additive genetic variation in trevally. It represents the ground work that will allow any further genomic investigations on this species. We acknowledge that parts of this work can still be enhanced. For instance:

- Increase the number of individuals used, which will allow:
 - the detection of loci of very small effect that might still have been undetected;

▸ to refine the effect of the size of the QTLs detected which were likely inflated because of LD between QTLs (whereby there might be more than one gene in that same region or multiple loci in one region).

- Better control of environmental and sampling variances.
- Increase the density of the linkage map or use a reference genome to increase the maximum number of detectable QTLs, which is set by the size of genetic maps.

The regions flanking the markers found in Chapter 3 did not correspond to known genes, indicating that they are most likely located in non-coding regions. This is still informative of the genetic architecture as it suggests that these loci could be located in intergenic regions affecting the expression of not just one gene but multiple genes. Moreover, the findings in Chapter 4 suggested that pleiotropy also plays a role in the genetic architecture of growth. Further investigation of these interactions will provide a more complete understanding of the genetic networks involved. Specifically, we recommend using:

- Using family based design and fitting general and generalised linear mixed models to investigate the effects of epistasis,;
- Using multivariate models to dissect pleiotropic QTL from closely linked QTL;
- Identifying further pleiotropic interaction among molecular networks through gene co-expression networks.

Genes (except their regulatory sequences) do not directly participate in molecular interactions. However, the potential interaction patterns between these networks are encoded in genomic DNA, and inherited from generation to generation. Additive and non-additive genetic interactions also combine with various types of molecular interactions between gene products, or between gene products and gene regulatory elements. Furthermore, both genes and gene products are subject to change by environmental factors and developmental histories (Figure 5.4). Linking loci to genes, gene to molecular networks, and investigating how these interconnected, interacting, layers are influenced by the environment is therefore only possible if datasets are combined. Interpreting results in an integrated manner will allow a more holistic and complete understanding of macro-phenotypes.

Figure 5.4: Integrating datasets. Genetic networks encode molecular networks, which in turn determine phenotypes. All these layers respond to and interact with the environment (modified from Han, 2008).

5.3.4 Climate change and implications for trevally

Genomics tools improve our understanding of how aquatic organisms adapt and respond to the environment, and also improve our ability to monitor environmental variation and exploited species. Using RNA-seq analysis as a strategy to assess transcriptional responses to environmental conditions, we studied the effect of different temperature regimes on the plasticity of growth (Chapter 4). Our study showed that key responses to thermal stress depend on the activation of complex molecular networks involved in metabolic, reproductive, immune and stress signal transduction pathways and the expression of specific sets of related genes. Given that the Intergovernmental Panel on Climate Change projects a global increase in water temperature, under all emission scenarios, of 1.5°C to 3°C by the end of this century (IPCC et al., 2018, Figure 5.5), the physiological constraints imposed on fishes are likely to cause significant limitations for aquaculture programmes worldwide.

Figure 5.5: The Intergovernmental Panel on Climate Change's projections under high (red) and low (blue) emissions. Any individual year may look quite different. Source: IPCC, 2018.

At an intraspecific level, the genetic variability that differentiates the plastic response among families can be harnessed to mitigate the effects of temperature through selection for shifted optimal temperature for growth (Sae-Lim et al., 2016). In Chapter 2, we found that seasonal fluctuations in temperature induced different responses across different genotypes (or families), which highlights a $G \times E$ interaction ($V_{G \times E}$). Since $G \times E$ interactions can result in re-rankings in a breeding programme, an interesting future experiment would be to rerun the temperature challenge of Chapter 4, but instead of treating the fish as a single population as was done in this research (i.e. a single genotype), separate the fish by family, to partition V_E from $V_{G \times E}$, and investigate the differences in plastic response mechanisms. Understanding $G \times E$ interactions will allow to select genotypes that will thrive better in different environments. Breeding programmes could potentially breed different strains acclimated to different temperature conditions to insure resilience in the face of climate change.

Findings from this thesis are also relevant to wild trevally populations. From a conservation and fisheries management perspective, climate change is shifting the geographic distribution and abundance of wild fish populations (Azzurro et al., 2019; Last

et al., 2011). Research has shown that the longevity of the trevally fishery around New Zealand could be compromised by temperature shifts (Brooks, 2020). Trevally populations are predicted to decline from waters around the North Island by 2050 where the largest stocks are currently located, and be mostly located in the South Island 2100 (Figure 5.6).

Figure 5.6: Recorded modern distribution of trevally (A) and probability distribution projections for in 2050 (B), and 2100 (C). Colours reflect probability of presence (POP) estimates between 0 and 1. Darker green indicates high POP, while red indicates low POP (modified from Brooks, 2020).

Southward extensions or shifts of populations have been shown to be mainly driven by sea surface temperature (Brooks, 2020). Again, integrating information and datasets will allow to widen the scope of our results. Information on temperature tolerance in trevally could be used to inform ecological models (Wilson et al., 2010) and predict the response of the species to further shifts in temperature. It could also bring insight into the complexities of and constraints on evolutionary dynamics under realistic ecological conditions (Ellegren & Sheldon, 2008; Wolak & Reid, 2017).

5.4 Conclusions

This study represents the first in-depth genomic investigation of growth rates for a captive trevally population in New Zealand. The genome-wide marker datasets combined with the reference genome will be useful for a wide range of future applied and fundamental research. Investigating the basic genetic structure of a founding population yields fundamental insights into the biology of a species, and is of primary importance when establishing a long-term breeding programme. Furthermore, understanding the underlying genetic mechanisms of growth can support decisions about how to selectively breed this species to fast-track gains, while at the same time ensuring the long-term viability of the breeding programme (e.g. avoiding inbreeding). Finally, the analysis of molecular networks will help conservation efforts by predicting the response of trevally to further temperature changes.

A Supplementary material

A.2 Chapter 2

Supplementary Figure 2. 1: FastQC quality scores across the 100 bp reads for each of the twelve pooled sequencing plates of offspring.

Supplementary Figure 2. 2: Filtering effects on the whole-genome resequenced F_0 (green) and Genotyped-by-Sequencing F_1 (yellow) data sets. Missing data per individual (indv) are plotted before and after filtering for the F_0 (1 and 5) and the F_1 (2 and 6) respectively; missing data per locus are plotted for unfiltered and filtered data sets for the F_0 (3 and 7) and the F_1 (4 and 8). The red dashed lines indicate the mean proportion of missing data.

Supplementary Figure 2.3: Distributions of the mean values for height at 25% of the peduncle length (H25), B) height at 50% of the peduncle length (H50), C) height at 75% of the peduncle length (H75), D) peduncle length (PL), E) estimated weight (EW), F) net gain in H25 ($\Delta H25$), G) net gain in H50 ($\Delta H50$), H) net gain in H75 ($\Delta H75$), I) net gain in PL (ΔPL), J) net gain in EW (ΔEW) of captive silver trevally. The 25th, 50th and 75th percentiles are represented by horizontal lines and the 5th and 95th percentiles are represented by error bars. Dots represent outliers.

Supplementary Figure 2.4: Distribution of the coefficient of variation of height at 25% of the peduncle length (H25), B) height at 50% of the peduncle length (H50), C) height at 75% of the peduncle length (H75), D) peduncle length (PL), E) estimated weight (EW), F) net gain in H25 ($\Delta H25$), G) net gain in H50 ($\Delta H50$), H) net gain in H75 ($\Delta H75$), I) net gain in PL (ΔPL), J) net gain in EW (ΔEW) of captive silver trevally. The 25th, 50th and 75th percentiles are represented by horizontal lines and the 5th and 95th percentiles are represented by error bars. Dots represent outliers.

Supplementary Figure 2.5: Growth rates per family for 10 morphometric traits A) height at 25% of the peduncle length (H25), B) height at 50% of the peduncle length (H50), C) height at 75% of the peduncle length (H75), D) peduncle length (PL), E) estimated weight (EW), F) net gain in H25 ($\Delta H25$), G) net gain in H50 ($\Delta H50$), H) net gain in H75 ($\Delta H75$), I) net gain in PL (ΔPL), J) net gain in EW (ΔEW)

Supplementary Table 2.1: Sequencing metrics and filtering effects on the whole-genome resequenced F_0 and Genotyped-by-Sequencing F_1 datasets.

	F_0	F_1
Read length	125	100
Sequence reads (billion)	1.23	3.05
Coverage	13	0.42
Failed samples	0	29
Raw calling yields (millions of SNPs)	20.8	2.1
Indel filtering (millions of SNPs)	17.8	1.8
Removed individuals with missing data	0	47
Average read depth	181.99	5,126.33
Depth filtering (millions of SNPs)	17.7	1.1
Filter missing SNPs (millions of SNPs)	17.1	0.2
Maximum depth filter (millions of SNPs)	16.9	0.2
Total shared SNPs	171,923	

Supplementary Table 2. 2: Comparison of the differences of the mean F statistic (F_{IS}) between the F1 families. Upper triangle shows the p -values adjusted for multiple comparison using a post hoc Tukey correction. The diagonal shows the means inbreeding value for each family. The bottom line shows the number (n) of offspring in the family.

	11_10	11_2	11_3	11_4	11_5	11_6	11_9	12_10	12_2	12_3	12_4	12_5	12_6	12_9	8_10	8_2	8_3	8_4	8_5	8_6	8_9
11_10	-0.05	1.00	1.00	1.00	0.91	0.98	0.98	1.00	0.95	1.00	0.20	0.31	0.98	1.00	0.22	<.0001	0.52	0.29	0.34	0.01	0.19
11_2		-0.04	1.00	1.00	0.97	1.00	1.00	1.00	0.98	1.00	0.56	0.66	0.99	1.00	0.62	<.0001	0.74	0.70	0.68	0.06	0.49
11_3			-0.03	1.00	<.0001	1.00	1.00	0.16	0.00	0.03	<.0001	<.0001	0.00	1.00	0.59	<.0001	0.80	0.69	0.71	0.04	0.50
11_4				-0.04	<.0001	0.97	0.98	0.60	0.00	0.15	<.0001	<.0001	0.00	1.00	0.25	<.0001	0.61	0.32	0.41	0.01	0.23
11_5					-0.09	<.0001	<.0001	0.44	1.00	1.00	0.80	0.96	1.00	0.07	0.00	<.0001	0.10	0.00	0.01	<.0001	0.00
11_6						-0.02	1.00	0.03	<.0001	0.00	<.0001	<.0001	<.0001	0.93	0.78	<.0001	0.88	0.86	0.85	0.09	0.66
11_9							-0.02	0.09	0.00	0.02	<.0001	<.0001	0.00	0.94	0.87	<.0001	0.92	0.92	0.90	0.13	0.75
12_10								-0.07	0.69	1.00	0.00	0.01	0.92	1.00	0.03	<.0001	0.29	0.05	0.10	0.00	0.04
12_2									-0.09	1.00	0.66	0.90	1.00	0.13	0.00	<.0001	0.11	0.00	0.02	<.0001	0.01
12_3										-0.08	0.29	0.53	1.00	0.83	0.01	<.0001	0.19	0.02	0.04	0.00	0.02
12_4											-0.11	1.00	0.63	0.00	<.0001	<.0001	0.03	0.00	0.00	<.0001	0.00
12_5												-0.11	0.87	0.00	0.00	<.0001	0.04	0.00	0.00	<.0001	0.00
12_6													-0.08	0.29	0.00	<.0001	0.13	0.01	0.02	<.0001	0.01
12_9														-0.05	0.19	<.0001	0.54	0.26	0.34	0.01	0.18
8_10															0.05	0.91	1.00	1.00	1.00	1.00	1.00
8_2																0.13	1.00	0.87	1.00	1.00	1.00
8_3																	0.09	1.00	1.00	1.00	1.00
8_4																		0.05	1.00	1.00	1.00
8_5																			0.07	1.00	1.00
8_6																				0.11	1.00
8_9																					0.08
n	11	5	31	64	89	28	18	74	80	42	60	55	59	29	3	5	1	3	2	2	2

Supplementary Table 2.3: Count, mean, standard deviation (Stdv), and coefficient of variation (CV) for each continuous trait for in November 2017 (Nov17), March 2018 (Mar18), June 2018 (Jun18), October 2018 (Oct18), January 2019 (Jan19), April 2019 (Apr19), August 2019 (Aug19) and November 2019 (Nov19).

Time measure	Trait	Samples	Mean	Stdv	CV
17-Nov	H25 (mm)	1093	49.71	9.03	0.18
	H50 (mm)	1093	59.45	11.02	0.19
	H75 (mm)	1093	48.58	9.91	0.20
	PL (mm)	1093	156.79	23.93	0.15
	EW (g)	1093	89.90	41.87	0.47
18-Mar	H25 (mm)	748	57.01	11.79	0.21
	H50 (mm)	748	66.95	14.81	0.22
	H75 (mm)	748	54.08	13.25	0.24
	PL (mm)	748	172.64	31.98	0.19
	EW (g)	748	123.64	68.88	0.56
	Δ H25	743	6.06	6.07	1.00
	Δ H50	743	5.87	6.85	1.17
	Δ H75	743	4.01	6.91	1.72
	Δ PL	743	12.55	14.00	1.12
	Δ EW	743	28.44	34.41	1.21
18-Jun	H25 (mm)	742	62.09	12.85	0.21
	H50 (mm)	742	74.12	15.74	0.21
	H75 (mm)	742	59.22	13.33	0.23
	PL (mm)	742	186.81	34.32	0.18
	EW (g)	742	156.22	85.26	0.55
	Δ H25	737	10.99	7.14	0.65
	Δ H50	737	12.84	7.80	0.61
	Δ H75	737	9.00	7.25	0.81
	Δ PL	737	26.18	16.81	0.64
	Δ EW	737	60.15	50.73	0.84
18-Oct	H25 (mm)	719	66.94	13.52	0.20
	H50 (mm)	719	79.99	16.40	0.21
	H75 (mm)	719	61.26	12.70	0.21
	PL (mm)	719	200.64	35.38	0.18
	EW (g)	719	191.93	99.96	0.52
	Δ H25	715	15.85	7.80	0.49
	Δ H50	715	18.75	8.43	0.45
	Δ H75	715	11.08	6.78	0.61
	Δ PL	715	40.03	18.02	0.45
	Δ EW	715	95.83	64.50	0.67
19-Jan	H25 (mm)	704	77.03	14.74	0.19
	H50 (mm)	704	91.21	17.68	0.19
	H75 (mm)	704	69.50	14.07	0.20
	PL (mm)	704	228.87	38.95	0.17

	EW (g)	704	282.36	135.04	0.48
	$\Delta H25$	699	26.05	9.51	0.36
	$\Delta H50$	699	30.05	10.62	0.35
	$\Delta H75$	699	19.43	8.20	0.42
	ΔPL	699	68.34	23.48	0.34
	ΔEW	699	186.65	101.57	0.54
19-Apr	H25 (mm)	698	84.29	15.35	0.18
	H50 (mm)	698	98.29	17.85	0.18
	H75 (mm)	698	75.85	14.32	0.19
	PL (mm)	698	245.27	38.80	0.16
	EW (g)	698	343.40	150.48	0.44
	$\Delta H25$	693	33.25	10.64	0.32
	$\Delta H50$	693	37.03	11.50	0.31
	$\Delta H75$	693	25.67	8.93	0.35
	ΔPL	693	84.58	24.82	0.29
	ΔEW	693	247.52	118.42	0.48
19-Aug	H25 (mm)	630	89.24	15.42	0.17
	H50 (mm)	630	103.84	17.63	0.17
	H75 (mm)	630	82.65	14.62	0.18
	PL (mm)	630	256.25	38.77	0.15
	EW (g)	630	389.10	163.00	0.42
	$\Delta H25$	627	38.20	10.76	0.28
	$\Delta H50$	627	42.67	11.33	0.27
	$\Delta H75$	627	32.57	9.20	0.28
	ΔPL	627	95.81	25.45	0.27
	ΔEW	627	293.68	132.22	0.45
19-Nov	H25 (mm)	694	90.10	15.71	0.17
	H50 (mm)	694	105.53	18.67	0.18
	H75 (mm)	694	85.63	15.81	0.18
	PL (mm)	694	264.09	40.47	0.15
	EW (g)	694	426.34	178.06	0.42
	$\Delta H25$	689	39.08	11.22	0.29
	$\Delta H50$	689	44.34	12.37	0.28
	$\Delta H75$	689	35.54	10.20	0.29
	ΔPL	689	103.59	27.33	0.26
	ΔEW	689	330.78	147.23	0.45

Supplementary Table 2.4: Heritability estimates (in bold), Pearson's phenotypic (above the diagonal) and genetic correlations (below the diagonal) in November 2017 (Nov17), March 2018 (Mar18), June 2018 (Jun18), October 2018 (Oct18), January 2019 (Jan19), April 2019 (Apr19), August 2019 (Aug19) and November 2019 (Nov19). Means $\pm$ standard deviations (SD) are indicated for height at 25% (H25), height at 50% (H50), height at 75% (H75), peduncle length (PL), estimated weight (EW), delta height 25% (Δ H25), delta height 50% (Δ H50), delta height 75% (Δ H75), delta peduncle length (Δ PL) and delta estimated weight (Δ EW). The last row in italic reports the additive genetic variance (vA) of each trait.

17- Nov	H25	H50	H75	PL	EW					
H25	0.67$\pm$ 0.05	0.92 $\pm$ 0.01	0.85 $\pm$ 0.02	0.89 $\pm$ 0.04	0.88 $\pm$ 0.07					
H50	1.00 $\pm$ 0.00	0.75$\pm$ 0.05	0.95 $\pm$ 0.01	0.96 $\pm$ 0.02	0.95 $\pm$ 0.04					
H75	0.98 $\pm$ 0.01	0.99 $\pm$ 0.00	0.73$\pm$ 0.05	0.93 $\pm$ 0.03	0.91 $\pm$ 0.05					
PL	1.00 $\pm$ 0.00	0.98 $\pm$ 0.01	0.94 $\pm$ 0.01	0.74$\pm$ 0.05	0.99 $\pm$ 0.01					
EW	0.99 $\pm$ 0.01	0.97 $\pm$ 0.01	0.94 $\pm$ 0.00	1.00 $\pm$ 0.00	0.75$\pm$ 0.05					
vA	25.89	29.23	24.8	149.8 7	458.5 9					

18- Mar	H25	H50	H75	PL	EW	Δ H25	Δ H50	Δ H75	Δ PL	Δ EW
H25	0.75$\pm$ 0.06	0.98 $\pm$ 0.01	0.94 $\pm$ 0.01	0.98 $\pm$ 0.02	0.97 $\pm$ 0.05	0.65 $\pm$ 0.01	0.69 $\pm$ 0.02	0.55 $\pm$ 0.02	0.73 $\pm$ 0.03	0.84 $\pm$ 0.06
H50	0.99 $\pm$ 0.00	0.74$\pm$ 0.06	0.96 $\pm$ 0.01	0.98 $\pm$ 0.02	0.96 $\pm$ 0.05	0.64 $\pm$ 0.01	0.74 $\pm$ 0.01	0.60 $\pm$ 0.01	0.74 $\pm$ 0.02	0.83 $\pm$ 0.05
H75	0.97 $\pm$ 0.01	0.98 $\pm$ 0.01	0.71$\pm$ 0.06	0.94 $\pm$ 0.03	0.91 $\pm$ 0.08	0.64 $\pm$ 0.01	0.73 $\pm$ 0.01	0.70 $\pm$ 0.01	0.74 $\pm$ 0.03	0.82 $\pm$ 0.05
PL	0.99 $\pm$ 0.00	0.99 $\pm$ 0.00	0.96 $\pm$ 0.01	0.76$\pm$ 0.06	0.98 $\pm$ 0.02	0.62 $\pm$ 0.01	0.68 $\pm$ 0.01	0.55 $\pm$ 0.01	0.74 $\pm$ 0.01	0.84 $\pm$ 0.02
EW	0.99 $\pm$ 0.00	0.98 $\pm$ 0.01	0.94 $\pm$ 0.02	0.99 $\pm$ 0.00	0.75$\pm$ 0.06	0.6 $\pm$ 0. 00	0.66 $\pm$ 0.00	0.52 $\pm$ 0.00	0.73 $\pm$ 0.01	0.87 $\pm$ 0.01
Δ H2	0.83 $\pm$ 0.07	0.86 $\pm$ 0.06	0.89 $\pm$ 0.06	0.83 $\pm$ 0.07	0.78 $\pm$ 0.08	0.28$\pm$ 0.07	0.76 $\pm$ 0.03	0.58 $\pm$ 0.03	0.69 $\pm$ 0.06	0.69 $\pm$ 0.15
Δ H5	0.77 $\pm$ 0.06	0.81 $\pm$ 0.05	0.84 $\pm$ 0.05	0.79 $\pm$ 0.06	0.75 $\pm$ 0.07	1.00 $\pm$ 0.01	0.42$\pm$ 0.07	0.84 $\pm$ 0.02	0.88 $\pm$ 0.04	0.83 $\pm$ 0.10
Δ H7	0.60 $\pm$ 0.10	0.65 $\pm$ 0.09	0.71 $\pm$ 0.08	0.59 $\pm$ 0.10	0.53 $\pm$ 0.11	1.00 $\pm$ 0.03	0.99 $\pm$ 0.02	0.35$\pm$ 0.08	0.77 $\pm$ 0.05	0.70 $\pm$ 0.13
Δ PL	0.83 $\pm$ 0.05	0.86 $\pm$ 0.05	0.88 $\pm$ 0.04	0.84 $\pm$ 0.05	0.80 $\pm$ 0.06	1.00 $\pm$ 0.02	1.00 $\pm$ 0.01	0.97 $\pm$ 0.05	0.46$\pm$ 0.08	0.95 $\pm$ 0.03
Δ E	0.93 $\pm$ 0.02	0.93 $\pm$ 0.02	0.93 $\pm$ 0.03	0.93 $\pm$ 0.02	0.93 $\pm$ 0.02	0.96 $\pm$ 0.03	0.95 $\pm$ 0.03	0.84 $\pm$ 0.07	0.95 $\pm$ 0.01	0.56$\pm$ 0.07
W										
vA	32.03	52.68	46.27	243.2 7	1159. 21	24.37	24.26	29.48	98.7	477.9 8

18- Jun	H25	H50	H75	PL	EW	Δ H25	Δ H50	Δ H75	Δ PL	Δ EW
H25	0.74$\pm$ 0.06	0.99 $\pm$ 0.01	0.94 $\pm$ 0.01	0.99 $\pm$ 0.02	0.98 $\pm$ 0.05	0.72 $\pm$ 0.01	0.75 $\pm$ 0.01	0.56 $\pm$ 0.02	0.78 $\pm$ 0.03	0.91 $\pm$ 0.06

A. SUPPLEMENTARY MATERIAL

H50	0.99± 0.00	0.73± 0.06	0.96± 0.01	0.98± 0.02	0.96± 0.05	0.7±0. 01	0.78± 0.01	0.58± 0.01	0.77± 0.03	0.89± 0.05
H75	0.98± 0.01	0.99± 0.00	0.72± 0.06	0.93± 0.03	0.92± 0.09	0.68± 0.01	0.75± 0.01	0.70± 0.01	0.74± 0.03	0.85± 0.07
PL	1.00± 0.00	0.99± 0.00	0.97± 0.01	0.69± 0.07	0.98± 0.02	0.70± 0.01	0.74± 0.01	0.54± 0.01	0.78± 0.01	0.91± 0.02
EW	0.99± 0.00	0.98± 0.01	0.97± 0.01	0.99± 0.00	0.69± 0.06	0.69± 0.00	0.73± 0.00	0.53± 0.00	0.76± 0.00	0.93± 0.01
ΔH2 5	0.87± 0.05	0.85± 0.05	0.86± 0.06	0.87± 0.05	0.84± 0.06	0.37± 0.08	0.81± 0.02	0.59± 0.03	0.79± 0.05	0.78± 0.17
ΔH5 0	0.87± 0.05	0.87± 0.04	0.86± 0.05	0.87± 0.05	0.87± 0.05	1.00± 0.01	0.41± 0.08	0.81± 0.02	0.91± 0.03	0.86± 0.12
ΔH7 5	0.75± 0.09	0.76± 0.08	0.77± 0.08	0.76± 0.09	0.75± 0.09	0.97± 0.04	0.98± 0.02	0.27± 0.07	0.72± 0.06	0.66± 0.19
ΔPL	0.85± 0.05	0.83± 0.05	0.83± 0.05	0.85± 0.05	0.85± 0.05	0.99± 0.01	0.97± 0.02	0.93± 0.05	0.41± 0.08	0.93± 0.04
ΔE W	0.95± 0.02	0.93± 0.02	0.93± 0.02	0.95± 0.02	0.96± 0.01	0.95± 0.03	0.95± 0.02	0.86± 0.07	0.96± 0.02	0.57± 0.07
νA	41.36	64.68	45.64	346.0 3	2126. 6	29.73	32.1	33.25	149.4 1	997.8

18- Oct	H25	H50	H75	PL	EW	ΔH25	ΔH50	ΔH75	ΔPL	ΔEW
H25	0.74± 0.06	0.99± 0.01	0.95± 0.01	0.98± 0.02	0.97± 0.06	0.75± 0.01	0.79± 0.01	0.54± 0.02	0.78± 0.03	0.93± 0.07
H50	0.99± 0.00	0.75± 0.06	0.96± 0.01	0.98± 0.02	0.96± 0.06	0.74± 0.01	0.79± 0.01	0.55± 0.01	0.78± 0.03	0.92± 0.06
H75	0.98± 0.01	0.99± 0.01	0.68± 0.07	0.94± 0.04	0.93± 0.11	0.71± 0.02	0.77± 0.02	0.64± 0.02	0.74± 0.04	0.88± 0.09
PL	0.99± 0.00	0.99± 0.00	0.97± 0.01	0.7±0. 06	0.98± 0.02	0.72± 0.01	0.77± 0.01	0.53± 0.01	0.79± 0.01	0.94± 0.02
EW	0.99± 0.00	0.98± 0.01	0.96± 0.01	0.99± 0.00	0.7±0. 06	0.70± 0.00	0.74± 0.00	0.50± 0.00	0.76± 0.00	0.95± 0.01
ΔH2 5	0.86± 0.05	0.85± 0.05	0.85± 0.06	0.85± 0.05	0.83± 0.06	0.46± 0.08	0.86± 0.02	0.64± 0.02	0.81± 0.05	0.78± 0.19
ΔH5 0	0.88± 0.04	0.88± 0.04	0.86± 0.04	0.87± 0.04	0.86± 0.05	1.00± 0.01	0.52± 0.08	0.82± 0.02	0.94± 0.03	0.87± 0.14
ΔH7 5	0.66± 0.11	0.65± 0.10	0.67± 0.1	0.66± 0.11	0.64± 0.11	0.94± 0.04	0.90± 0.05	0.28± 0.07	0.75± 0.07	0.64± 0.27
ΔPL	0.85± 0.05	0.83± 0.05	0.81± 0.06	0.85± 0.05	0.83± 0.05	0.99± 0.01	0.98± 0.01	0.88± 0.06	0.47± 0.08	0.91± 0.05
ΔE W	0.96± 0.01	0.95± 0.01	0.93± 0.02	0.96± 0.01	0.97± 0.01	0.93± 0.03	0.94± 0.03	0.75± 0.09	0.94± 0.02	0.63± 0.07
νA	44.53	63.09	46.72	348.6 7	2791. 05	31.19	31.57	32.3	156.9 4	1407. 77

19- Jan	H25	H50	H75	PL	EW	ΔH25	ΔH50	ΔH75	ΔPL	ΔEW
H25	0.730. 06	0.98± 0.01	0.96± 0.01	0.98± 0.02	0.97± 0.09	0.80± 0.01	0.79± 0.02	0.68± 0.02	0.80± 0.04	0.94± 0.09
H50	0.99± 0.00	0.72± 0.06	0.97± 0.01	0.98± 0.02	0.96± 0.08	0.80± 0.01	0.82± 0.01	0.71± 0.01	0.80± 0.03	0.94± 0.08
H75	0.97± 0.01	0.99± 0.00	0.67± 0.07	0.95± 0.03	0.94± 0.13	0.76± 0.02	0.79± 0.02	0.74± 0.01	0.78± 0.04	0.91± 0.11

PL	0.99± 0.00	0.98± 0.00	0.96± 0.01	0.7±0. 07	0.98± 0.03	0.78± 0.01	0.79± 0.01	0.68± 0.01	0.82± 0.01	0.96± 0.03
EW	0.98± 0.01	0.97± 0.01	0.95± 0.01	0.99± 0.00	0.71± 0.06	0.75± 0.00	0.75± 0.00	0.64± 0.00	0.78± 0.00	0.97± 0.01
ΔH2	0.88± 0.04	0.87± 0.04	0.82± 0.05	0.84± 0.04	0.80± 0.05	0.54± 0.08	0.90± 0.02	0.77± 0.02	0.87± 0.05	0.82± 0.23
ΔH5	0.90± 0.04	0.90± 0.03	0.85± 0.04	0.87± 0.04	0.84± 0.05	1.00± 0.00	0.5±0. 08	0.90± 0.01	0.95± 0.03	0.85± 0.19
ΔH7	0.90± 0.06	0.90± 0.05	0.86± 0.06	0.88± 0.06	0.85± 0.07	1.00± 0.01	0.99± 0.02	0.32± 0.08	0.87± 0.05	0.75± 0.31
ΔPL	0.91± 0.04	0.89± 0.04	0.85± 0.05	0.9±0. 04	0.88± 0.04	0.98± 0.01	0.98± 0.01	0.99± 0.01	0.46± 0.08	0.89± 0.07
ΔE	0.98± 0.01	0.96± 0.01	0.94± 0.02	0.98± 0.01	0.99± 0.00	0.86± 0.04	0.90± 0.03	0.91± 0.05	0.93± 0.02	0.64± 0.07
W										
νA	56.37	84.17	61.28	440.3 6	5100. 25	41.43	56.06	45.67	287.9 1	3507. 55

19- Apr	H25	H50	H75	PL	EW	ΔH25	ΔH50	ΔH75	ΔPL	ΔEW
H25	0.72± 0.06	0.99± 0.01	0.95± 0.01	0.98± 0.02	0.97± 0.09	0.81± 0.02	0.80± 0.02	0.69± 0.02	0.79± 0.04	0.96± 0.09
H50	1.00± 0.00	0.72± 0.06	0.97± 0.01	0.98± 0.02	0.96± 0.09	0.81± 0.01	0.81± 0.01	0.71± 0.01	0.80± 0.03	0.95± 0.08
H75	0.97± 0.01	0.98± 0.01	0.68± 0.07	0.95± 0.03	0.94± 0.14	0.77± 0.02	0.78± 0.02	0.75± 0.02	0.76± 0.04	0.92± 0.12
PL	0.99± 0.00	0.98± 0.00	0.96± 0.01	0.7±0. 07	0.98± 0.03	0.79± 0.01	0.78± 0.01	0.69± 0.01	0.81± 0.01	0.97± 0.03
EW	0.98± 0.01	0.98± 0.01	0.95± 0.01	0.99± 0.00	0.72± 0.06	0.75± 0.00	0.74± 0.00	0.65± 0.00	0.76± 0.00	0.98± 0.01
ΔH2	0.89± 0.04	0.88± 0.04	0.84± 0.05	0.86± 0.04	0.82± 0.05	0.52± 0.08	0.91± 0.02	0.79± 0.02	0.89± 0.04	0.83± 0.24
ΔH5	0.90± 0.04	0.89± 0.04	0.86± 0.05	0.88± 0.04	0.87± 0.05	0.97± 0.04	0.48± 0.08	0.91± 0.01	0.95± 0.02	0.84± 0.21
ΔH7	0.82± 0.06	0.82± 0.06	0.82± 0.06	0.81± 0.07	0.78± 0.07	0.98± 0.02	0.96± 0.02	0.38± 0.08	0.88± 0.05	0.75± 0.33
ΔPL	0.89± 0.04	0.87± 0.04	0.84± 0.05	0.89± 0.04	0.87± 0.04	0.98± 0.01	0.98± 0.01	0.96± 0.02	0.46± 0.08	0.87± 0.09
ΔE	0.98± 0.01	0.97± 0.01	0.95± 0.01	0.99± 0.00	0.99± 0.00	0.88± 0.04	0.92± 0.03	0.84± 0.06	0.93± 0.03	0.66± 0.07
W										
νA	65.4	90.5	64.27	450.7	6379. 85	57	71.92	51.43	335.8 6	4711. 91

19- Aug	H25	H50	H75	PL	EW	ΔH25	ΔH50	ΔH75	ΔPL	ΔEW
H25	0.7±0. 07	0.99± 0.01	0.93± 0.01	0.98± 0.02	0.96± 0.11	0.83± 0.02	0.81± 0.02	0.70± 0.02	0.80± 0.04	0.95± 0.10
H50	0.99± 0.00	0.71± 0.07	0.96± 0.01	0.97± 0.02	0.96± 0.11	0.81± 0.01	0.82± 0.01	0.73± 0.01	0.79± 0.04	0.94± 0.10
H75	0.97± 0.01	0.98± 0.01	0.69± 0.07	0.92± 0.04	0.91± 0.18	0.74± 0.02	0.77± 0.02	0.78± 0.02	0.73± 0.05	0.90± 0.16
PL	0.99± 0.00	0.98± 0.01	0.95± 0.01	0.69± 0.07	0.98± 0.03	0.80± 0.01	0.80± 0.01	0.70± 0.01	0.82± 0.02	0.97± 0.03
EW	0.98± 0.01	0.97± 0.01	0.95± 0.02	0.99± 0.00	0.72± 0.07	0.76± 0.00	0.75± 0.00	0.66± 0.00	0.77± 0.00	0.98± 0.01

A. SUPPLEMENTARY MATERIAL

$\Delta H2$	$0.90\pm$	$0.88\pm$	$0.86\pm$	$0.88\pm$	$0.86\pm$	$0.48\pm$	$0.92\pm$	$0.77\pm$	$0.89\pm$	$0.83\pm$
5	0.03	0.04	0.05	0.04	0.05	0.08	0.02	0.02	0.04	0.27
$\Delta H5$	$0.90\pm$	$0.89\pm$	$0.87\pm$	$0.89\pm$	$0.87\pm$	$1.00\pm$	$0.48\pm$	$0.90\pm$	$0.95\pm$	$0.84\pm$
0	0.04	0.04	0.05	0.04	0.05	0.00	0.08	0.01	0.03	0.25
$\Delta H7$	$0.85\pm$	$0.85\pm$	$0.84\pm$	$0.84\pm$	$0.82\pm$	$0.98\pm$	$0.98\pm$	$0.41\pm$	$0.85\pm$	$0.74\pm$
5	0.06	0.05	0.05	0.06	0.07	0.02	0.01	0.08	0.06	0.39
ΔPL	$0.86\pm$	$0.83\pm$	$0.80\pm$	$0.87\pm$	$0.85\pm$	$0.98\pm$	$0.98\pm$	$0.96\pm$	$0.5\pm 0.$	$0.87\pm$
	0.04	0.05	0.06	0.04	0.05	0.01	0.01	0.02	08	0.10
ΔE	$0.97\pm$	$0.96\pm$	$0.94\pm$	$0.99\pm$	$0.99\pm$	$0.91\pm$	$0.92\pm$	$0.87\pm$	$0.92\pm$	$0.68\pm$
W	0.01	0.01	0.02	0.00	0.00	0.03	0.03	0.05	0.03	0.07
νA	71.69	93.33	67.95	471.7	7713.	62.68	70.63	53.42	339.3	5792.
				4	83				3	1

19- Nov	H25	H50	H75	PL	EW	$\Delta H25$	$\Delta H50$	$\Delta H75$	ΔPL	ΔEW
H25	$0.73\pm$	$0.99\pm$	$0.95\pm$	$0.98\pm$	$0.97\pm$	$0.82\pm$	$0.82\pm$	$0.75\pm$	$0.79\pm$	$0.95\pm$
	0.06	0.01	0.01	0.02	0.11	0.02	0.02	0.02	0.04	0.11
H50	$0.99\pm$	$0.72\pm$	$0.97\pm$	$0.97\pm$	$0.96\pm$	$0.81\pm$	$0.83\pm$	$0.77\pm$	$0.79\pm$	$0.95\pm$
	0.00	0.06	0.01	0.02	0.10	0.01	0.01	0.01	0.03	0.10
H75	$0.95\pm$	$0.98\pm$	$0.68\pm$	$0.94\pm$	$0.93\pm$	$0.75\pm$	$0.78\pm$	$0.80\pm$	$0.75\pm$	$0.91\pm$
	0.01	0.01	0.07	0.03	0.16	0.02	0.02	0.01	0.04	0.15
PL	$0.99\pm$	$0.98\pm$	$0.94\pm$	$0.69\pm$	$0.98\pm$	$0.80\pm$	$0.81\pm$	$0.76\pm$	$0.82\pm$	$0.97\pm$
	0.00	0.00	0.01	0.07	0.03	0.01	0.01	0.01	0.01	0.03
EW	$0.98\pm$	$0.97\pm$	$0.93\pm$	$1.00\pm$	$0.72\pm$	$0.76\pm$	$0.77\pm$	$0.72\pm$	$0.78\pm$	$0.98\pm$
	0.01	0.01	0.02	0.00	0.06	0.00	0.00	0.00	0.00	0.01
$\Delta H2$	$0.89\pm$	$0.88\pm$	$0.82\pm$	$0.89\pm$	$0.86\pm$	$0.56\pm$	$0.92\pm$	$0.81\pm$	$0.89\pm$	$0.83\pm$
5	0.03	0.04	0.05	0.04	0.04	0.08	0.02	0.02	0.04	0.28
$\Delta H5$	$0.88\pm$	$0.88\pm$	$0.84\pm$	$0.88\pm$	$0.87\pm$	$1.00\pm$	$0.54\pm$	$0.92\pm$	$0.96\pm$	$0.85\pm$
0	0.04	0.03	0.04	0.04	0.04	0.00	0.08	0.01	0.02	0.24
$\Delta H7$	$0.88\pm$	$0.89\pm$	$0.86\pm$	$0.88\pm$	$0.86\pm$	$0.97\pm$	$0.99\pm$	$0.43\pm$	$0.89\pm$	$0.79\pm$
5	0.05	0.05	0.05	0.05	0.06	0.05	0.01	0.08	0.05	0.33
ΔPL	$0.85\pm$	$0.83\pm$	$0.77\pm$	$0.86\pm$	$0.86\pm$	$0.99\pm$	$0.98\pm$	$0.95\pm$	$0.51\pm$	$0.87\pm$
	0.04	0.04	0.06	0.04	0.04	0.01	0.01	0.02	0.08	0.10
ΔE	$0.97\pm$	$0.96\pm$	$0.91\pm$	$0.99\pm$	$0.99\pm$	$0.92\pm$	$0.92\pm$	$0.90\pm$	$0.92\pm$	$0.68\pm$
W	0.01	0.01	0.02	0.00	0.00	0.03	0.03	0.05	0.03	0.07
νA	66.05	98.03	78.91	524.0	9178.	59.67	74.68	63.01	388.1	7194.
				9	11				2	48

A.3 Chapter 3

Supplementary Table 3.1: Details of the 113 loci associated with growth identified via Genome-Wide Association Study (GWAS) in trevally. LG: linkage group; SNP: Single Nucleotide Polymorphism; MAF: Minor Allele Frequency; FDR: False Discovery Rate; PVE: Percentage of Variance Explained.

Time	Trait	L G	SNP	Position	<i>p</i> -value	MA F	n	FDR Adjusted <i>p</i> - value	PVE
Nov 2017	H75	1	trevally000070_63741 29	6374129	1.38E- 08	0.19	102 4	2.08E- 04	3.47
Oct 2018	PL	1	trevally000070_97319 65	9731965	7.81E- 08	0.18	672	1.18E- 03	4.75
Nov 2017	H50	1	trevally000070_18446 546	1844654	2.04E- 09	0.11	102 4	1.23E- 04	3.83
Nov 2019	ΔE	1	trevally000070_18794 305	1879430	1.31E- 09	0.33	648	1.39E- 04	6.12
Nov 2017	EW	1	trevally000070_20239 525	2023952	2.69E- 08	0.13	102 4	3.36E- 04	3.35
Nov 2017	H75	1	trevally000070_20362 359	2036235	5.43E- 09	0.22	102 4	1.15E- 04	3.65
Oct 2018	ΔPL	2	trevally000099_56968 9	569689	9.82E- 09	0.11	669	2.08E- 04	5.36
Oct 2018	H25	2	trevally000099_12443 59	1244359	9.60E- 08	0.16	672	1.45E- 03	4.70
Oct 2018	PL	2	trevally000099_12443 59	1244359	7.32E- 09	0.16	672	1.30E- 04	5.42
Oct 2018	EW	2	trevally000099_12443 59	1244359	1.13E- 08	0.16	672	1.20E- 03	5.30
Nov 2017	EW	2	trevally000099_37931 17	3793117	1.87E- 08	0.15	102 4	2.82E- 04	3.42
Nov 2019	H50	2	trevally000099_72251 08	7225108	5.78E- 08	0.17	652	8.72E- 04	4.98
Nov 2019	H50	2	trevally000099_82096 85	8209685	2.04E- 08	0.29	652	3.60E- 04	5.29
Nov 2017	H75	2	trevally000099_11783 136	1178313	8.99E- 08	0.07	102 4	9.50E- 04	3.12
Nov 2017	H75	2	trevally000099_12134 953	1213495	2.55E- 08	0.10	102 4	3.00E- 04	3.36
Oct 2018	H25	2	trevally000099_12159 700	1215970	6.91E- 11	0.33	672	7.31E- 06	6.73
Oct 2018	PL	2	trevally000099_12159 700	1215970	6.94E- 10	0.33	672	3.67E- 05	6.08
Nov 2017	H25	2	trevally000099_15042 047	1504204	1.04E- 08	0.43	102 4	3.66E- 04	3.53
Nov 2017	PL	3	trevally000114_20796 97	2079697	4.22E- 09	0.46	102 4	1.49E- 04	3.70
Nov 2017	H50	3	trevally000114_23934 42	2393442	6.59E- 08	0.03	102 4	9.95E- 04	3.18
Nov 2019	PL	3	trevally000114_49564 08	4956408	4.87E- 10	0.11	652	2.57E- 05	6.37
Oct 2018	H50	3	trevally000114_17345 627	1734562	3.90E- 08	0.08	672	1.03E- 03	4.95
Nov 2019	H75	3	trevally000114_17886 771	1788677	2.33E- 08	0.09	652	4.92E- 04	5.25

A. SUPPLEMENTARY MATERIAL

Nov 2017	H50	3	trevally000114_18608059	18608059	2.42E-08	0.47	1024	6.38E-04	3.37
Nov 2019	Δ H2	4	trevally000118_3914965	3914965	8.11E-09	0.24	648	4.46E-04	5.59
Oct 2018	Δ H5	4	trevally000118_15037078	15037078	2.17E-08	0.28	669	4.38E-04	5.14
Nov 2019	H75	5	trevally000147_51190	51190	2.23E-09	0.35	652	7.85E-05	5.93
Nov 2019	EW	5	trevally000147_171904	171904	6.89E-17	0.09	652	7.28E-12	10.79
Oct 2018	Δ E	5	trevally000147_518680	518680	2.61E-10	0.15	669	1.38E-05	6.38
Oct 2018	W	5	trevally000147_518680	518680	6.18E-09	0.15	672	1.30E-04	5.47
Nov 2019	Δ E	5	trevally000147_1384576	1384576	4.98E-09	0.09	648	2.63E-04	5.73
Nov 2019	W	5	trevally000147_1800159	1800159	1.98E-15	0.08	652	2.10E-10	9.86
Nov 2019	H25	5	trevally000147_1800159	1800159	2.45E-11	0.08	652	2.59E-06	7.22
Nov 2019	H50	5	trevally000147_1800159	1800159	1.17E-14	0.08	652	1.23E-09	9.37
Nov 2017	EW	5	trevally000147_7503742	7503742	1.38E-11	0.10	1024	7.31E-07	4.77
Nov 2017	PL	5	trevally000147_15390868	15390868	1.04E-11	0.09	1024	1.10E-06	4.82
Nov 2017	PL	5	trevally000147_18467804	18467804	4.06E-12	0.10	1024	2.15E-07	4.99
Oct 2018	H25	5	trevally000147_19868051	19868051	2.05E-08	0.24	672	4.74E-04	5.13
Oct 2018	H25	5	trevally000147_19868051	19868051	2.03E-08	0.24	672	7.15E-04	5.14
Nov 2017	H50	6	trevally000312_5886000	5886000	4.24E-08	0.15	1024	8.97E-04	3.26
Nov 2017	PL	6	trevally000312_5886000	5886000	7.11E-08	0.15	1024	1.07E-03	3.16
Nov 2019	H75	6	trevally000312_21977368	21977368	1.12E-11	0.08	652	1.19E-06	7.44
Oct 2018	H25	6	trevally000312_22187402	22187402	1.02E-09	0.13	672	3.58E-05	5.98
Oct 2018	H25	6	trevally000312_23789636	23789636	7.63E-10	0.11	672	3.58E-05	6.06
Oct 2018	PL	7	trevally000314_4776498	4776498	7.36E-09	0.11	672	1.30E-04	5.42
Nov 2019	H50	7	trevally000314_9897116	9897116	6.07E-09	0.27	652	1.60E-04	5.64
Nov 2019	H75	7	trevally000314_16022716	16022716	6.10E-09	0.11	652	1.61E-04	5.64
Nov 2017	EW	7	trevally000314_17150364	17150364	8.42E-12	0.12	1024	7.31E-07	4.86
Oct 2018	Δ E	7	trevally000314_17944233	17944233	1.15E-10	0.16	669	1.21E-05	6.61
Nov 2017	W	7	trevally000314_19873866	19873866	1.85E-12	0.11	1024	1.96E-07	5.14
Nov 2017	H25	9	trevally000429_863530	863530	7.17E-11	0.09	1024	7.58E-06	4.46
Nov 2017	H25	9	trevally000429_23757012	23757012	3.42E-09	0.10	1024	1.81E-04	3.74
Nov 2017	H75	9	trevally000429_23757012	23757012	6.65E-09	0.10	1024	1.17E-04	3.61

Nov 2017	EW	9	trevally000429_23757012	23757012	2.86E-08	0.10	1024	3.36E-04	3.34
Nov 2017	H50	10	trevally000453_1834346	1834346	1.04E-08	0.17	1024	3.65E-04	3.53
Nov 2017	H75	10	trevally000453_6611189	6611189	6.55E-10	0.18	1024	2.31E-05	4.05
Nov 2017	H25	10	trevally000453_6795704	6795704	4.43E-08	0.34	1024	9.36E-04	3.25
Nov 2017	EW	10	trevally000453_25808027	25808027	3.57E-08	0.35	1024	3.77E-04	3.29
Nov 2019	EW	10	trevally000453_26669143	26669143	5.74E-08	0.09	652	2.02E-03	4.99
Oct 2018	$\Delta H2$	11	trevally000657_5815843	5815843	1.77E-08	0.07	669	9.36E-04	5.20
Oct 2018	ΔPL	11	trevally000657_1753543	1753543	5.22E-08	0.10	669	7.88E-04	4.89
Nov 2017	EW	11	trevally000657_8005070	8005070	6.13E-08	0.10	1024	5.89E-04	3.19
Oct 2018	ΔPL	11	trevally000657_21597270	21597270	3.44E-13	0.11	669	1.82E-08	8.22
Oct 2018	$\Delta H5$	11	trevally000657_23735033	23735033	2.49E-08	0.12	669	4.38E-04	5.10
Nov 2017	PL	12	trevally000836_956217	956217	5.57E-08	0.08	1024	9.81E-04	3.21
Nov 2017	EW	12	trevally000836_956217	956217	1.15E-10	0.08	1024	4.06E-06	4.37
Nov 2019	ΔPL	12	trevally000836_5995009	5995009	7.67E-09	0.15	648	4.05E-04	5.60
Oct 2018	H25	12	trevally000836_7695006	7695006	2.24E-08	0.16	672	4.74E-04	5.11
Oct 2018	H50	12	trevally000836_8494592	8494592	9.52E-11	0.09	672	5.03E-06	6.64
Oct 2018	H75	12	trevally000836_8494592	8494592	8.12E-08	0.09	672	8.58E-03	4.74
Oct 2018	PL	12	trevally000836_8494592	8494592	2.70E-12	0.09	672	2.85E-07	7.62
Nov 2017	EW	12	trevally000836_9949138	9949138	1.24E-08	0.29	1024	2.19E-04	3.49
Nov 2017	EW	12	trevally000836_20462163	20462163	4.18E-10	0.16	1024	1.10E-05	4.13
Nov 2019	EW	12	trevally000836_22237578	22237578	6.39E-09	0.27	652	3.38E-04	5.62
Nov 2017	H75	13	trevally000874_3396466	3396466	3.80E-09	0.49	1024	1.00E-04	3.72
Nov 2017	H50	14	trevally001025_997527	997527	5.28E-08	0.04	1024	9.31E-04	3.22
Nov 2019	H50	14	trevally001025_1848158	1848158	7.61E-10	0.07	652	2.68E-05	6.24
Nov 2019	H25	14	trevally001025_8332327	8332327	4.03E-08	0.07	652	1.42E-03	5.09
Nov 2019	H25	14	trevally001025_17510482	17510482	1.71E-08	0.10	652	9.05E-04	5.34
Nov 2019	H50	14	trevally001025_18642018	18642018	1.02E-08	0.13	652	2.16E-04	5.49
Oct 2018	PL	14	trevally001025_22043161	22043161	1.51E-09	0.08	672	5.32E-05	5.87
Nov 2019	$\Delta H5$	14	trevally001025_25939455	25939455	3.99E-08	0.04	648	2.17E-03	5.12
Nov 2019	ΔPL	14	trevally001025_25939455	25939455	2.28E-08	0.04	648	8.03E-04	5.29

A. SUPPLEMENTARY MATERIAL

Oct 2018	ΔPL	14	trevally001025_26800 655	2680065 5	5.26E- 13	0.08	669	1.85E- 08	8.11
Nov 2017	PL	15	trevally001072_34222 7	342227	9.28E- 08	0.05	4	1.23E- 03	3.11
Oct 2018	ΔH5	15	trevally001072_36591 35	3659135	1.54E- 10	0.18	669	8.14E- 06	6.53
Nov 2019	ΔH2	15	trevally001072_18228 519	1822851 9	8.44E- 09	0.11	648	4.46E- 04	5.58
Nov 2019	ΔPL	15	trevally001072_18228 519	1822851 9	4.14E- 08	0.11	648	1.09E- 03	5.11
Nov 2019	ΔH5	16	trevally001123_39056 2	390562	5.16E- 08	0.23	648	2.17E- 03	5.05
Nov 2017	EW	16	trevally001123_11985 91	1198591	5.36E- 10	0.07	4	1.13E- 05	4.08
Nov 2017	PL	17	trevally001154_78974 63	7897463	1.36E- 09	0.17	4	7.19E- 05	3.91
Nov 2017	H75	17	trevally001154_15664 384	1566438	2.36E- 08	0.17	4	3.00E- 04	3.37
Oct 2018	ΔH5	17	trevally001154_27539 969	2753996 9	5.43E- 09	0.10	669	1.91E- 04	5.53
Nov 2019	PL	18	trevally001200_14360 58	1436058	1.06E- 08	0.09	652	3.73E- 04	5.48
Oct 2018	H25	18	trevally001200_16214 03	1621403	4.69E- 08	0.14	672	8.26E- 04	4.90
Oct 2018	ΔPL	19	trevally001460_25100 20	2510020	2.08E- 08	0.12	669	3.67E- 04	5.15
Oct 2018	ΔH5	19	trevally001460_29282 235	2928223 5	7.68E- 09	0.20	669	2.03E- 04	5.43
Oct 2018	ΔPL	19	trevally001460_29282 235	2928223 5	2.92E- 15	0.20	669	3.09E- 10	9.52
Nov 2017	PL	20	trevally001584_32910 34	3291034	6.79E- 09	0.21	4	1.79E- 04	3.61
Oct 2018	ΔH2	20	trevally001584_83154 64	8315464	2.58E- 09	0.16	669	2.73E- 04	5.74
Oct 2018	ΔH5	20	trevally001584_83154 64	8315464	1.36E- 25	0.16	669	1.44E- 20	15.7 3
Oct 2018	ΔPL	20	trevally001584_83154 64	8315464	4.88E- 09	0.16	669	1.29E- 04	5.56
Oct 2018	ΔE	21	trevally001701_34607 60	3460760	3.15E- 08	0.06	669	1.11E- 03	5.03
Nov 2019	ΔH2	21	trevally001701_73194 01	7319401	6.84E- 08	0.22	648	2.41E- 03	4.96
Nov 2017	H25	21	trevally001701_10218 874	1021887 4	2.26E- 08	0.42	4	5.97E- 04	3.38
Nov 2017	H50	21	trevally001701_10218 874	1021887 4	2.32E- 09	0.42	4	1.23E- 04	3.81
Nov 2019	H50	21	trevally001701_13079 733	1307973 3	1.40E- 10	0.10	652	7.40E- 06	6.72
Nov 2019	ΔH5	22	trevally001800_18509 540	1850954 0	6.15E- 08	0.11	648	2.17E- 03	5.00
Nov 2019	ΔPL	22	trevally001800_18509 540	1850954 0	4.87E- 09	0.11	648	4.05E- 04	5.74
Oct 2018	H50	22	trevally001800_19257 649	1925764 9	5.46E- 12	0.07	672	5.77E- 07	7.43
Nov 2017	PL	22	trevally001800_22226 404	2222640 4	4.06E- 08	0.27	4	8.59E- 04	3.27
Nov 2019	PL	23	trevally001891_89006 02	8900602	3.40E- 08	0.31	652	8.98E- 04	5.14
Nov 2019	H75	23	trevally001891_21809 900	2180990 0	7.37E- 11	0.50	652	3.89E- 06	6.91

A.4 Chapter 4

Supplementary Figure 4.1: Comparison of the normalisation factors calculated by either the trimmed mean of M-values (TMM) or the TMM with singleton pairing (TMMwsp) methods for pituitary (PIT) and liver (LIV) samples in the high and low treatments.

Supplementary Figure 4.2: Pairwise exploratory plot of the statistical metrics from the *Danio rerio* BLASTX report. This plot shows the degree of correlation between different BLASTX metrics. pident: percentage of identity; evaluate: the expect value; bitscore: the statistical metric defining similarity (related to the number of identity matches between two sets of sequences).

Supplementary Figure 4.3: Changes in library sizes after filtering. This plot shows the raw library sizes (x-axis) compared with the library sizes after filtering (y-axis) for genes that have sufficiently large counts to be retained in a statistical analysis.

Supplementary Figure 4.5: Exploratory plots after normalisation using the TMM method. (A) Normalisation factors used for pituitary (PIT) and liver (LIV) samples in the high and low treatments, (B) Distribution of the normalisation factors in function of tissue and treatment.

Supplementary Figure 4.6: Dispersion estimates in the pituitary (A) and in the liver (B) samples. Genewise biological coefficients of variation (y-axis) are plotted against gene abundance (x-axis). CPM: Count per Million.

Supplementary Figure 4.7: Enrichment of Gene Ontology (GO) terms associated with the genes that were differentially expressed in the pituitary gland and in the liver between the two temperature treatments. GO-terms enrichment for genes up- or down-regulated in the pituitary (A and B, respectively) and in the liver (C and D, respectively). For each GO-term, the colour saturation indicates the $\log_{10}$ Fisher's p -value of enrichment from smaller (blue) to higher (red) along the x-axis and the semantic space y after multidimensional scaling of GO-terms semantic similarities performed by REVIGO in the y -axis. The log size indicates the number of GO-terms represented.

Supplementary Table 4.1: Summary mapping the samples to Next-Generation Sequencing files and numbers of raw and aligned reads.

Temperature	Tissue	Fish_ID	Prefix	Raw reads	Aligned reads
High	LIV	1	LIV1_HLJ7VDRXX_TTGGTA-TAGGGC_L002	8437401	5576427
High	LIV	2	LIV2_HLJ7VDRXX_GTTACC-TAGGGC_L002	15001477	9189667
High	LIV	3	LIV3_HLJ7VDRXX_CGCAAC-TAGGGC_L002	7137781	4652641
High	LIV	4	LIV4_HLJ7VDRXX_TGGCGA-TAGGGC_L002	9106321	5344004
High	LIV	6	LIV6_HLJ7VDRXX_ACCGTG-TAGGGC_L002	7331929	4433695
High	LIV	7	LIV7_HLJ7VDRXX_CAACAG-TAGGGC_L002	7515064	4871432
High	LIV	8	LIV8_HLJ7VDRXX_GATTGT-TAGGGC_L002	7166218	4251776
High	LIV	9	LIV9_HLJ7VDRXX_ATGGCG-TAGGGC_L002	7763032	4578677
High	LIV	10	LIV10_HJK3MDRXX_TCAGGA-CGCCAT_L001	3974885	2272254
High	LIV	11	LIV11_HJK3MDRXX_CGGTTA-CGCCAT_L001	5605095	3793449
High	LIV	31	LIV31_HJK3MDRXX_GAGTCC-CGCCAT_L001	4909647	2683163
High	LIV	32	LIV32_HJK3MDRXX_GGAGGT-CGCCAT_L001	4960173	2163308
High	LIV	33	LIV33_HJK3MDRXX_CACACT-CGCCAT_L001	5406318	2396143
High	LIV	34	LIV34_HJK3MDRXX_CCGCAA-CGCCAT_L001	5152834	2673470
High	LIV	35	LIV35_HJK3MDRXX_TTTATG-CGCCAT_L001	4174840	3018963
High	LIV	36	LIV36_HJK3MDRXX_AACGCC-CGCCAT_L001	4864172	2574904
High	LIV	37	LIV37_HJK3MDRXX_CAAGCA-CGCCAT_L001	4843682	2629572
High	LIV	38	LIV38_HJK3MDRXX_GCTCGA-CGCCAT_L001	4945868	3190366
High	LIV	39	LIV39_HJK3MDRXX_GCGAAT-CGCCAT_L001	4952246	2923579
High	LIV	40	LIV40_HJK3MDRXX_TGGATT-CGCCAT_L001	5848518	2984547
High	LIV	46	LIV46_HJK3MDRXX_ACCTAC-CGCCAT_L001	4987724	3041861
High	LIV	47	LIV47_HLJ7VDRXX_ATTGGT-TAGGGC_L002	9465255	6670320
High	LIV	48	LIV48_HJK3MDRXX_AGATAG-CGCCAT_L001	3489651	2089499
High	LIV	49	LIV49_HJK3MDRXX_TTGGTA-CGCCAT_L001	4571797	2514551
High	LIV	50	LIV50_HJK3MDRXX_GTTACC-CGCCAT_L001	4422446	2374783
High	LIV	51	LIV51_HJK3MDRXX_CGCAAC-CGCCAT_L001	4317092	2444950
High	LIV	52	LIV52_HJK3MDRXX_TGGCGA-CGCCAT_L001	3958906	1877836

High	LIV	53	LIV53_HJK3MDRXX_ACCGTG-CGCCAT_L001	4235397	1961790
High	LIV	54	LIV54_HJK3MDRXX_CAACAG-CGCCAT_L001	5101179	2689711
High	LIV	55	R-LIV55_HK5YKDRXX_CCTAAG-GCAACG_L001	5151796	2511818
High	LIV	91	LIV91_HJK3MDRXX_GAAGTG-CGCCAT_L001	4764930	2917972
High	LIV	92	LIV92_HJK3MDRXX_CAATGC-CGCCAT_L001	4947005	2442157
High	LIV	93	LIV93_HJK3MDRXX_ACGTCT-CGCCAT_L001	4962543	2682900
High	LIV	94	LIV94_HJK3MDRXX_CAGGAC-CGCCAT_L001	5725461	2738814
High	LIV	95	LIV95_HJK3MDRXX_AAGCTC-CGCCAT_L001	4133716	2454381
High	LIV	96	LIV96_HJK3MDRXX_GACGAT-CGCCAT_L001	6073796	3145864
High	LIV	97	LIV97_HJK3MDRXX_TCGTTC-CGCCAT_L001	4629739	2304375
High	LIV	98	LIV98_HJK3MDRXX_CCAATT-CGCCAT_L001	5545528	2951165
High	LIV	99	LIV99_HJK3MDRXX_AGTTGA-CGCCAT_L001	5191835	2752175
High	LIV	100	LIV100_HJK3MDRXX_AACCGA-CGCCAT_L001	4577712	1986026
High	LIV	136	LIV136_HJK3MDRXX_GCGCTG-CGCCAT_L001	6943494	3893053
High	LIV	137	LIV137_HJK3MDRXX_GAACCT-CGCCAT_L001	6719869	4020331
High	LIV	138	LIV138_HJK3MDRXX_TTCGAG-CGCCAT_L001	6551743	3814532
High	LIV	139	LIV139_HJK3MDRXX_AGAATC-CGCCAT_L001	5645071	3542549
High	LIV	140	LIV140_HJK3MDRXX_AGGCAT-CGCCAT_L001	7020815	4744813
High	LIV	141	LIV141_HLJ7VDRXX_CTCATA-TAGGGC_L002	9392167	6873618
High	LIV	142	LIV142_HLJ7VDRXX_CCGACC-TAGGGC_L002	8469523	5870859
High	LIV	143	LIV143_HLJ7VDRXX_GGCCAA-TAGGGC_L002	9611133	6790572
High	LIV	144	LIV144_HLJ7VDRXX_AGACCA-TAGGGC_L002	10974812	7588885
High	LIV	145	LIV145_HLJ7VDRXX_CGCGGA-TAGGGC_L002	8277286	5562401
Total LIV				308956922	183456598
High	PIT	1	PIT1_HLJ7VDRXX_CAGCGT-ATTTTA_L002	13432277	8879464
High	PIT	2	PIT2_HK5YKDRXX_GATCAC-ATTTTA_L001	4853238	3633884
High	PIT	3	PIT3_HK5YKDRXX_ACCAGT-ATTTTA_L001	5137268	3115574
High	PIT	4	PIT4_HK5YKDRXX_TGCACG-ATTTTA_L001	5400811	3424312
High	PIT	6	PIT6_HK5YKDRXX_ACATTA-ATTTTA_L001	4845146	3111794
High	PIT	7	PIT7_HLJ7VDRXX_GTG TAG-ATTTTA_L002	10886061	6897660

A. SUPPLEMENTARY MATERIAL

High	PIT	8	PIT8_HLJ7VDRXX_CTAGTC- ATTTTA_L002	9822216	6437852
High	PIT	9	PIT9_HLJ7VDRXX_TGTGCA- ATTTTA_L002	11813512	7657798
High	PIT	10	PIT10_HLJ7VDRXX_TCAGGA- ATTTTA_L002	11618137	7467565
High	PIT	11	PIT11_HK5YKDRXX_CGGTTA- ATTTTA_L001	6437083	4749544
High	PIT	31	PIT31_HK5YKDRXX_GAGTCC- ATTTTA_L001	3981069	2463893
High	PIT	32	PIT32_HK5YKDRXX_GGAGGT- ATTTTA_L001	4331897	2070447
High	PIT	33	PIT33_HLJ7VDRXX_CACACT- ATTTTA_L002	9586997	5544328
High	PIT	34	PIT34_HLJ7VDRXX_CCGCAA- ATTTTA_L002	10474169	6446749
High	PIT	35	PIT35_HK5YKDRXX_TTTATG- ATTTTA_L001	4796600	3607182
High	PIT	36	PIT36_HK5YKDRXX_AACGCC- ATTTTA_L001	5435342	2909860
High	PIT	37	PIT37_HK5YKDRXX_CAAGCA- ATTTTA_L001	4837776	2357764
High	PIT	38	PIT38_HK5YKDRXX_GCTCGA- ATTTTA_L001	4802814	2365927
High	PIT	39	PIT39_HK5YKDRXX_GCGAAT- ATTTTA_L001	5481132	3354317
High	PIT	40	PIT40_HK5YKDRXX_TGGATT- ATTTTA_L001	6476090	4514125
High	PIT	46	PIT46_HLJ7VDRXX_ACCTAC- ATTTTA_L002	9433598	6296230
High	PIT	47	PIT47_HLJ7VDRXX_CGAAGG- ATTTTA_L002	12341945	8307862
High	PIT	48	PIT48_HK5YKDRXX_AGATAG- ATTTTA_L001	6233049	4324411
High	PIT	49	PIT49_HK5YKDRXX_TTGGA- ATTTTA_L001	4803215	2996510
High	PIT	50	PIT50_HK5YKDRXX_GTTACC- ATTTTA_L001	4570258	2787105
High	PIT	51	PIT51_HK5YKDRXX_CGCAAC- ATTTTA_L001	5316921	3058794
High	PIT	52	PIT52_HK5YKDRXX_TGGCGA- ATTTTA_L001	5321161	3326371
High	PIT	53	PIT53_HLJ7VDRXX_ACCGTG- ATTTTA_L002	11470368	7045917
High	PIT	54	PIT54_HK5YKDRXX_CAACAG- ATTTTA_L001	4401846	2721774
High	PIT	55	PIT55_HLJ7VDRXX_GATTGT- ATTTTA_L002	11584560	7553929
High	PIT	91	PIT91_HK5YKDRXX_GAAGTG- ATTTTA_L001	5122838	3560813
High	PIT	92	PIT92_HK5YKDRXX_CAATGC- ATTTTA_L001	4454962	2752380
High	PIT	93	PIT93_HK5YKDRXX_ACGTCT- ATTTTA_L001	5765159	3796254
High	PIT	94	PIT94_HK5YKDRXX_CAGGAC- ATTTTA_L001	5601203	3961990
High	PIT	95	PIT95_HK5YKDRXX_AAGCTC- ATTTTA_L001	5075791	2826763
High	PIT	96	PIT96_HK5YKDRXX_GACGAT- ATTTTA_L001	5497602	3113473

High	PIT	97	PIT97_HK5YKDRXX_TCGTTC- ATTTTA_L001	4953838	3194847
High	PIT	98	PIT98_HK5YKDRXX_CCAATT- ATTTTA_L001	4620210	2941476
High	PIT	99	PIT99_HK5YKDRXX_AGTTGA- ATTTTA_L001	4719593	2789018
High	PIT	100	PIT100_HK5YKDRXX_AACCGA- ATTTTA_L001	4436333	2917712
High	PIT	136	PIT136_HK5YKDRXX_GAACCT- ATTTTA_L001	5622671	3655631
High	PIT	137	PIT137_HK5YKDRXX_TTCGAG- ATTTTA_L001	6295154	4392843
High	PIT	138	PIT138_HK5YKDRXX_AGAATC- ATTTTA_L001	5891870	3902628
High	PIT	139	PIT139_HK5YKDRXX_AGGCAT- ATTTTA_L001	6736636	3687532
High	PIT	140	PIT140_HLJ7VDRXX_ACACGC- ATTTTA_L002	9914550	6174432
High	PIT	141	PIT141_HLJ7VDRXX_GCGCTG- TAGGGC_L002	9071190	6348118
High	PIT	142	PIT142_HLJ7VDRXX_GAACCT- TAGGGC_L002	2480151	1632468
High	PIT	143	PIT143_HLJ7VDRXX_TTCGAG- TAGGGC_L002	8374894	5760544
High	PIT	144	PIT144_HLJ7VDRXX_AGAATC- TAGGGC_L002	9096278	6278823
High	PIT	145	PIT145_HLJ7VDRXX_AGGCAT- TAGGGC_L002	9652198	6754247
Total PIT				343309677	219870934
Low	LIV	16	LIV16_HJK3MDRXX_TTAAC- CGCCAT_L001	4823833	2403971
Low	LIV	17	LIV17_HJK3MDRXX_ATGAAC- CGCCAT_L001	4337448	1648408
Low	LIV	18	LIV18_HJK3MDRXX_CCTAAG- CGCCAT_L001	4671360	1958490
Low	LIV	19	LIV19_HJK3MDRXX_AATCCG- CGCCAT_L001	3999766	1814881
Low	LIV	20	LIV20_HJK3MDRXX_GGCTGC- CGCCAT_L001	5382017	2568944
Low	LIV	21	LIV21_HJK3MDRXX_TACCTT- CGCCAT_L001	5741947	3620733
Low	LIV	22	LIV22_HJK3MDRXX_TCTTAA- CGCCAT_L001	3489434	2142362
Low	LIV	23	LIV23_HJK3MDRXX_GTCAGG- CGCCAT_L001	4743705	2527064
Low	LIV	24	LIV24_HJK3MDRXX_ATACTG- CGCCAT_L001	4459765	2430626
Low	LIV	25	LIV25_HJK3MDRXX_TATGTC- CGCCAT_L001	3803018	2081202
Low	LIV	62	LIV62_HLJ7VDRXX_GCCACA- TAGGGC_L002	8477747	5987908
Low	LIV	63	LIV63_HJK3MDRXX_TGACAC- CGCCAT_L001	4166252	2372257
Low	LIV	65	LIV65_HJK3MDRXX_AAGACA- CGCCAT_L001	3876363	1763965
Low	LIV	66	LIV66_HJK3MDRXX_ACAGAT- CGCCAT_L001	3530523	1562356
Low	LIV	67	LIV67_HJK3MDRXX_TAGGCT- CGCCAT_L001	4534424	2187272

Low	LIV	68	LIV68_HJK3MDRXX_CTCCAT-CGCCAT_L001	4993988	2700680
Low	LIV	69	LIV69_HJK3MDRXX_GCATGG-CGCCAT_L001	4311364	2250275
Low	LIV	70	LIV70_HJK3MDRXX_AATAGC-CGCCAT_L001	3711173	2500527
Low	LIV	71	LIV71_HLJ7VDRXX_CATCTA-TAGGGC_L002	8165073	5820597
Low	LIV	72	LIV72_HJK3MDRXX_TCGAGG-CGCCAT_L001	5260440	3078936
Low	LIV	76	LIV76_HJK3MDRXX_CACTAA-CGCCAT_L001	5579751	3277151
Low	LIV	77	LIV77_HJK3MDRXX_GGTATA-CGCCAT_L001	4501929	2457252
Low	LIV	78	LIV78_HJK3MDRXX_CGCCTG-CGCCAT_L001	6472332	4030989
Low	LIV	79	LIV79_HJK3MDRXX_AATGAA-CGCCAT_L001	5099585	2914427
Low	LIV	80	LIV80_HJK3MDRXX_ACAACG-CGCCAT_L001	5483025	2777073
Low	LIV	81	LIV81_HJK3MDRXX_ATATCC-CGCCAT_L001	5224837	2767303
Low	LIV	82	LIV82_HLJ7VDRXX_AACAAG-TAGGGC_L002	9164420	6586139
Low	LIV	83	LIV83_HJK3MDRXX_ATAAGA-CGCCAT_L001	5067208	3107311
Low	LIV	84	LIV84_HJK3MDRXX_GGTGAG-CGCCAT_L001	5567849	3275886
Low	LIV	85	LIV85_HJK3MDRXX_TTCCGC-CGCCAT_L001	5343102	3374881
Low	LIV	106	LIV106_HJK3MDRXX_CAGATG-CGCCAT_L001	4958651	2634890
Low	LIV	107	LIV107_HJK3MDRXX_GTAGAA-CGCCAT_L001	4333018	2810050
Low	LIV	108	LIV108_HJK3MDRXX_GACATC-CGCCAT_L001	7185312	4406526
Low	LIV	109	LIV109_HJK3MDRXX_CGATCT-CGCCAT_L001	6004230	3608176
Low	LIV	110	LIV110_HLJ7VDRXX_GCAGCC-TAGGGC_L002	7554138	4902910
Low	LIV	111	LIV111_HJK3MDRXX_ATGGCG-CGCCAT_L001	5030287	3039609
Low	LIV	112	LIV112_HJK3MDRXX_ATTGGT-CGCCAT_L001	5217047	1960290
Low	LIV	113	LIV113_HJK3MDRXX_GCCACA-CGCCAT_L001	5115480	2772022
Low	LIV	114	LIV114_HLJ7VDRXX_ACTCTT-TAGGGC_L002	8552909	5841585
Low	LIV	115	LIV115_HJK3MDRXX_AACAAG-CGCCAT_L001	5186807	3258396
Low	LIV	121	LIV121_HLJ7VDRXX_TGCTAT-TAGGGC_L002	10154338	7387751
Low	LIV	122	LIV122_HJK3MDRXX_ACTCTT-CGCCAT_L001	6501352	3992482
Low	LIV	123	LIV123_HJK3MDRXX_TGCTAT-CGCCAT_L001	5788153	2693540
Low	LIV	124	LIV124_HJK3MDRXX_AAGTGG-CGCCAT_L001	5253671	2729245
Low	LIV	125	LIV125_HJK3MDRXX CTCATA-CGCCAT_L001	5884769	2980405

Low	LIV	126	LIV126_HJK3MDRXX_CCGACC-CGCCAT_L001	5670948	2763997
Low	LIV	127	LIV127_HJK3MDRXX_GGCCAA-CGCCAT_L001	5136309	3063042
Low	LIV	128	LIV128_HLJ7VDRXX_AAGTGG-TAGGGC_L002	7808118	4861731
Low	LIV	129	LIV129_HJK3MDRXX_CGCGGA-CGCCAT_L001	3796004	1334639
Low	LIV	130	LIV130_HJK3MDRXX_CCTGCT-CGCCAT_L001	5958422	3292556
Total LIV				275073641	158323708
Low	PIT	16	PIT16_HK5YKDRXX_TTAAC-ATTTTA_L001	5467022	4009780
Low	PIT	17	PIT17_HK5YKDRXX_ATGAAC-ATTTTA_L001	4669964	2284705
Low	PIT	18	PIT18_HK5YKDRXX_CCTAAG-ATTTTA_L001	4151268	2795965
Low	PIT	19	PIT19_HK5YKDRXX_AATCCG-ATTTTA_L001	3911401	2234322
Low	PIT	20	PIT20_HLJ7VDRXX_GGCTGC-ATTTTA_L002	10269705	7319872
Low	PIT	21	PIT21_HLJ7VDRXX_TACCTT-ATTTTA_L002	14558889	9169010
Low	PIT	22	PIT22_HLJ7VDRXX_TCTTAA-ATTTTA_L002	9544396	6107403
Low	PIT	23	PIT23_HK5YKDRXX_GTCAGG-ATTTTA_L001	4762037	2508317
Low	PIT	24	PIT24_HK5YKDRXX_ATACTG-ATTTTA_L001	3898742	2030679
Low	PIT	25	PIT25_HK5YKDRXX_TATGTC-ATTTTA_L001	3863957	1897083
Low	PIT	62	PIT62_HLJ7VDRXX_CCTGCT-TAGGGC_L002	9094323	6045749
Low	PIT	63	PIT63_HK5YKDRXX_TGACAC-ATTTTA_L001	5721696	4019789
Low	PIT	65	PIT65_HK5YKDRXX_AAGACA-ATTTTA_L001	4874301	2658988
Low	PIT	66	PIT66_HK5YKDRXX_ACAGAT-ATTTTA_L001	5141674	2834253
Low	PIT	67	PIT67_HK5YKDRXX_TAGGCT-ATTTTA_L001	4269848	2699476
Low	PIT	68	PIT68_HK5YKDRXX_CTCCAT-ATTTTA_L001	5723746	3627050
Low	PIT	69	PIT69_HLJ7VDRXX_GCATGG-ATTTTA_L002	7881872	5051975
Low	PIT	70	PIT70_HLJ7VDRXX_AATAGC-ATTTTA_L002	11570251	8571817
Low	PIT	71	PIT71_HLJ7VDRXX_GTGCCA-ATTTTA_L002	10110292	6499127
Low	PIT	72	PIT72_HK5YKDRXX_TCGAGG-ATTTTA_L001	5391656	3080797
Low	PIT	76	PIT76_HK5YKDRXX_CACTAA-ATTTTA_L001	5181429	3145134
Low	PIT	77	PIT77_HK5YKDRXX_GGTATA-ATTTTA_L001	4945911	3146382
Low	PIT	78	PIT78_HK5YKDRXX_CGCCTG-ATTTTA_L001	5162344	3161321
Low	PIT	79	PIT79_HLJ7VDRXX_AATGAA-ATTTTA_L002	9729050	4845352

A. SUPPLEMENTARY MATERIAL

Low	PIT	80	PIT80_HLJ7VDRXX_ACAACG- ATTTTA_L002	8284310	5053056
Low	PIT	81	PIT81_HLJ7VDRXX_ATATCC- ATTTTA_L002	9902914	5710433
Low	PIT	82	PIT82_HLJ7VDRXX_AGTACT- ATTTTA_L002	12329918	8102298
Low	PIT	83	PIT83_HK5YKDRXX_ATAAGA- ATTTTA_L001	6341812	4444123
Low	PIT	84	PIT84_HK5YKDRXX_GGTGAG- ATTTTA_L001	5556267	3855729
Low	PIT	85	PIT85_HK5YKDRXX_TTCCGC- ATTTTA_L001	4759376	3445445
Low	PIT	106	PIT106_HK5YKDRXX_CAGATG- ATTTTA_L001	4893071	3367863
Low	PIT	107	PIT107_HLJ7VDRXX_GTAGAA- ATTTTA_L002	10573942	6331547
Low	PIT	108	PIT108_HK5YKDRXX_GACATC- ATTTTA_L001	5632139	4040833
Low	PIT	109	PIT109_HK5YKDRXX_CGATCT- ATTTTA_L001	5938762	3600529
Low	PIT	110	PIT110_HK5YKDRXX_CGTCGC- ATTTTA_L001	4984011	3268796
Low	PIT	111	PIT111_HK5YKDRXX_ATGGCG- ATTTTA_L001	4589250	2723159
Low	PIT	112	PIT112_HK5YKDRXX_ATTGGT- ATTTTA_L001	4337834	2814677
Low	PIT	113	PIT113_HK5YKDRXX_GCCACA- ATTTTA_L001	4678972	2459022
Low	PIT	114	PIT114_HK5YKDRXX_CATCTA- ATTTTA_L001	4804212	3203260
Low	PIT	115	PIT115_HLJ7VDRXX_AACAAG- ATTTTA_L002	12856011	9149586
Low	PIT	121	PIT121_HLJ7VDRXX_GCAGCC- ATTTTA_L002	10940143	6916660
Low	PIT	122	PIT122_HK5YKDRXX_ACTCTT- ATTTTA_L001	6017608	4083129
Low	PIT	123	PIT123_HLJ7VDRXX_TGCTAT- ATTTTA_L002	14314889	7929005
Low	PIT	124	PIT124_HK5YKDRXX_AAGTGG- ATTTTA_L001	4724754	1856561
Low	PIT	125	PIT125_HK5YKDRXX_CTCATA- ATTTTA_L001	4202101	2363070
Low	PIT	126	PIT126_HK5YKDRXX_CCGACC- ATTTTA_L001	5086820	2973554
Low	PIT	127	PIT127_HK5YKDRXX_GGCCAA- ATTTTA_L001	4746343	2559534
Low	PIT	128	PIT128_HK5YKDRXX_CGCGGA- ATTTTA_L001	5763678	3833416
Low	PIT	129	PIT129_HK5YKDRXX_CCTGCT- ATTTTA_L001	5125328	3670833
Low	PIT	130	PIT130_HK5YKDRXX_GCGCTG- ATTTTA_L001	4928419	3342851
Total PIT				336208658	210843315
Total High				652266599	403327532
Total Low				611282299	369167023
Total reads				1263548898	772494555

Supplementary Table 4.2: List of the 39 genes found differentially expressed ($|\log_2FC| > 0.26$, p -value < 0.05) in the pituitary gland, in the high versus low condition.

Gene	ACCNUM	ENTREZ ID	SYMBOL	Log ₂ FC High/Low	log CP M	LR	E- value	p- value	FDR adjusted p- value
g12011	CCE39580	402917	lhb	1.67	10.50	60.58	1.41E-45	2.00E-12	1.72E-08
g14247	XP_017213687	30259	prph	-1.53	3.69	56.18	2.92E-146	2.25E-11	9.67E-08
g16082	AHE93330	101887161	tet2	-1.33	4.73	54.93	1.70E-62	1.13E-09	3.23E-06
g27508	A3KQ55	561384	mepce	-1.94	3.32	48.53	0.00E+00	2.30E-09	4.94E-06
g17089	AAI63519	403114	adm2a	-1.57	3.06	46.06	0.00E+00	4.02E-08	6.92E-05
g18798	AAV31152	402919	fshb	1.68	12.52	40.62	4.15E-137	6.48E-08	9.28E-05
g2742	XP_017207109	566645	espnlb	-1.96	2.54	40.62	2.54E-17	9.94E-08	1.22E-04
g369	XP_005172678	563644	kdm6bb	-0.81	8.79	37.52	0.00E+00	3.06E-07	3.29E-04
g24133	AAH49416	393952	sigmar1	-0.92	4.55	36.57	0.00E+00	3.45E-07	3.29E-04
g27177	XP_001334673	30746	tie1	-2.00	2.36	36.08	0.00E+00	4.44E-07	3.82E-04
g20265	XP_001920640	100150283	loxhd1b	-0.90	5.57	35.36	1.10E-109	9.16E-07	6.76E-04
g14339	CAD60658	368823	fkbp11	0.64	8.84	35.24	1.14E-23	9.44E-07	6.76E-04
g23237	XP_005174156	100330283	LOC100330283	-1.09	4.03	35.46	3.13E-28	1.59E-06	1.05E-03
g10829	AAH59702	793658	arl13a	-1.33	4.18	34.39	9.07E-52	3.18E-06	1.95E-03
g10971	XP_009289566	103908779	c14h4orf45	-0.82	6.18	34.08	1.36E-20	3.95E-06	2.26E-03
g12861	XP_021325885	565402	col11a1a	1.22	3.55	33.73	6.23E-09	4.41E-06	2.37E-03
g25215	NP_956951	393630	thyn1	-0.88	4.20	32.17	2.38E-72	5.74E-06	2.87E-03
g27504	XP_017210159	100331067	LOC100331067	-0.81	5.28	32.12	0.00E+00	6.01E-06	2.87E-03
g4773	AAH92763	550548	stmnlb	0.78	5.93	31.72	1.95E-110	1.33E-05	6.02E-03
g25621	XP_017207641	108180053	LOC108180053	0.76	5.27	31.47	1.28E-24	1.90E-05	8.15E-03
g24169	XP_021324735	793912	nefla	1.18	4.22	31.35	5.27E-32	2.94E-05	1.20E-02
g10394	NP_001131027	798788	gpx3	0.78	8.91	31.05	0.00E+00	3.16E-05	1.23E-02
g10440	NP_001338645	568796	pcdh20	0.98	3.94	28.65	1.04E-68	3.32E-05	1.24E-02
g24427	ABB60051	493623	gpsm1b	-0.63	6.58	28.35	0.00E+00	3.82E-05	1.37E-02
g25508	NP_001119871	561195	trpv1	0.92	4.76	27.31	0.00E+00	4.33E-05	1.49E-02
g24324	AAH92881	550498	iscub	-1.06	5.26	26.64	0.00E+00	6.49E-05	2.14E-02

A. SUPPLEMENTARY MATERIAL

g268 5	AAI25914	777641	scg2b	-0.99	5.53	26.0 0	7.48E- 44	7.79E- 05	2.48E- 02
g187 77	XP_005166 545	564462	lrrc4cb	0.93	4.36	26.0 6	3.53E- 155	8.44E- 05	2.59E- 02
g434	AAH92842	572669	dnajb13	-1.41	3.39	25.3 7	3.30E- 19	9.33E- 05	2.76E- 02
g157 80	XP_005171 649	793920	il2rb	1.50	2.16	23.4 6	0.00E+ 00	9.63E- 05	2.76E- 02
g681 8	XP_005173 575	572411	ephb6	0.66	6.10	23.3 4	0.00E+ 00	1.50E- 04	4.16E- 02
g194 48	AAH61705	494154	lancl1	-0.78	4.25	22.7 1	0.00E+ 00	1.55E- 04	4.16E- 02
g119 27	AAD47423	30645	cd74b	0.90	5.00	22.5 7	0.00E+ 00	1.81E- 04	4.70E- 02
g114 19	NP_001026 843	791878	cst3	-1.02	8.18	22.1 7	0.00E+ 00	1.88E- 04	4.72E- 02
g149 35	AAH56531	321818	ctsa	-0.80	4.69	21.7 0	0.00E+ 00	1.98E- 04	4.72E- 02
g256 91	XP_009292 131	1000343 87	sphkap	-0.77	5.00	21.8 0	3.62E- 82	1.99E- 04	4.72E- 02
g289 9	XP_017207 706	563528	prss56	-1.29	2.59	21.7 2	0.00E+ 00	2.03E- 04	4.72E- 02
g176 78	XP_009301 186	1039113 43	si:ch1073- 153i20.4	0.58	6.79	21.5 3	0.00E+ 00	2.14E- 04	4.85E- 02
g492 1	BAF47195	564619	gstr	-0.97	6.75	21.1 8	4.30E- 173	2.22E- 04	4.90E- 02

Supplementary Table 4.3: List of the 238 genes found differentially expressed in the liver ($|\log_2FC| > 0.26$, p -value < 0.05) in the high versus low condition.

Gene	ACCNUM	ENTREZ ID	SYMBOL	Log ₂ F C High/ Low	log CP M	LR	E-value	p- value	FDR adjusted p- value
g7896	AAI24470	558677	zp3a.2	-1.76	6.51	131.08	3.57E-114	4.25E-24	3.66E-20
g10950	AAH59459	317738	elovl6	-2.46	6.97	109.04	0.00E+00	2.04E-22	8.78E-19
g4294	NP_942110	386661	scd	-1.92	7.09	79.51	1.14E-163	4.29E-16	1.23E-12
g11736	A4QP75	606657	trmt2b	-1.60	7.00	79.84	0.00E+00	2.70E-15	5.80E-12
g3745	XP_009305081	559001	fasn	-2.66	3.45	80.31	0.00E+00	3.76E-14	6.46E-11
g25374	AAI62122	568167	thrsp	-1.92	10.97	74.52	0.00E+00	5.07E-13	7.27E-10
g9018	AAX45073	394021	shmt1	-1.17	5.95	69.32	1.01E-76	8.54E-13	1.05E-09
g21643	AAF79948	58128	fabp7a	-2.22	6.44	69.01	5.20E-13	1.32E-12	1.42E-09
g25034	XP_005157555	100037332	zgc:162608	1.27	8.24	67.84	6.29E-84	4.16E-12	3.98E-09
g3123	AAI50352	567406	ahsg2	1.13	11.00	67.14	8.16E-158	7.52E-12	6.46E-09
g19174	XP_003199513	571189	abcb11b	-1.20	5.37	66.75	3.18E-58	1.83E-11	1.43E-08
g20728	XP_005171883	406312	zfand5a	1.03	7.22	65.63	2.21E-36	4.03E-11	2.86E-08
g6991	AAH56293	793727	gnl2	0.83	6.41	63.90	0.00E+00	4.33E-11	2.86E-08
g9808	AAH76484	436922	aclya	-3.36	3.84	63.85	0.00E+00	7.85E-11	4.82E-08
g4032	AAH56596	321618	h1f0	1.03	5.38	61.61	1.36E-113	8.44E-11	4.84E-08
g25035	XP_696211	567816	LOC567816	1.11	8.60	61.22	8.91E-40	1.24E-10	6.67E-08
g14566	XP_693183	564761	itih1	-1.42	3.47	61.08	0.00E+00	1.80E-10	9.11E-08
g12756	AAI17630	554998	pdia4	-1.19	5.17	57.54	1.35E-123	2.15E-10	1.03E-07
g24011	CAI99155	560202	ghrb	1.13	4.91	57.26	1.94E-83	2.85E-10	1.29E-07
g3399	BAE78824	402819	cnpy1	-1.04	6.07	56.92	0.00E+00	7.95E-10	3.27E-07
g23522	AAH46004	322420	chia.6	1.67	3.39	56.77	2.26E-84	7.98E-10	3.27E-07
g14734	XP_002663836	100334148	si:dkey-57n24.6	1.55	3.03	54.11	0.00E+00	2.03E-09	7.92E-07
g24462	XP_017208526	559835	alpk2	-2.10	2.78	54.04	0.00E+00	3.35E-09	1.25E-06
g20065	XP_005165133	393984	aacs	-1.55	5.58	52.78	0.00E+00	4.71E-09	1.69E-06
g8240	AAH91470	560651	ces3	0.94	7.48	52.27	2.24E-100	6.32E-09	2.17E-06
g28139	CAB64945	30708	fabp2	-1.26	5.84	49.78	0.00E+00	6.67E-09	2.21E-06

A. SUPPLEMENTARY MATERIAL

g20278	AAH83399	450001	zgc:103482	-1.55	3.61	48.31	0.00E+00	7.65E-09	2.44E-06
g15238	AAH54574	64608	alas1	1.04	6.75	48.20	1.23E-105	1.48E-08	4.56E-06
g20667	NP_001352946	110437731	LOC110437731	0.68	13.17	48.10	2.76E-69	2.30E-08	6.80E-06
g8257	NP_001093469	560780	si:dkey-238o13.4	-2.90	3.22	47.48	0.00E+00	2.85E-08	8.16E-06
g11458	ABY90095	407621	disc1	0.67	6.35	47.13	9.70E-173	3.88E-08	1.08E-05
g27504	XP_017210159	100331067	LOC100331067	-1.01	4.97	47.06	0.00E+00	5.12E-08	1.38E-05
g8568	AAP47138	386590	hsp90b1	-0.92	7.62	46.70	0.00E+00	5.69E-08	1.48E-05
g8006	XP_005159361	100321139	alpk3a	-1.32	3.60	46.18	0.00E+00	5.96E-08	1.51E-05
g9529	XP_021323416	100007164	cbln14	1.38	6.44	45.61	2.48E-57	7.70E-08	1.89E-05
g15317	ABM68033	767678	cish	1.23	5.93	44.95	0.00E+00	8.74E-08	2.08E-05
g7092	AAH76516	798836	tspan13a	-1.17	4.33	44.88	0.00E+00	8.96E-08	2.08E-05
g9286	AAI71350	573110	cbx4	-0.85	5.83	44.52	1.57E-54	9.50E-08	2.15E-05
g19261	NP_001314912	100537632	spp2	0.96	11.24	44.17	1.36E-20	1.02E-07	2.25E-05
g4478	AAH81587	447860	acsl5	-1.05	7.97	43.85	0.00E+00	1.25E-07	2.69E-05
g18149	NP_001038917	751742	uraha	0.86	5.89	43.64	3.90E-166	1.80E-07	3.77E-05
g8109	XP_005174313	101887155	pnpla2	1.14	3.97	43.61	0.00E+00	2.21E-07	4.52E-05
g16082	AHE93330	101887161	tet2	-1.10	4.57	43.52	9.61E-23	2.64E-07	5.28E-05
g387	XP_017211598	559403	acaca	-1.19	4.23	42.11	0.00E+00	3.27E-07	6.39E-05
g655	NP_001313465	556700	smpd1	-1.01	5.08	41.94	0.00E+00	4.34E-07	8.30E-05
g21862	XP_009291550	101883329	si:dkey-206p8.1	1.17	4.41	41.77	0.00E+00	4.45E-07	8.32E-05
g18087	AAH65657	324010	hnf4b	0.98	5.30	41.48	0.00E+00	5.10E-07	9.29E-05
g26929	AAH76466	436604	uox	0.75	11.12	41.23	0.00E+00	5.18E-07	9.29E-05
g4039	XP_005162937	406623	decr2	-0.73	7.13	41.04	0.00E+00	6.20E-07	1.09E-04
g16143	AAI27404	777738	vps37c	0.97	4.41	40.80	2.92E-109	7.49E-07	1.29E-04
g27511	AAQ97775	378963	eno3	0.72	9.86	40.45	3.21E-39	7.81E-07	1.32E-04
g4159	AAH59706	406298	ndufa4	-1.40	3.94	40.07	0.00E+00	9.75E-07	1.61E-04
g3385	CBN61608	405829	dpydb	0.64	9.27	40.04	1.44E-80	1.01E-06	1.63E-04
g11479	AAI22336	563048	LOC563048	0.58	11.16	39.56	0.00E+00	1.15E-06	1.82E-04
g1602	XP_685872	564300	lrrc58b	0.75	6.57	39.49	0.00E+00	1.20E-06	1.87E-04
g23029	AAH44196	406762	pgd	-1.38	3.52	39.30	0.00E+00	1.25E-06	1.92E-04

g303 9	XP_005163 400	1018833 96	LOC1018833 96	-1.02	4.22	39.0 3	0.00E+ 00	1.29E- 06	1.95E- 04
g962 3	XP_005164 295	572061	pde4a	-0.71	6.01	38.5 2	2.46E- 118	1.33E- 06	1.97E- 04
g228 93	AAI33979	449685	zgc:136410	-0.87	11.4 4	38.3 9	0.00E+ 00	1.53E- 06	2.22E- 04
g214 55	AAI24343	767694	slc25a21	-0.69	6.78	38.4 0	5.75E- 73	1.74E- 06	2.49E- 04
g568 6	AAI28792	664756	eef1a1l2	-1.20	3.80	38.1 0	1.73E- 59	2.35E- 06	3.31E- 04
g137 01	XP_002665 852	1003304 35	apobb.2	-1.62	2.93	37.5 2	4.70E- 49	2.55E- 06	3.54E- 04
g325	XP_009299 903	393246	ncor1	0.61	7.49	37.5 1	2.78E- 159	2.65E- 06	3.61E- 04
g186 98	AAQ97835	393922	psmd13	-0.93	5.10	37.3 3	0.00E+ 00	2.78E- 06	3.74E- 04
g244 09	AAT68067	378848	hsa5	-1.36	3.55	37.2 1	0.00E+ 00	3.44E- 06	4.54E- 04
g163 72	XP_021332 635	1005357 13	LOC1005357 13	-0.72	5.67	37.0 6	0.00E+ 00	3.50E- 06	4.56E- 04
g245 40	AAH97172	327642	lrrc8aa	-1.04	4.28	36.9 3	1.45E- 46	3.57E- 06	4.58E- 04
g282 7	XP_699339	570735	dhx30	-1.37	3.36	36.6 0	7.83E- 175	3.92E- 06	4.96E- 04
g265 90	NP_694486	259306	cyp2ad2	0.61	9.22	36.5 9	0.00E+ 00	4.73E- 06	5.89E- 04
g222 83	XP_005158 606	556848	crybg1a	1.01	4.79	36.3 1	4.36E- 59	5.65E- 06	6.94E- 04
g168 32	ACR83585	407086	junba	0.99	7.59	36.2 8	1.51E- 134	5.83E- 06	7.05E- 04
g181 01	AAH53243	393997	lipca	0.73	6.30	36.1 6	0.00E+ 00	7.08E- 06	8.46E- 04
g172 88	AAH71313	415138	akr1b1.1	0.73	4.96	36.0 2	1.40E+ 00	7.86E- 06	9.26E- 04
g274 88	AAH76027	436717	decr1	0.68	7.05	35.8 8	4.38E- 27	8.33E- 06	9.67E- 04
g107 1	AAH81582	447800	serpinf1	0.88	6.72	35.8 2	0.00E+ 00	8.44E- 06	9.67E- 04
g252 15	NP_956951	393630	thyn1	-1.01	4.08	35.2 3	4.41E- 24	9.74E- 06	1.10E- 03
g694 3	CAB64946	30314	apoeb	-1.33	3.45	35.2 3	6.73E- 149	1.01E- 05	1.13E- 03
g258 70	AAI24635	569462	pnkp	0.64	6.34	35.0 2	0.00E+ 00	1.04E- 05	1.14E- 03
g317 5	AAH49037	406736	mgst3a	-0.86	5.55	34.6 6	4.42E- 149	1.11E- 05	1.21E- 03
g202 87	AAU14811	402991	gadd45ga	1.16	3.68	34.1 1	0.00E+ 00	1.32E- 05	1.42E- 03
g970 5	XP_021326 410	553550	tsen54	-1.92	3.25	33.5 8	5.85E- 151	1.48E- 05	1.55E- 03
g199 02	AAI25920	777717	bmp3	0.91	5.13	33.5 1	0.00E+ 00	1.48E- 05	1.55E- 03
g183 97	NP_001139 486	555721	sall1a	0.65	8.92	33.4 3	2.80E+ 00	1.51E- 05	1.56E- 03
g147 50	AAH93358	550453	pdr1	-1.36	3.42	33.4 4	2.19E- 130	1.72E- 05	1.76E- 03
g264 51	AAH48051	321193	paics	0.69	7.47	33.2 8	0.00E+ 00	1.77E- 05	1.79E- 03
g258 69	AAI39710	563828	si:ch1073- 280e3.1	0.63	6.03	33.2 8	1.26E- 52	1.84E- 05	1.84E- 03

A. SUPPLEMENTARY MATERIAL

g956	AAH47807	327133	hyou1	-0.71	5.61	32.9 9	0.00E+ 00	2.18E- 05	2.15E- 03
g879	XP_002663 594	378983	postna	1.10	3.08	32.8 3	2.36E- 141	2.21E- 05	2.15E- 03
g246 48	XP_021331 726	404040	dab2	-0.72	4.86	32.8 3	2.11E- 142	2.22E- 05	2.15E- 03
g200 74	AAH57478	321621	gldc	-0.84	4.12	32.7 2	0.00E+ 00	2.34E- 05	2.22E- 03
g111 53	XP_005156 656	492518	haao	0.68	5.62	32.6 9	0.00E+ 00	2.37E- 05	2.22E- 03
g253 56	XP_009289 773	559212	gbe1b	-1.00	3.29	32.1 2	0.00E+ 00	2.37E- 05	2.22E- 03
g136 54	AAH60895	335799	sod2	-0.96	4.40	32.0 3	2.09E- 80	2.48E- 05	2.29E- 03
g697 2	XP_005157 915	553294	ube3d	-1.27	2.60	31.8 4	1.86E- 41	2.51E- 05	2.30E- 03
g258 00	NP_001107 918	573117	tbrg1	-1.01	3.12	31.4 8	9.55E- 10	2.62E- 05	2.36E- 03
g450 3	F1QGH9	322265	psmd11b	-1.31	2.45	31.4 0	1.03E- 146	2.64E- 05	2.36E- 03
g267 44	ABF18955	393108	cyp2p6	-0.75	5.91	30.5 4	0.00E+ 00	2.90E- 05	2.57E- 03
g810 2	AAG25710	140615	fads2	-2.88	2.25	30.4 3	0.00E+ 00	2.98E- 05	2.61E- 03
g164 03	XP_001921 144	799340	si:ch211- 207j7.2	0.87	6.03	30.2 9	0.00E+ 00	3.02E- 05	2.63E- 03
g234 8	AAH75752	436592	stard3nl	-0.61	7.59	30.2 2	3.88E- 144	3.22E- 05	2.77E- 03
g105 78	AAQ94592	321664	aldob	0.61	12.8 1	30.2 4	0.00E+ 00	3.31E- 05	2.82E- 03
g549 0	CAK04314	338267	arl4ab	0.97	3.73	30.1 6	1.10E- 76	3.50E- 05	2.93E- 03
g145	ABA54856	566735	insb	0.85	5.84	30.0 8	0.00E+ 00	3.51E- 05	2.93E- 03
g243 24	AAH92881	550498	iscub	-1.14	4.92	29.9 1	0.00E+ 00	4.20E- 05	3.47E- 03
g170 79	AAH53187	393941	psmc2	-0.60	7.80	29.7 9	2.19E- 61	4.28E- 05	3.50E- 03
g232 32	AAI62754	1000058 54	zgc:194242	0.87	4.30	29.6 3	2.29E- 85	4.50E- 05	3.65E- 03
g211 32	AAI53957	561171	cbln8	-1.16	11.6 0	29.5 0	2.99E- 64	4.63E- 05	3.72E- 03
g155 9	AAH71315	321611	mpc2	-0.70	6.32	29.4 0	0.00E+ 00	4.89E- 05	3.89E- 03
g159 59	AAI25969	558851	cyp4v8	0.84	5.61	29.2 2	7.14E- 56	5.30E- 05	4.18E- 03
g234 24	AAH65350	402927	zgc:77375	-1.50	2.17	29.1 5	0.00E+ 00	5.80E- 05	4.50E- 03
g506 6	ACX31197	560713	ccn4a	-1.27	2.68	28.9 8	5.68E- 130	5.82E- 05	4.50E- 03
g137 75	XP_021322 709	1005365 00	LOC1005365 00	-1.92	4.34	28.8 1	3.40E- 149	6.08E- 05	4.66E- 03
g215 31	AAH79506	445316	cdkl1	0.83	4.33	28.7 8	0.00E+ 00	6.94E- 05	5.28E- 03
g973 6	AAH81493	447942	cox6a2	0.65	7.70	28.7 0	0.00E+ 00	7.04E- 05	5.31E- 03
g492 1	BAF47195	564619	gstr	-0.93	8.82	28.5 7	2.69E- 46	7.44E- 05	5.56E- 03
g633 8	NP_001313 526	792958	LOC792958	-0.84	5.26	28.2 8	1.70E- 62	7.64E- 05	5.66E- 03

g160 22	ABA54452	326102	hhp	0.73	4.57	28.2 2	3.10E- 132	7.75E- 05	5.69E- 03
g176 58	AAI65335	114367	nots	-1.41	2.79	28.2 1	1.07E- 123	8.01E- 05	5.84E- 03
g106 91	XP_021336 737	767699	pmt	0.61	8.26	28.1 6	1.70E- 145	8.68E- 05	6.27E- 03
g151 33	AAI33741	1000047 00	cyp24a1	0.89	6.22	28.1 4	0.00E+ 00	9.20E- 05	6.59E- 03
g438 0	Q6PBP3	323739	selenou1a	-0.90	7.74	27.8 3	2.78E- 133	9.29E- 05	6.60E- 03
g919 3	AAH71424	405830	lgals2a	-0.79	6.78	27.7 3	2.55E- 99	9.48E- 05	6.68E- 03
g211 31	XP_021322 670	557984	si:ch211- 215k15.4	0.63	11.3 4	27.6 3	0.00E+ 00	9.86E- 05	6.89E- 03
g258 5	XP_021335 998	1104401 42	LOC1104401 42	-1.12	3.77	27.5 9	0.00E+ 00	1.12E- 04	7.76E- 03
g959 2	XP_005164 296	1018828 99	LOC1018828 99	0.60	6.04	27.4 7	0.00E+ 00	1.29E- 04	8.77E- 03
g183 49	AAH93190	550462	ubl7a	-1.26	2.22	27.1 4	0.00E+ 00	1.29E- 04	8.77E- 03
g853 2	AAH76206	445033	psma1	-0.73	5.20	26.8 4	0.00E+ 00	1.30E- 04	8.77E- 03
g823 0	AAH65322	406571	utp4	0.79	3.99	26.8 0	0.00E+ 00	1.32E- 04	8.83E- 03
g172 16	XP_003200 166	1005378 09	LOC1005378 09	-1.26	2.58	26.6 7	1.95E- 110	1.38E- 04	9.19E- 03
g835 3	AAX48021	415191	tnni4b.2	-0.99	4.92	26.0 3	8.75E- 25	1.39E- 04	9.19E- 03
g266 13	XP_005170 499	373089	tjp3	0.61	5.18	25.9 7	9.01E- 25	1.46E- 04	9.59E- 03
g239 31	AAI35051	64278	psmb7	-0.61	6.72	25.8 3	1.08E- 21	1.50E- 04	9.76E- 03
g166 07	XP_002660 623	1003314 28	c3a.4	0.63	8.42	25.8 3	0.00E+ 00	1.51E- 04	9.78E- 03
g215 16	NP_001177 311	1004629 64	selenol	0.90	4.01	25.7 1	0.00E+ 00	1.53E- 04	9.83E- 03
g87	AAH44395	406828	faxdc2	-0.75	5.78	25.6 3	0.00E+ 00	1.55E- 04	9.86E- 03
g140 5	AAH56563	327588	hpxa	0.82	10.5 2	25.5 1	0.00E+ 00	1.70E- 04	1.07E- 02
g262 83	AAH68336	30248	calr3a	-0.77	4.34	25.3 4	2.61E- 104	1.71E- 04	1.07E- 02
g895	XP_021335 093	562316	kl	1.09	3.00	25.1 7	0.00E+ 00	1.81E- 04	1.13E- 02
g137 90	AAI63181	792608	knstrn	-1.53	2.60	24.8 7	0.00E+ 00	1.83E- 04	1.13E- 02
g274 4	AAH56520	79378	agxtb	0.62	9.65	24.7 6	0.00E+ 00	1.86E- 04	1.14E- 02
g199 88	AAL75953	140614	xbp1	0.63	5.41	24.5 9	0.00E+ 00	1.94E- 04	1.18E- 02
g248 93	XP_005172 658	1018865 67	rph3aa	0.58	7.32	24.6 0	5.24E- 172	2.08E- 04	1.26E- 02
g621 2	AAI60631	1001445 69	dpys	-0.76	4.63	24.5 9	0.00E+ 00	2.14E- 04	1.29E- 02
g111 10	XP_021336 675	436640	slc25a48	-1.03	3.45	24.5 1	8.77E- 153	2.16E- 04	1.29E- 02
g115 85	AAN77156	393425	elov15	-1.10	2.74	24.4 8	1.51E- 88	2.18E- 04	1.29E- 02
g136 95	XP_017207 922	321070	rock2b	-0.75	4.06	24.4 1	0.00E+ 00	2.19E- 04	1.29E- 02

g240 33	XP_009300 100	1003317 45	npr2	0.67	4.55	24.3 4	0.00E+ 00	2.47E- 04	1.44E- 02
g202 45	AAH91862	541493	lifra	0.71	4.13	24.2 0	1.54E- 147	2.53E- 04	1.46E- 02
g166 18	XP_001345 197	1000064 64	si:dkey-27j5.5	-1.10	2.89	23.9 7	0.00E+ 00	2.53E- 04	1.46E- 02
g102 96	XP_005157 299	563853	mtus1a	-0.70	4.41	23.9 0	2.03E- 51	2.61E- 04	1.49E- 02
g268 27	XP_005166 980	795505	eps15	-0.62	5.18	23.8 9	0.00E+ 00	2.74E- 04	1.56E- 02
g135 34	AAH65466	394198	fosab	1.24	3.87	23.8 3	0.00E+ 00	2.83E- 04	1.60E- 02
g262 16	Q5U3U3	1000057 17	cpt2	-1.22	2.52	23.5 8	0.00E+ 00	2.98E- 04	1.68E- 02
g107 2	AAI28865	563663	serpinf2b	0.52	9.70	23.5 6	3.64E- 130	3.01E- 04	1.68E- 02
g126 0	AAH62829	406531	arl4cb	0.99	3.52	23.5 5	1.29E- 84	3.16E- 04	1.75E- 02
g175 19	XP_005164 939	563056	hcls1	-0.64	6.28	23.5 0	0.00E+ 00	3.26E- 04	1.80E- 02
g188 74	XP_021333 072	322372	fah	-0.95	3.11	23.4 6	1.19E- 168	3.29E- 04	1.80E- 02
g191 21	XP_005165 919	393122	dnpep	-0.75	5.00	23.2 9	1.38E- 95	3.32E- 04	1.81E- 02
g240 76	XP_021327 915	558050	si:ch211- 127i16.2	-1.04	2.79	23.1 2	2.06E- 116	3.38E- 04	1.83E- 02
g187 79	AAS58450	322626	hsd17b12b	-0.73	3.89	23.0 9	6.73E- 164	3.62E- 04	1.95E- 02
g182 00	AAI60629	799650	ddx21	0.47	12.0 7	23.0 4	0.00E+ 00	3.84E- 04	2.04E- 02
g147 35	AAI17590	724002	acad9	0.58	5.56	22.9 8	1.01E- 101	3.85E- 04	2.04E- 02
g165 10	AAH42325	321898	psmd12	-0.73	4.25	22.9 0	1.42E- 124	3.87E- 04	2.04E- 02
g148 88	AJG05938	767691	manf	-1.13	2.55	22.8 0	3.32E- 103	3.97E- 04	2.08E- 02
g219 1	AAH81413	447937	zgc:101851	-1.23	2.27	22.7 9	4.00E- 03	4.00E- 04	2.08E- 02
g213 16	AAI52167	553553	cfid	0.98	3.82	22.7 7	4.59E- 83	4.06E- 04	2.10E- 02
g193 48	AAI16550	692285	pspc1	0.57	5.43	22.7 5	1.02E- 103	4.13E- 04	2.12E- 02
g961 9	NP_001315 082	1079902 73	si:dkey- 114l24.2	-0.98	4.04	22.6 2	0.00E+ 00	4.15E- 04	2.13E- 02
g229 62	AAI24536	556929	zgc:154075	0.96	3.89	22.6 1	0.00E+ 00	4.19E- 04	2.13E- 02
g155 89	XP_005155 982	664760	tbce	0.73	4.18	22.5 6	3.03E- 110	4.22E- 04	2.14E- 02
g204 88	NP_001119 925	793882	git2a	-1.10	2.64	22.5 7	0.00E+ 00	4.27E- 04	2.15E- 02
g814	AAI15276	678552	f9b	0.47	9.04	22.5 8	0.00E+ 00	4.29E- 04	2.15E- 02
g721 3	XP_005158 307	558292	scap	-1.03	2.73	22.3 2	0.00E+ 00	4.38E- 04	2.18E- 02
g262 22	NP_001038 663	570393	hook1	-1.17	2.74	22.2 4	0.00E+ 00	4.43E- 04	2.19E- 02
g219 59	XP_002660 625	1003201 76	si:ch73- 343l4.8	0.50	8.00	22.0 8	1.24E- 52	4.63E- 04	2.28E- 02
g260 15	AAH56516	393518	psmd2	-0.47	7.85	22.0 6	1.10E- 90	4.66E- 04	2.28E- 02

g233 54	AAF63256	791729	rh50	-1.09	3.45	21.8 1	0.00E+ 00	5.02E- 04	2.44E- 02
g851 4	AAH75928	436610	pnpla3	0.86	3.94	21.7 3	0.00E+ 00	5.15E- 04	2.47E- 02
g280 46	XP_021333 686	555504	pcdh7b	0.83	3.83	21.7 1	0.00E+ 00	5.15E- 04	2.47E- 02
g242 09	XP_021331 766	445388	rcl1	0.55	8.36	21.7 0	0.00E+ 00	5.22E- 04	2.49E- 02
g168 87	AAH49010	573095	psmb3	-0.55	6.61	21.6 6	3.18E- 57	5.26E- 04	2.50E- 02
g147 51	ACG75896 83	1005266 83	aldh1l1	-0.60	7.05	21.6 0	0.00E+ 00	5.38E- 04	2.54E- 02
g237 2	XP_009295 541	557738	ccny	0.59	4.74	21.5 5	1.92E- 57	5.42E- 04	2.54E- 02
g175 60	Q7SXX5	402826	si:ch73- 71d17.1	0.76	3.68	21.5 3	0.00E+ 00	5.51E- 04	2.57E- 02
g175 82	NP_001314 754	406280	itih2	0.51	11.7 0	21.5 2	1.12E- 103	5.66E- 04	2.63E- 02
g143 39	CAD60658	368823	fkbp11	0.55	7.85	21.5 1	0.00E+ 00	5.76E- 04	2.66E- 02
g215 32	AAH46029	393187	dmac2l	0.84	4.12	21.4 8	0.00E+ 00	6.12E- 04	2.81E- 02
g330 5	AAI50400	566527	dad1	-0.72	5.72	21.3 8	0.00E+ 00	6.16E- 04	2.81E- 02
g252 74	XP_005157 459	1001260 24	zgc:174895	-1.22	3.23	21.3 6	2.64E- 64	6.19E- 04	2.81E- 02
g231 39	XP_005162 282	30634	llcama	0.94	3.52	21.3 5	1.80E- 02	6.34E- 04	2.87E- 02
g117 34	P59723	373126	hif1an	-1.15	2.88	21.2 0	0.00E+ 00	6.62E- 04	2.98E- 02
g184 35	BAB90841	140634	cyp1a	-0.84	3.98	21.2 0	0.00E+ 00	6.77E- 04	3.03E- 02
g608 9	AAF66961	58122	mcl1a	0.47	8.16	21.1 1	0.00E+ 00	7.05E- 04	3.14E- 02
g241	AAH46072	266797	stt3a	-0.63	5.36	21.0 9	1.01E- 169	7.23E- 04	3.21E- 02
g215 8	AAH64707	406314	ubxn4	-0.52	5.97	20.9 1	2.90E+ 00	7.49E- 04	3.30E- 02
g217 94	BAD67593	335195	ywhaqb	0.48	7.18	20.8 1	2.33E- 83	7.55E- 04	3.31E- 02
g202 39	AAI09473	322228	bhmt	0.53	12.0 9	20.7 2	0.00E+ 00	7.81E- 04	3.41E- 02
g137 30	XP_017207 953	436657	cpsf2	-0.72	4.02	20.6 5	8.25E- 84	7.92E- 04	3.44E- 02
g123 55	AAF05817	30647	psme2	-0.64	5.50	20.5 8	4.07E- 90	8.01E- 04	3.46E- 02
g197 66	XP_002667 717	1003328 41	LOC1003328 41	-1.10	2.85	20.5 5	0.00E+ 00	8.08E- 04	3.47E- 02
g145 73	AAH54602	393902	rab7a	-0.69	3.92	20.5 1	0.00E+ 00	8.14E- 04	3.48E- 02
g517 8	NP_001093 492	566469	si:dkeyp- 120h9.1	-0.95	2.74	20.4 5	0.00E+ 00	8.22E- 04	3.50E- 02
g175 59	NP_001038 575	566596	si:dkey- 153k10.9	0.69	4.22	20.3 7	0.00E+ 00	8.40E- 04	3.54E- 02
g871 6	XP_009292 028	793125	si:dkey- 47k20.3	-0.83	5.46	20.3 5	9.97E- 118	8.44E- 04	3.54E- 02
g973 3	XP_001924 016	569097	LOC569097	0.76	3.18	20.2 7	1.45E- 178	8.44E- 04	3.54E- 02
g166 5	AAH65443	322544	dnajc3a	-0.75	4.19	20.2 6	0.00E+ 00	8.53E- 04	3.55E- 02

g249 5	NP_001264 046	560020	acss2l	-0.73	4.36	20.2 5	3.53E- 35	8.55E- 04	3.55E- 02
g775 1	XP_021329 551	797251	si:ch211- 236l14.4	-0.68	3.97	20.2 3	0.00E+ 00	8.59E- 04	3.55E- 02
g315 9	AAH45968	327057	uck2b	0.57	4.80	20.1 0	4.61E- 60	8.93E- 04	3.67E- 02
g215 61	A1L2F3	567446	nusap1	-0.91	3.58	19.9 9	0.00E+ 00	9.11E- 04	3.73E- 02
g258 58	CAA74004	30355	apoal1a	0.64	12.3 1	19.8 7	0.00E+ 00	9.16E- 04	3.73E- 02
g163 99	NP_001121 843	1001478 56	illr2	0.47	11.9 5	19.7 6	0.00E+ 00	9.19E- 04	3.73E- 02
g200 60	XP_005166 846	335159	acsb2g	-0.66	5.11	19.7 6	0.00E+ 00	9.50E- 04	3.83E- 02
g231 01	AA4Y40842	565685	fance	0.69	4.97	19.7 7	0.00E+ 00	9.54E- 04	3.83E- 02
g358 2	AAH83309	449786	polr3glb	-1.07	2.24	19.6 4	5.97E- 62	9.57E- 04	3.83E- 02
g125 96	XP_005162 731	322795	csmp1b	0.50	6.69	19.6 4	7.65E- 06	9.68E- 04	3.85E- 02
g163 98	XP_021336 864	1000030 29	fgl1	0.48	12.3 5	19.5 0	2.09E- 79	9.88E- 04	3.92E- 02
g244 77	AAH68340	286746	sdad1	0.47	6.94	19.4 6	0.00E+ 00	1.01E- 03	3.97E- 02
g926 1	ABI33876	793274	srebf1	-1.21	2.31	19.3 7	9.17E- 20	1.01E- 03	3.97E- 02
g264 94	AAI15130	555526	si:ch211- 201h21.5	0.67	4.58	19.3 6	0.00E+ 00	1.03E- 03	4.02E- 02
g302 3	NP_001243 132	492647	angptl4	0.56	6.53	19.1 7	7.01E- 86	1.04E- 03	4.05E- 02
g228 52	XP_009295 191	564967	zbtb17	0.77	3.37	19.0 1	0.00E+ 00	1.05E- 03	4.05E- 02
g277	AAI63387	566852	pdzd3b	0.69	4.25	18.9 9	0.00E+ 00	1.07E- 03	4.10E- 02
g121 86	NP_956936	393615	zgc:64201	-1.32	2.36	18.7 9	0.00E+ 00	1.07E- 03	4.10E- 02
g74	AAX83257	378966	ncf1	1.12	2.29	18.7 6	7.61E- 33	1.08E- 03	4.11E- 02
g208 81	NP_001116 221	563963	dgr8	-0.62	4.68	18.6 8	1.83E- 61	1.09E- 03	4.15E- 02
g175 35	AAH45345	322605	chpt1	0.68	5.69	18.6 8	1.55E- 124	1.10E- 03	4.18E- 02
g121 25	AAH60926	393807	ppp1r3cb	0.82	3.36	18.6 8	0.00E+ 00	1.12E- 03	4.22E- 02
g972 5	XP_009297 793	327462	hsd3b7	-0.67	4.72	18.5 7	0.00E+ 00	1.24E- 03	4.64E- 02
g122 29	AAH53263	406409	cyb5a	0.66	5.07	18.5 6	3.07E- 67	1.24E- 03	4.64E- 02
g430 5	AAH95058	554142	rgrb	0.75	3.82	18.5 5	1.18E- 83	1.25E- 03	4.64E- 02
g685 0	NP_001116 330	560277	masp2	0.47	11.2 2	18.5 4	3.28E- 59	1.25E- 03	4.64E- 02
g632 2	ABG35925	777792	sult2st3	-1.04	2.53	18.5 2	2.96E- 10	1.26E- 03	4.64E- 02
g154 83	XP_005173 986	1018823 55	LOC1018823 55	-0.82	3.24	18.4 8	2.29E- 123	1.26E- 03	4.64E- 02
g809 5	AAH54696	337225	chid1	-0.92	2.76	18.4 7	0.00E+ 00	1.28E- 03	4.69E- 02
g167 04	NP_001186 962	794625	atp5md	-0.67	6.64	18.4 0	8.56E- 157	1.34E- 03	4.87E- 02

g144 38	AAI42794	553214	EIF4B	0.43	7.42	18.3 9	0.00E+ 00	1.34E- 03	4.87E- 02
g650 7	XP_009304 581	558046	TRIM2B	-1.09	2.27	18.3 1	0.00E+ 00	1.36E- 03	4.90E- 02

Supplementary Table 4.4: Gene Ontology (GO) annotation for up- and down-regulated pathways for the pituitary gland (p -value < 0.01). ID refers to the universal identifiers of the GO terms, the Ont column shows the ontology domain that each GO term belongs to. The three domains are: biological process (BP), cellular component (CC) and molecular function (MF). The N column represents the total number of genes annotated with each GO term. The Up and Down columns indicate the number of genes within the GO term that are significantly up- and down-regulated in high versus low treatment, respectively.

ID	N	Up	Down	p -value Up	p -value Down	Ont	Term
GO:0060319	10	0	3	1.00	9.57E-05	BP	primitive erythrocyte differentiation
GO:0002262	57	1	5	0.37	1.98E-04	BP	myeloid cell homeostasis
GO:0005737	2845	20	43	0.78	2.19E-04	CC	cytoplasm
GO:0048872	63	1	5	0.39	3.18E-04	BP	homeostasis of number of cells
GO:0043295	4	0	2	1.00	5.33E-04	MF	glutathione binding
GO:1900750	4	0	2	1.00	5.33E-04	MF	oligopeptide binding
GO:0030218	51	0	4	1.00	1.37E-03	BP	erythrocyte differentiation
GO:0034101	52	0	4	1.00	1.47E-03	BP	erythrocyte homeostasis
GO:0060215	28	0	3	1.00	2.31E-03	BP	primitive hemopoiesis
GO:0044444	1934	12	30	0.87	2.57E-03	CC	cytoplasmic part
GO:0042246	64	1	4	0.40	3.17E-03	BP	tissue regeneration
GO:0009311	10	0	2	1.00	3.85E-03	BP	oligosaccharide metabolic process
GO:0006898	35	0	3	1.00	4.40E-03	BP	receptor-mediated endocytosis
GO:0051084	11	0	2	1.00	4.68E-03	BP	'de novo' posttranslational protein folding
GO:0051085	11	0	2	1.00	4.68E-03	BP	chaperone cofactor-dependent protein refolding
GO:0006458	12	0	2	1.00	5.58E-03	BP	'de novo' protein folding
GO:0004364	12	0	2	1.00	5.58E-03	MF	glutathione transferase activity
GO:0035162	40	0	3	1.00	6.41E-03	BP	embryonic hemopoiesis
GO:0044092	80	1	4	0.47	7.03E-03	BP	negative regulation of molecular function
GO:0009611	127	2	5	0.27	7.17E-03	BP	response to wounding
GO:0045765	42	0	3	1.00	7.35E-03	BP	regulation of angiogenesis
GO:0070988	14	0	2	1.00	7.59E-03	BP	demethylation

GO:0040007	185	3	6	0.18	8.30E-03	BP	growth
GO:0061077	15	0	2	1.00	8.71E-03	BP	chaperone-mediated protein folding
GO:0045766	15	0	2	1.00	8.71E-03	BP	positive regulation of angiogenesis
GO:1904018	15	0	2	1.00	8.71E-03	BP	positive regulation of vasculature development
GO:0030099	86	0	4	1.00	9.04E-03	BP	myeloid cell differentiation
GO:0030729	1	0	1	1.00	9.54E-03	MF	acetoacetate-CoA ligase activity
GO:0007195	1	0	1	1.00	9.54E-03	BP	adenylate cyclase-inhibiting dopamine receptor signaling pathway
GO:0004557	1	0	1	1.00	9.54E-03	MF	alpha-galactosidase activity
GO:1904158	1	0	1	1.00	9.54E-03	BP	axonemal central apparatus assembly
GO:0004952	1	0	1	1.00	9.54E-03	MF	dopamine neurotransmitter receptor activity
GO:0001591	1	0	1	1.00	9.54E-03	MF	dopamine neurotransmitter receptor activity, coupled via Gi/Go
GO:0071603	1	0	1	1.00	9.54E-03	BP	endothelial cell-cell adhesion
GO:0030228	1	0	1	1.00	9.54E-03	MF	lipoprotein particle receptor activity
GO:0005041	1	0	1	1.00	9.54E-03	MF	low-density lipoprotein particle receptor activity
GO:1990124	1	0	1	1.00	9.54E-03	CC	messenger ribonucleoprotein complex
GO:1903966	1	0	1	1.00	9.54E-03	BP	monounsaturated fatty acid biosynthetic process
GO:1903964	1	0	1	1.00	9.54E-03	BP	monounsaturated fatty acid metabolic process
GO:1903170	1	0	1	1.00	9.54E-03	BP	negative regulation of calcium ion transmembrane transport
GO:1901020	1	0	1	1.00	9.54E-03	BP	negative regulation of calcium ion transmembrane transporter activity
GO:2001258	1	0	1	1.00	9.54E-03	BP	negative regulation of cation channel activity
GO:2000766	1	0	1	1.00	9.54E-03	BP	negative regulation of cytoplasmic translation
GO:1901386	1	0	1	1.00	9.54E-03	BP	negative regulation of voltage-gated calcium channel activity
GO:0032896	1	0	1	1.00	9.54E-03	MF	palmitoyl-CoA 9-desaturase activity
GO:0004615	1	0	1	1.00	9.54E-03	MF	phosphomannomutase activity

A. SUPPLEMENTARY MATERIAL

GO:0090118	1	0	1	1.00	9.54E-03	BP	receptor-mediated endocytosis involved in cholesterol transport
GO:0004768	1	0	1	1.00	9.54E-03	MF	stearoyl-CoA 9-desaturase activity
GO:0045098	1	0	1	1.00	9.54E-03	CC	type III intermediate filament
GO:0072341	16	0	2	1.00	9.89E-03	MF	modified amino acid binding
GO:1901342	47	0	3	1.00	1.00E-02	BP	regulation of vasculature development
GO:0042403	3	2	0	1.84E-04	1.00	BP	thyroid hormone metabolic process
GO:0051129	68	5	0	1.89E-04	1.00	BP	negative regulation of cellular component organization
GO:0008585	4	2	0	3.66E-04	1.00	BP	female gonad development
GO:0032878	4	2	0	3.66E-04	1.00	BP	regulation of establishment or maintenance of cell polarity
GO:0005179	21	3	1	5.69E-04	0.18	MF	hormone activity
GO:0046545	5	2	0	6.07E-04	1.00	BP	development of primary female sexual characteristics
GO:0046660	5	2	0	6.07E-04	1.00	BP	female sex differentiation
GO:0031492	5	2	0	6.07E-04	1.00	MF	nucleosomal DNA binding
GO:0016584	5	2	0	6.07E-04	1.00	BP	nucleosome positioning
GO:0048232	24	3	0	8.51E-04	1.00	BP	male gamete generation
GO:0007283	24	3	0	8.51E-04	1.00	BP	spermatogenesis
GO:0031936	6	2	0	9.06E-04	1.00	BP	negative regulation of chromatin silencing
GO:0060969	6	2	0	9.06E-04	1.00	BP	negative regulation of gene silencing
GO:0009755	25	3	0	9.61E-04	1.00	BP	hormone-mediated signaling pathway
GO:0008406	7	2	0	1.26E-03	1.00	BP	gonad development
GO:0045815	7	2	0	1.26E-03	1.00	BP	positive regulation of gene expression, epigenetic
GO:0007530	7	2	0	1.26E-03	1.00	BP	sex determination
GO:0045137	8	2	0	1.67E-03	1.00	BP	development of primary sexual characteristics
GO:0007064	8	2	0	1.67E-03	1.00	BP	mitotic sister chromatid cohesion
GO:0045910	8	2	0	1.67E-03	1.00	BP	negative regulation of DNA recombination
GO:0031935	8	2	0	1.67E-03	1.00	BP	regulation of chromatin silencing

GO:0018958	9	2	0	2.14E-03	1.00	BP	phenol-containing compound metabolic process
GO:0048608	9	2	0	2.14E-03	1.00	BP	reproductive structure development
GO:0061458	9	2	0	2.14E-03	1.00	BP	reproductive structure development
GO:0007548	9	2	0	2.14E-03	1.00	BP	sex differentiation
GO:0003006	35	3	0	2.59E-03	1.00	BP	developmental process involved in reproduction
GO:0048018	74	4	3	2.72E-03	0.03	MF	receptor ligand activity
GO:0030545	76	4	3	3.00E-03	0.04	MF	receptor regulator activity
GO:1905268	11	2	0	3.24E-03	1.00	BP	negative regulation of chromatin organization
GO:0040029	38	3	0	3.28E-03	1.00	BP	regulation of gene expression, epigenetic
GO:0030261	12	2	0	3.87E-03	1.00	BP	chromosome condensation
GO:0000018	12	2	0	3.87E-03	1.00	BP	regulation of DNA recombination
GO:0060968	12	2	0	3.87E-03	1.00	BP	regulation of gene silencing
GO:0051704	135	5	1	4.19E-03	0.73	BP	multi-organism process
GO:0008630	13	2	0	4.55E-03	1.00	BP	intrinsic apoptotic signaling pathway in response to DNA damage
GO:0007276	44	3	0	4.97E-03	1.00	BP	gamete generation
GO:0016458	44	3	0	4.97E-03	1.00	BP	gene silencing
GO:0048609	44	3	0	4.97E-03	1.00	BP	multicellular organismal reproductive process
GO:0005576	339	8	6	5.05E-03	0.10	CC	extracellular region
GO:0032504	45	3	0	5.30E-03	1.00	BP	multicellular organism reproduction
GO:0010639	46	3	0	5.64E-03	1.00	BP	negative regulation of organelle organization
GO:0140014	47	3	0	5.99E-03	1.00	BP	mitotic nuclear division
GO:0005102	213	6	3	6.58E-03	0.33	MF	signaling receptor binding
GO:0031491	16	2	0	6.89E-03	1.00	MF	nucleosome binding
GO:0031490	17	2	1	7.77E-03	0.15	MF	chromatin DNA binding
GO:0000786	17	2	0	7.77E-03	1.00	CC	nucleosome
GO:0007062	17	2	0	7.77E-03	1.00	BP	sister chromatid cohesion
GO:0003860	1	1	0	7.91E-03	1.00	MF	3-hydroxyisobutyryl-CoA hydrolase activity

A. SUPPLEMENTARY MATERIAL

GO:0003877	1	1	0	7.91E-03	1.00	MF	ATP adenylyltransferase activity
GO:0007414	1	1	0	7.91E-03	1.00	BP	axonal defasciculation
GO:0045575	1	1	0	7.91E-03	1.00	BP	basophil activation
GO:0002276	1	1	0	7.91E-03	1.00	BP	basophil activation involved in immune response
GO:0005432	1	1	0	7.91E-03	1.00	MF	calcium:sodium antiporter activity
GO:0071481	1	1	0	7.91E-03	1.00	BP	cellular response to X-ray
GO:0015960	1	1	0	7.91E-03	1.00	BP	diadenosine polyphosphate biosynthetic process
GO:0015966	1	1	0	7.91E-03	1.00	BP	diadenosine tetraphosphate biosynthetic process
GO:0015965	1	1	0	7.91E-03	1.00	BP	diadenosine tetraphosphate metabolic process
GO:0003347	1	1	0	7.91E-03	1.00	BP	epicardial cell to mesenchymal cell transition
GO:0070287	1	1	0	7.91E-03	1.00	MF	ferritin receptor activity
GO:0016913	1	1	0	7.91E-03	1.00	MF	follicle-stimulating hormone activity
GO:0016914	1	1	0	7.91E-03	1.00	CC	follicle-stimulating hormone complex
GO:0036230	1	1	0	7.91E-03	1.00	BP	granulocyte activation
GO:0043564	1	1	0	7.91E-03	1.00	CC	Ku70:Ku80 complex
GO:0004824	1	1	0	7.91E-03	1.00	MF	lysine-tRNA ligase activity
GO:0006430	1	1	0	7.91E-03	1.00	BP	lysyl-tRNA aminoacylation
GO:0042116	1	1	0	7.91E-03	1.00	BP	macrophage activation
GO:0048496	1	1	0	7.91E-03	1.00	BP	maintenance of animal organ identity
GO:1903392	1	1	0	7.91E-03	1.00	BP	negative regulation of adherens junction organization
GO:0001607	1	1	0	7.91E-03	1.00	MF	neuromedin U receptor activity
GO:0030728	1	1	0	7.91E-03	1.00	BP	ovulation
GO:0061696	1	1	0	7.91E-03	1.00	CC	pituitary gonadotropin complex
GO:0043032	1	1	0	7.91E-03	1.00	BP	positive regulation of macrophage activation
GO:0010893	1	1	0	7.91E-03	1.00	BP	positive regulation of steroid biosynthetic process

GO:0043030	1	1	0	7.91E-03	1.00	BP	regulation of macrophage activation
GO:0050810	1	1	0	7.91E-03	1.00	BP	regulation of steroid biosynthetic process
GO:0040040	1	1	0	7.91E-03	1.00	BP	thermosensory behavior
GO:0006590	1	1	0	7.91E-03	1.00	BP	thyroid hormone generation
GO:0102524	1	1	0	7.91E-03	1.00	MF	tRNAPhe (7-(3-amino-3-carboxypropyl)wyosine37-C2)-hydroxylase activity
GO:0031591	1	1	0	7.91E-03	1.00	BP	wybutosine biosynthetic process
GO:0031590	1	1	0	7.91E-03	1.00	BP	wybutosine metabolic process
GO:0043066	52	3	0	7.94E-03	1.00	BP	negative regulation of apoptotic process
GO:0019953	52	3	0	7.94E-03	1.00	BP	sexual reproduction
GO:0044703	53	3	0	8.36E-03	1.00	BP	multi-organism reproductive process
GO:0043069	53	3	0	8.36E-03	1.00	BP	negative regulation of programmed cell death
GO:0003684	18	2	0	8.69E-03	1.00	MF	damaged DNA binding
GO:0000049	18	2	0	8.69E-03	1.00	MF	tRNA binding
GO:0005615	228	6	4	9.07E-03	0.17	CC	extracellular space
GO:0006259	229	6	1	9.26E-03	0.89	BP	DNA metabolic process
GO:0044815	19	2	0	9.67E-03	1.00	CC	DNA packaging complex
GO:0051053	19	2	0	9.67E-03	1.00	BP	negative regulation of DNA metabolic process
GO:0006955	107	4	1	1.00E-02	0.64	BP	immune response

Supplementary Table 4. 5: Gene Ontology (GO) annotation for up- and down-regulated pathways for liver (p -value < 0.01). ID refers to the universal identifiers of the GO terms, the Ont column shows the ontology domain that each GO term belongs to. The three domains are: biological process (BP), cellular component (CC) and molecular function (MF). The N column represents the total number of genes annotated with each GO term. The Up and Down columns indicate the number of genes within the GO term that are significantly up- and down-regulated in high versus low treatment, respectively.

ID	N	Up	Down	p -value Up	p -value Down	Ont	Term
GO:1905369	42	1	24	0.90	1.37E-22	CC	endopeptidase complex
GO:0000502	42	1	24	0.90	1.37E-22	CC	proteasome complex
GO:1905368	54	1	24	0.95	3.38E-19	CC	peptidase complex
GO:0022624	21	0	14	1.00	5.21E-15	CC	proteasome accessory complex
GO:0005838	18	0	13	1.00	9.99E-15	CC	proteasome regulatory particle
GO:1901565	362	23	48	0.19	1.59E-12	BP	organonitrogen compound catabolic process
GO:0030163	255	10	38	0.86	1.18E-11	BP	protein catabolic process
GO:1901575	566	39	60	0.04	3.59E-11	BP	organic substance catabolic process
GO:0009056	657	45	65	0.04	9.96E-11	BP	catabolic process
GO:0043632	211	6	32	0.97	3.46E-10	BP	modification-dependent macromolecule catabolic process
GO:0044248	592	43	59	0.02	6.39E-10	BP	cellular catabolic process
GO:0005839	14	0	9	1.00	7.62E-10	CC	proteasome core complex
GO:0019941	206	6	31	0.96	8.05E-10	BP	modification-dependent protein catabolic process
GO:0051603	232	9	33	0.86	1.00E-09	BP	proteolysis involved in cellular protein catabolic process
GO:0044257	234	10	33	0.79	1.25E-09	BP	cellular protein catabolic process
GO:0006511	200	4	30	0.99	1.66E-09	BP	ubiquitin-dependent protein catabolic process
GO:0010499	15	1	9	0.55	1.83E-09	BP	proteasomal ubiquitin-independent protein catabolic process
GO:0004298	15	0	9	1.00	1.83E-09	MF	threonine-type endopeptidase activity
GO:0070003	15	0	9	1.00	1.83E-09	MF	threonine-type peptidase activity
GO:0003824	2768	133	171	0.90	6.45E-09	MF	catalytic activity
GO:0008540	9	0	7	1.00	8.65E-09	CC	proteasome regulatory particle, base subcomplex
GO:0009057	347	14	40	0.88	8.90E-09	BP	macromolecule catabolic process
GO:0005783	336	8	39	1.00	1.14E-08	CC	endoplasmic reticulum
GO:0008541	7	0	6	1.00	4.13E-08	CC	proteasome regulatory particle, lid subcomplex

GO:0010498	125	1	21	1.00	6.82E-08	BP	proteasomal protein catabolic process
GO:0043161	116	0	20	1.00	9.02E-08	BP	proteasome-mediated ubiquitin-dependent protein catabolic process
GO:0004576	8	0	6	1.00	1.59E-07	MF	oligosaccharyl transferase activity
GO:0044265	313	13	34	0.84	5.18E-07	BP	cellular macromolecule catabolic process
GO:0005737	2845	161	167	0.11	5.68E-07	CC	cytoplasm
GO:0004579	6	0	5	1.00	8.33E-07	MF	dolichyl-diphosphooligosaccharide-protein glycotransferase activity
GO:2000144	6	0	5	1.00	8.33E-07	BP	positive regulation of DNA-templated transcription, initiation
GO:0060261	6	0	5	1.00	8.33E-07	BP	positive regulation of transcription initiation from RNA polymerase II promoter
GO:0036402	6	0	5	1.00	8.33E-07	MF	proteasome-activating ATPase activity
GO:2000142	6	0	5	1.00	8.33E-07	BP	regulation of DNA-templated transcription, initiation
GO:0060260	6	0	5	1.00	8.33E-07	BP	regulation of transcription initiation from RNA polymerase II promoter
GO:0032787	114	11	18	0.04	1.54E-06	BP	monocarboxylic acid metabolic process
GO:0006508	458	30	42	0.12	2.29E-06	BP	proteolysis
GO:0016491	374	35	36	0.00	4.26E-06	MF	oxidoreductase activity
GO:0044444	1934	115	120	0.06	4.56E-06	CC	cytoplasmic part
GO:0055114	353	30	34	0.01	7.87E-06	BP	oxidation-reduction process
GO:0045899	5	0	4	1.00	1.63E-05	BP	positive regulation of RNA polymerase II transcriptional preinitiation complex assembly
GO:0045898	5	0	4	1.00	1.63E-05	BP	regulation of RNA polymerase II transcriptional preinitiation complex assembly
GO:0006631	67	7	12	0.06	2.38E-05	BP	fatty acid metabolic process
GO:0006082	297	26	29	0.01	2.91E-05	BP	organic acid metabolic process
GO:0008610	116	10	16	0.08	3.39E-05	BP	lipid biosynthetic process
GO:0008250	6	0	4	1.00	4.73E-05	CC	oligosaccharyltransferase complex
GO:0019773	6	0	4	1.00	4.73E-05	CC	proteasome core complex, alpha-subunit complex
GO:0005215	387	14	34	0.95	5.40E-05	MF	transporter activity
GO:0012505	733	19	54	1.00	5.42E-05	CC	endomembrane system

A. SUPPLEMENTARY MATERIAL

GO:0019752	278	19	27	0.14	6.04E-05	BP	carboxylic acid metabolic process
GO:0015711	75	4	12	0.56	7.59E-05	BP	organic anion transport
GO:0006629	282	21	27	0.06	7.72E-05	BP	lipid metabolic process
GO:0004364	12	1	5	0.47	8.87E-05	MF	glutathione transferase activity
GO:0043436	286	19	27	0.17	9.82E-05	BP	oxoacid metabolic process
GO:0006633	27	3	7	0.16	1.09E-04	BP	fatty acid biosynthetic process
GO:0044432	212	3	22	1.00	1.12E-04	CC	endoplasmic reticulum part
GO:0046942	47	2	9	0.71	1.47E-04	BP	carboxylic acid transport
GO:0015849	48	2	9	0.72	1.74E-04	BP	organic acid transport
GO:0016209	29	1	7	0.79	1.77E-04	MF	antioxidant activity
GO:0018196	8	0	4	1.00	2.06E-04	BP	peptidyl-asparagine modification
GO:0042623	164	3	18	0.99	2.33E-04	MF	ATPase activity, coupled
GO:0003674	6511	369	308	0.00	2.33E-04	MF	molecular_function
GO:1902494	560	14	42	1.00	2.69E-04	CC	catalytic complex
GO:0006820	111	4	14	0.84	2.75E-04	BP	anion transport
GO:0055085	324	10	28	0.98	3.27E-04	BP	transmembrane transport
GO:0022857	343	12	29	0.95	3.70E-04	MF	transmembrane transporter activity
GO:0042737	43	9	8	0.00	4.21E-04	BP	drug catabolic process
GO:0005789	189	3	19	1.00	4.87E-04	CC	endoplasmic reticulum membrane
GO:0043231	2943	144	157	0.85	4.93E-04	CC	intracellular membrane-bounded organelle
GO:0098827	190	3	19	1.00	5.20E-04	CC	endoplasmic reticulum subcompartment
GO:0005622	4340	234	218	0.25	5.37E-04	CC	intracellular
GO:0044424	4340	234	218	0.25	5.37E-04	CC	intracellular part
GO:0006694	25	3	6	0.14	5.39E-04	BP	steroid biosynthetic process
GO:0042175	191	3	19	1.00	5.55E-04	CC	nuclear outer membrane-endoplasmic reticulum membrane network
GO:0043248	10	0	4	1.00	5.77E-04	BP	proteasome assembly
GO:0042626	57	0	9	1.00	6.62E-04	MF	ATPase activity, coupled to transmembrane movement of substances
GO:0046943	36	1	7	0.86	7.32E-04	MF	carboxylic acid transmembrane transporter activity
GO:0031597	5	0	3	1.00	7.41E-04	CC	cytosolic proteasome complex
GO:0033764	5	1	3	0.24	7.41E-04	MF	steroid dehydrogenase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor
GO:0006811	344	11	28	0.97	8.50E-04	BP	ion transport
GO:0098656	37	0	7	1.00	8.70E-04	BP	anion transmembrane transport

GO:0005342	37	1	7	0.86	8.70E-04	MF	organic acid transmembrane transporter activity
GO:0006636	11	1	4	0.45	8.76E-04	BP	unsaturated fatty acid biosynthetic process
GO:0044255	215	19	20	0.02	9.40E-04	BP	cellular lipid metabolic process
GO:0006457	72	2	10	0.90	9.54E-04	BP	protein folding
GO:0015405	60	0	9	1.00	9.71E-04	MF	P-P-bond-hydrolysis-driven transmembrane transporter activity
GO:1905039	19	0	5	1.00	1.01E-03	BP	carboxylic acid transmembrane transport
GO:0022626	71	31	1	1.20E-21	0.96	CC	cytosolic ribosome
GO:0003735	123	36	0	4.16E-18	1.00	MF	structural constituent of ribosome
GO:0022625	39	21	0	1.97E-17	1.00	CC	cytosolic large ribosomal subunit
GO:0005840	145	37	1	2.09E-16	1.00	CC	ribosome
GO:0044445	108	32	4	2.16E-16	0.69	CC	cytosolic part
GO:0044391	119	32	1	4.76E-15	0.99	CC	ribosomal subunit
GO:0005198	252	42	9	1.25E-11	0.76	MF	structural molecule activity
GO:0015934	70	21	0	2.68E-11	1.00	CC	large ribosomal subunit
GO:0005615	228	38	7	1.25E-10	0.87	CC	extracellular space
GO:0005576	339	48	13	1.58E-10	0.71	CC	extracellular region
GO:0006412	286	43	6	2.23E-10	0.99	BP	translation
GO:0043043	289	43	6	3.13E-10	0.99	BP	peptide biosynthetic process
GO:0005829	468	58	30	3.82E-10	0.02	CC	cytosol
GO:0072376	20	11	1	7.67E-10	0.59	BP	protein activation cascade
GO:0043604	320	44	12	2.49E-09	0.73	BP	amide biosynthetic process
GO:0006518	335	45	9	3.43E-09	0.96	BP	peptide metabolic process
GO:0042273	46	15	0	5.34E-09	1.00	BP	ribosomal large subunit biogenesis
GO:0044421	260	38	8	5.94E-09	0.88	CC	extracellular region part
GO:0043603	396	49	17	1.06E-08	0.54	BP	cellular amide metabolic process
GO:1990904	390	48	11	1.81E-08	0.95	CC	ribonucleoprotein complex
GO:0006956	14	8	1	1.19E-07	0.46	BP	complement activation
GO:0046503	15	8	2	2.42E-07	0.13	BP	glycerolipid catabolic process
GO:1901566	597	60	26	4.90E-07	0.51	BP	organonitrogen compound biosynthetic process

A. SUPPLEMENTARY MATERIAL

GO:0005506	57	14	8	8.93E-07	0.00	MF	iron ion binding
GO:0022627	30	10	1	1.60E-06	0.73	CC	cytosolic small ribosomal subunit
GO:0006959	20	8	1	3.76E-06	0.59	BP	humoral immune response
GO:0042254	153	23	2	3.98E-06	0.99	BP	ribosome biogenesis
GO:0000027	16	7	0	7.68E-06	1.00	BP	ribosomal large subunit assembly
GO:0002252	43	11	2	8.78E-06	0.56	BP	immune effector process
GO:0004497	37	10	4	1.33E-05	0.07	MF	monooxygenase activity
GO:0046464	12	6	2	1.39E-05	0.09	BP	acylglycerol catabolic process
GO:0046461	12	6	2	1.39E-05	0.09	BP	neutral lipid catabolic process
GO:1901576	1553	116	62	1.63E-05	0.77	BP	organic substance biosynthetic process
GO:0042738	8	5	2	1.87E-05	0.04	BP	exogenous drug catabolic process
GO:0008395	8	5	2	1.87E-05	0.04	MF	steroid hydroxylase activity
GO:0019433	8	5	2	1.87E-05	0.04	BP	triglyceride catabolic process
GO:0044249	1512	113	58	2.14E-05	0.86	BP	cellular biosynthetic process
GO:0009058	1580	117	66	2.22E-05	0.63	BP	biosynthetic process
GO:0006639	25	8	3	2.56E-05	0.09	BP	acylglycerol metabolic process
GO:0006638	25	8	3	2.56E-05	0.09	BP	neutral lipid metabolic process
GO:0044271	1239	96	42	2.64E-05	0.97	BP	cellular nitrogen compound biosynthetic process
GO:0008152	3717	236	189	2.84E-05	0.00	BP	metabolic process
GO:0006641	19	7	3	2.95E-05	0.05	BP	triglyceride metabolic process
GO:0015935	49	11	1	3.35E-05	0.88	CC	small ribosomal subunit
GO:0006955	107	17	3	3.52E-05	0.85	BP	immune response
GO:0061134	58	12	4	3.53E-05	0.24	MF	peptidase regulator activity
GO:0045834	9	5	2	4.02E-05	0.05	BP	positive regulation of lipid metabolic process
GO:0022613	230	27	4	6.22E-05	0.99	BP	ribonucleoprotein complex biogenesis
GO:0007596	28	8	0	6.41E-05	1.00	BP	blood coagulation
GO:0006144	15	6	3	6.56E-05	0.02	BP	purine nucleobase metabolic process
GO:0042255	36	9	0	7.10E-05	1.00	BP	ribosome assembly
GO:0071704	3465	220	172	7.95E-05	0.01	BP	organic substance metabolic process

GO:0002181	45	10	2	8.37E-05	0.58	BP	cytoplasmic translation
GO:0007599	29	8	0	8.45E-05	1.00	BP	hemostasis
GO:0050878	29	8	0	8.45E-05	1.00	BP	regulation of body fluid levels
GO:1901564	2013	139	110	1.03E-04	0.00	BP	organonitrogen compound metabolic process
GO:0050817	30	8	0	1.10E-04	1.00	BP	coagulation
GO:0070887	360	36	17	1.22E-04	0.38	BP	cellular response to chemical stimulus
GO:0020037	47	10	4	1.24E-04	0.14	MF	heme binding
GO:0016712	11	5	3	1.35E-04	0.01	MF	oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen, reduced flavin or flavoprotein as one donor, and incorporation of one atom of oxygen
GO:0005577	3	3	0	1.42E-04	1.00	CC	fibrinogen complex
GO:1901998	3	3	0	1.42E-04	1.00	BP	toxin transport
GO:0005575	6350	364	292	1.50E-04	0.01	CC	cellular_component
GO:0009112	24	7	4	1.61E-04	0.02	BP	nucleobase metabolic process
GO:0050778	40	9	1	1.71E-04	0.83	BP	positive regulation of immune response
GO:1901652	40	9	1	1.71E-04	0.83	BP	response to peptide
GO:0043434	40	9	1	1.71E-04	0.83	BP	response to peptide hormone
GO:0004866	49	10	2	1.79E-04	0.63	MF	endopeptidase inhibitor activity
GO:0046113	7	4	2	2.27E-04	0.03	BP	nucleobase catabolic process
GO:0004252	60	11	0	2.34E-04	1.00	MF	serine-type endopeptidase activity
GO:0034645	1213	90	39	2.45E-04	0.98	BP	cellular macromolecule biosynthetic process
GO:0044242	51	10	6	2.53E-04	0.02	BP	cellular lipid catabolic process
GO:0030414	51	10	2	2.53E-04	0.65	MF	peptidase inhibitor activity
GO:0050776	51	10	1	2.53E-04	0.89	BP	regulation of immune response
GO:0046906	51	10	4	2.53E-04	0.18	MF	tetrapyrrole binding
GO:0009059	1218	90	39	2.83E-04	0.99	BP	macromolecule biosynthetic process
GO:0051346	34	8	1	2.84E-04	0.78	BP	negative regulation of hydrolase activity
GO:0002682	104	15	3	3.04E-04	0.83	BP	regulation of immune system process

A. SUPPLEMENTARY MATERIAL

GO:0042737	43	9	8	3.07E-04	0.00	BP	drug catabolic process
GO:0030073	13	5	0	3.45E-04	1.00	BP	insulin secretion
GO:0046883	13	5	0	3.45E-04	1.00	BP	regulation of hormone secretion
GO:0019843	27	7	0	3.61E-04	1.00	MF	rRNA binding
GO:0061135	54	10	4	4.11E-04	0.20	MF	endopeptidase regulator activity
GO:0002253	36	8	1	4.31E-04	0.80	BP	activation of immune response
GO:0034641	1917	130	70	4.33E-04	0.95	BP	cellular nitrogen compound metabolic process
GO:0003674	6511	369	308	4.71E-04	0.00	MF	molecular_function
GO:0043232	1023	77	36	4.88E-04	0.92	CC	intracellular non-membrane-bounded organelle
GO:0043228	1023	77	36	4.88E-04	0.92	CC	non-membrane-bounded organelle
GO:0030072	14	5	0	5.14E-04	1.00	BP	peptide hormone secretion
GO:0016491	374	35	36	5.37E-04	0.00	MF	oxidoreductase activity
GO:0010951	21	6	1	5.44E-04	0.60	BP	negative regulation of endopeptidase activity
GO:0016725	4	3	0	5.44E-04	1.00	MF	oxidoreductase activity, acting on CH or CH2 groups
GO:0006145	4	3	1	5.44E-04	0.16	BP	purine nucleobase catabolic process
GO:0048584	209	23	3	5.52E-04	0.99	BP	positive regulation of response to stimulus
GO:0009410	38	8	3	6.36E-04	0.22	BP	response to xenobiotic stimulus
GO:0002684	57	10	2	6.45E-04	0.71	BP	positive regulation of immune system process
GO:0072523	15	5	1	7.38E-04	0.48	BP	purine-containing compound catabolic process
GO:0006805	15	5	2	7.38E-04	0.13	BP	xenobiotic metabolic process
GO:0006577	9	4	0	7.50E-04	1.00	BP	amino-acid betaine metabolic process
GO:0050796	9	4	0	7.50E-04	1.00	BP	regulation of insulin secretion
GO:0052548	39	8	2	7.64E-04	0.51	BP	regulation of endopeptidase activity
GO:0044238	3305	205	158	8.17E-04	0.05	BP	primary metabolic process
GO:0006807	3120	195	143	8.25E-04	0.18	BP	nitrogen compound metabolic process
GO:0048518	816	63	26	8.90E-04	0.96	BP	positive regulation of biological process
GO:0017171	70	11	0	9.23E-04	1.00	MF	serine hydrolase activity
GO:0008236	70	11	0	9.23E-04	1.00	MF	serine-type peptidase activity

GO:0010466	23	6	1	9.26E-04	0.64	BP	negative regulation of peptidase activity
GO:0004867	23	6	2	9.26E-04	0.26	MF	serine-type endopeptidase inhibitor activity
GO:0046879	16	5	0	1.03E-03	1.00	BP	hormone secretion

Supplementary Table 4. 6: List of KEGG pathways for up- and down-regulated pathways for the pituitary gland. ID refers to the universal identifiers of the KEGG terms, the N column represents the total number of genes annotated with each KEGG pathway. The Up and Down columns indicate the number of genes within the KEGG pathway that are significantly up- and down-regulated in high versus low treatment, respectively

ID	ko	N	Up	Down	<i>p</i> - value Up	<i>p</i> -value Down	Pathway
dre04912	K05858	62	4	1	0.0014152	0.4492123	GnRH signaling pathway
dre04080	K05226	67	4	2	0.0018878	0.1338667	Neuroactive ligand-receptor interaction
dre04520	K05753	68	3	0	0.0164555	1	Adherens junction
dre04210	K02649	83	3	1	0.0277542	0.5504021	Apoptosis Adrenergic signaling in cardiomyocytes
dre04261	K05858	87	3	1	0.0313155	0.5674791	MAPK signaling pathway
dre04010	K05095	171	4	3	0.04608	0.2232692	Non-homologous end-joining
dre03450	K10884	6	1	0	0.0465484	1	Cell adhesion molecules
dre04514	K06088	48	2	0	0.0550964	1	Mitophagy - animal
dre04137	K08341	51	2	0	0.0613505	1	Cell cycle
dre04110	K06619	60	2	0	0.0814396	1	Oocyte meiosis
dre04114	K04441	62	2	2	0.0861485	0.1179447	Arachidonic acid metabolism
dre00590	K07418	20	1	0	0.1470196	1	Base excision repair
dre03410	K10798	23	1	0	0.1671516	1	beta-Alanine metabolism
dre00410	K00128	23	1	0	0.1671516	1	Propanoate metabolism
dre00640	K07514	27	1	0	0.1932676	1	Glutathione metabolism
dre00480	K00681	32	1	1	0.2247803	0.2645558	Aminoacyl-tRNA biosynthesis
dre00970	K01870	34	1	0	0.2370429	1	VEGF signaling pathway
dre04370	K02649	35	1	0	0.2431024	1	Protein processing in endoplasmic reticulum
dre04141	K10575	120	2	3	0.245355	0.1066185	Cysteine and methionine metabolism
dre00270	K08963	37	1	1	0.2550794	0.2991052	Lysosome
dre04142	K06501	90	0	5	1	0.0016286	Glyoxylate and dicarboxylate metabolism
dre00630	K01915	30	0	2	1	0.0329055	Glycosphingolipid biosynthesis -
dre00603	K03663	5	0	1	1	0.0468059	

							globo and isoglobo series
dre00190	K03967	87	0	3	1	0.0500817	Oxidative phosphorylation
dre01100	K00128	860	3	13	0.9721356	0.0626876	Metabolic pathways
dre00601	K07970	7	0	1	1	0.0649164	Glycosphingolipid biosynthesis - lacto and neolacto series
dre01212	K07514	44	0	2	1	0.0657289	Fatty acid metabolism
dre03320	K07514	54	1	2	0.3496255	0.0936223	PPAR signaling pathway
dre00531	K12309	11	0	1	1	0.1001236	Glycosaminoglycan degradation
dre04141	K10575	120	2	3	0.245355	0.1066185	Protein processing in endoplasmic reticulum
dre04540	K04157	59	0	2	1	0.1086435	Gap junction
dre04114	K04441	62	2	2	0.0861485	0.1179447	Oocyte meiosis
dre04080	K05226	67	4	2	0.0018878	0.1338667	Neuroactive ligand-receptor interaction
dre01040	K10203	15	0	1	1	0.1340209	Biosynthesis of unsaturated fatty acids
dre00650	K07514	15	0	1	1	0.1340209	Butanoate metabolism
dre00062	K01074	16	0	1	1	0.1422961	Fatty acid elongation
dre04145	K01365	73	0	2	1	0.1535658	Phagosome
dre04144	K05095	159	2	3	0.3593518	0.1936028	Endocytosis
dre00020	K00031	23	0	1	1	0.1980761	Citrate cycle (TCA cycle)
dre00982	K00699	24	0	1	1	0.2057464	Drug metabolism - cytochrome P450
dre01200	K07514	90	1	2	0.5125323	0.2118415	Carbon metabolism
dre00980	K00699	25	0	1	1	0.2133443	Metabolism of xenobiotics by cytochrome P450
dre04010	K05095	171	4	3	0.04608	0.2232692	MAPK signaling pathway
dre03040	K12862	95	0	2	1	0.2294162	Spliceosome
dre00520	K01183	28	0	1	1	0.2357097	Amino sugar and nucleotide sugar metabolism
dre00051	K17497	28	0	1	1	0.2357097	Fructose and mannose metabolism
dre00480	K00681	32	1	1	0.2247803	0.2645558	Glutathione metabolism

A. SUPPLEMENTARY MATERIAL

dre00260	K00315	36	0	1	1	0.2923261	Glycine, serine and threonine metabolism
dre00620	K00128	36	0	1	1	0.2923261	Pyruvate metabolism
dre00270	K08963	37	1	1	0.2550794	0.2991052	Cysteine and methionine metabolism
dre03050	K11598	37	0	1	1	0.2991052	Proteasome
dre00280	K00128	40	1	1	0.2726956	0.31906	Valine, leucine and isoleucine degradation
dre00983	K00699	43	0	1	1	0.3384534	Drug metabolism - other enzymes
dre04060	K04304	50	1	1	0.3285034	0.3816084	Cytokine-cytokine receptor interaction
dre04916	K05858	50	0	1	1	0.3816084	Melanogenesis
dre04146	K07514	53	0	1	1	0.3992409	Peroxisome
dre04914	K02649	53	0	1	1	0.3992409	Progesterone-mediated oocyte maturation
dre04260	K02265	54	1	1	0.3496255	0.4050073	Cardiac muscle contraction
dre05132	K10419	152	0	2	1	0.4273702	Salmonella infection
dre04012	K02649	58	0	1	1	0.4275311	ErbB signaling pathway

Supplementary Table 4. 7: List of KEGG pathways for up- and down-regulated pathways for the liver. ID refers to the universal identifiers of the KEGG terms, the N column represents the total number of genes annotated with each KEGG pathway. The Up and Down columns indicate the number of genes within the KEGG pathway that are significantly up- and down-regulated in high versus low treatment, respectively.

ID	ko	N	Up	Down	<i>p</i> - value Up	<i>p</i> -value Down	Pathway
dre03010	K02973	105	36	0	1.09E-20	1	Ribosome
dre03320	K07514	54	9	11	0.00174	1.461E-05	PPAR signaling pathway
dre00232	K00365	3	2	0	0.0078845	1	Caffeine metabolism
dre04210	K02649	83	10	2	0.010869	0.8784165	Apoptosis
dre00591	K07418	10	3	2	0.01	0.0661561	Linoleic acid metabolism
dre04260	K02265	54	6	4	0.06	0.2020317	Cardiac muscle contraction
dre04068	K08341	82	8	1	0.06	0.9733577	FoxO signaling pathway
dre00565	K16342	18	3	1	0.06	0.5474014	Ether lipid metabolism
dre00230	K00939	84	8	2	0.07	0.882582	Purine metabolism
dre00983	K00699	43	5	5	0.07	0.0361902	Drug metabolism - other enzymes
dre04115	K10140	33	4	0	0.09	1	p53 signaling pathway
dre00534	K01793	11	2	0	0.11	1	Glycosaminoglycan biosynthesis - heparan sulfate / heparin
dre00564	K00968	49	5	4	0.11	0.1587812	Glycerophospholipid metabolism
dre04150	K02649	83	7	2	0.14	0.8784165	mTOR signaling pathway
dre04810	K05095	131	10	3	0.15	0.9257988	Regulation of actin cytoskeleton
dre04216	K14714	26	3	5	0.15	0.0044765	Ferroptosis
dre00830	K00699	26	3	1	0.15	0.6819761	Retinol metabolism
dre00561	K00128	40	4	1	0.15	0.8286383	Glycerolipid metabolism
dre04620	K02649	41	4	0	0.16	1	Toll-like receptor signaling pathway
dre03008	K14525	56	5	1	0.17	0.9155784	Ribosome biogenesis in eukaryotes
dre03050	K11598	37	1	24	0.86	1.68E-24	Proteasome
dre01212	K07514	44	3	12	0.41	2.05E-07	Fatty acid metabolism
dre04141	K10575	120	3	19	0.95	7.513E-07	Protein processing in endoplasmic reticulum
dre01100	K00128	860	48	63	0.33	1.397E-05	Metabolic pathways

A. SUPPLEMENTARY MATERIAL

dre03320	K07514	54	9	11	0.00	1.461E-05	PPAR signaling pathway
dre01040	K10203	15	0	6	1.00	2.196E-05	Biosynthesis of unsaturated fatty acids
dre00061	K15013	13	0	5	1.00	0.000139	Fatty acid biosynthesis
dre00513	K00736	27	0	6	1.00	0.0008373	Various types of N-glycan biosynthesis
dre04145	K01365	73	3	10	0.74	0.0010644	Phagosome
dre00510	K00736	29	0	6	1.00	0.0012493	N-Glycan biosynthesis
dre04142	K06501	90	3	11	0.86	0.001586	Lysosome
dre00480	K00681	32	2	6	0.50	0.0021372	Glutathione metabolism
dre00982	K00699	24	2	5	0.36	0.0031042	Drug metabolism - cytochrome P450
dre04216	K14714	26	3	5	0.15	0.0044765	Ferroptosis
dre00592	K16342	9	1	3	0.38	0.0054732	alpha-Linolenic acid metabolism
dre01200	K07514	90	4	9	0.70	0.0150273	Carbon metabolism
dre00071	K00128	35	3	5	0.28	0.0161108	Fatty acid degradation
dre02010	K05665	24	0	4	1.00	0.0181249	ABC transporters
dre00980	K00699	25	1	4	0.74	0.0208638	Metabolism of xenobiotics by cytochrome P450
dre04146	K07514	53	4	6	0.30	0.0254687	Peroxisome
dre00062	K01074	16	1	3	0.58	0.0291849	Fatty acid elongation
dre00190	K03967	87	5	8	0.48	0.0331838	Oxidative phosphorylation
dre00983	K00699	43	5	5	0.07	0.0361902	Drug metabolism - other enzymes
dre00590	K07418	20	2	3	0.28	0.0523814	Arachidonic acid metabolism
dre00140	K00699	20	1	3	0.66	0.0523814	Steroid hormone biosynthesis
dre00350	K00431	20	1	3	0.66	0.0523814	Tyrosine metabolism
dre00591	K07418	10	3	2	0.01	0.0661561	Linoleic acid metabolism
dre00500	K01084	22	0	3	1.00	0.0664734	Starch and sucrose metabolism
dre00620	K00128	36	1	4	0.86	0.067376	Pyruvate metabolism
dre00410	K00128	23	1	3	0.71	0.0741012	beta-Alanine metabolism
dre00670	K13403	13	0	2	1.00	0.1054345	One carbon pool by folate
dre00640	K07514	27	0	3	1.00	0.1081439	Propanoate metabolism
dre00330	K00128	28	2	3	0.43	0.1174564	Arginine and proline metabolism

dre00600	K12357	28	0	3	1.00	0.1174564	Sphingolipid metabolism
dre00524	K00844	3	0	1	1.00	0.1236652	Neomycin, kanamycin and gentamicin biosynthesis
dre00650	K07514	15	2	2	0.18	0.1342785	Butanoate metabolism
dre00630	K01915	30	1	3	0.80	0.1369243	Glyoxylate and dicarboxylate metabolism
dre04144	K05095	159	4	10	0.97	0.1474093	Endocytosis
dre00340	K00128	16	2	2	0.20	0.1492919	Histidine metabolism
dre00564	K00968	49	5	4	0.11	0.1587812	Glycerophospholipid metabolism

B Appendix 1: Iteration process for QTL mapping

Construction of the genetic map (First round):

Using the pedigree information obtained with the variant call from Chapter 2, we selected the largest family with both parents genotyped (n=89) and used LepMap3. We filtered the loci of using the Filtering2 module (Number of called markers = 171923 (59019 informative) based on high segregation distortion (dataTolerance= 0.001 respectively). We then used the SeparateChromosomes2 module to clustered the loci into linkage groups (LGs) based on recombination rates by calculating logarithm of the odds (LOD) scores between all pairs of loci with the. The LOD scores were chosen by increasing the LOD value by one until the number of LGs stabilized. The final LOD scores used to generate the map was 12. We then added remaining markers using the JoinSingles2All module with a LOD threshold of 5. Loci were then ordered within LGs by utilizing the OrderMarkers2 module and sex-averaged map distances were computed with the options sexAveraged=1 and hypePhase=1. Finally we manually orientated the scaffolds using MapChart, and manually removed markers co-mapping at the same region.

QTL mapping (first round):

We converted the merged call to a plink .raw format, filtered it to keep only the individuals from the family and the markers in the map and converted the genotypes to AC, AD, BC, BD. We then loaded the file in R/qtl as a 4-way cross. This is where the problems started. Using this file, the threshold computed after permutations were high (>5) which meant that none of the peaks were significant. We checked for distortion in the markers maps and we saw that quite a few markers had a chi2 inflated, especially at the tails of the LGs and despite having filtered the markers in LepMap Filtering2 step.

We then decided to check the distortion of all the markers found after merging the parents and offspring calls. This resulted in a set of 33k markers.

Construction of the genetic map (second round):

We used the 33k markers as an input for LepMap and kept the same parameters than for the previous round. The Marey plots all looked fine.

QTL mapping (second round):

The same inflation happened in R/qtl. We use MapQTL at that point to check the plots, since R/qtl only allows to see the genome-wide permutations, and we wanted to check if the high significance threshold was due to only one LG. So we loaded the map, genotypes and phenotypes in MapQTL and run Interval Mapping, before running the permutation test. When we looked at the charts (LOD score over genetic distance for each cM) we saw that the LOD curve was not continuous, and it had multiple drops to 0 across all LG and no clear QTL peak. We then realized there was still something wrong with the map where some markers are still mis-positioned.

Construction of the genetic map (third round):

We input the 33k markers into JoinMap and manually check the markers that were segregating ABxAA, ABxBB, ABxAB, AAxAB and BBxAB. We used set the LOD parameters to go up to 20 and generated the 24 LGs as expected. But when looking closer, the LGs were all VERY long (~3,000cM and filtering for NNstress was only reducing the distance a little bit).

Construction of the genetic map (fourth round):

We thought that might be due to the fact markers from both parents were input in the same file so we generated a file for the mother containing the markers segregating ABxAA, ABxBB and ABxAB and another file with the father's markers segregating AAxAB, BBxAB and ABxAB. We got the same results of inflated LG lengths. There is a very large genetic distance between each markers, so the huge length of the LG is not due to gaps. There is one encouraging fact though: the grouping seems all right because we find correspondence between LG and super-scaffold.

Construction of the genetic map (fourth round):

We tried a regression algorithm, versus maximum likelihood that we were using before and which is the default in JoinMap that is trying markers pair by pair. But with the number of markers per LG that we have the computation is taking forever (one LG with ~350 markers has been running since Monday morning

Using freebayes call (fifth round):

Filtered for 2% missing data, SNPs with a mendelian error rate $> 2\%$ which yielded 48k SNPs, and a further filter for segregation distortion and got back a set of 13.3k SNPs. Same issue, high thresholds from R/qtl and long LG with spaced markers in JoinMap. A couple of Marey plots from the LepMap map are included. We started thinking there might be a few mistakes in the genome scaffolds as the y axis (cM) looks fine but there are big gaps in the middle on the x axis (bp). At this stage the genome assembly was not polished, so misplacement of markers in the marey plots might be due more to that than to the genetic map.

Marey plot of LG1 obtained using the freebayes call. The x-axis represents the physical position (bp) and the y-axis the genetic distance (cM).

Still issues, for instance, the first LG obtained with JoinMap was generated from the stringent filtered data and after 4 extra rounds of filtering for nearest neighbour stress. Only 79 markers were left yet the LG was 147cM. We then investigated again different ways to input the data into R/qtl. We were using a 4 way cross, but thought there might be another more appropriate option to this data set. The Marey plots were showing a reduced recombination rate around the centromere, which give that sort of “plateau” in the middle of the plot. The lack of recombination and hence very extended haplotypes through that region mean that once a SNP is in the linkage map, it captures all of the information through that region.

The crux!

Found in this tutorial: <https://avikarn.com/2019-04-17-Genetic-Mapping-in-Lep-MAP3/#-covnvert-genotype-data-into-4-way-cross-rqtl-input-data->

1	1	1	AC
1	2	2	BC
2	1	3	AD
2	2	4	BD

I realised that I was misunderstanding the way LepMap was outputting the genotypes and I was converting 1 2 into AD instead of BC and 2 1 into BC instead of AD...

Bibliography

- Abbink, W., Blanco Garcia, A., Roques, J. A. C., Partridge, G. J., Kloet, K., & Schneider, O. (2012). The effect of temperature and pH on the growth and physiological response of juvenile yellowtail kingfish *Seriola lalandi* in recirculating aquaculture systems. *Aquaculture*, 330–333, 130–135. <https://doi.org/10.1016/j.aquaculture.2011.11.043>
- Abram, P. K., Boivin, G., Moiroux, J., & Brodeur, J. (2017). Behavioural effects of temperature on ectothermic animals: Unifying thermal physiology and behavioural plasticity. *Biological Reviews*, 92(4), 1859–1876. <https://doi.org/10.1111/brv.12312>
- Agrawal, A. A. (1999). Induced plant defense: Evolution of induction and adaptive phenotypic plasticity. *Inducible Plant Defenses against Pathogens and Herbivores: Biochemistry, Ecology, and Agriculture*. American Phytopathological Society Press, St. Paul, MN, 251–268.
- Agrawal, A. A. (2001). Phenotypic Plasticity in the Interactions and Evolution of Species. *Science*, 294(5541), 321–326. <https://doi.org/10.1126/science.1060701>
- Albertin, W., Marullo, P., Bely, M., Aigle, M., Bourgaïs, A., Langella, O., Balliau, T., Chevret, D., Valot, B., da Silva, T., Dillmann, C., de Vienne, D., & Sicard, D. (2013). Linking Post-Translational Modifications and Variation of Phenotypic Traits*. *Molecular & Cellular Proteomics*, 12(3), 720–735. <https://doi.org/10.1074/mcp.M112.024349>
- Allison, D. B., Fernandez, J. R., Heo, M., Zhu, S., Etzel, C., Beasley, T. M., & Amos, C. I. (2002). Bias in estimates of quantitative-trait–locus effect in genome scans: Demonstration of the phenomenon and a method-of-moments procedure for reducing bias. *The American Journal of Human Genetics*, 70(3), 575–585.
- Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Research*, 25(17), 3389–3402.
- Anders, S., Pyl, P. T., & Huber, W. (2015). HTSeq—A Python framework to work with high-throughput sequencing data. *Bioinformatics*, 31(2), 166–169.
- Andrews, S. (2010). Babraham bioinformatics-FastQC a quality control tool for high throughput sequence data. *Babraham Inst.*
- Andrews, S. (2017). *FastQC: A quality control tool for high throughput sequence data*. 2010.

-
- Aronesty, E. (2013). Comparison of Sequencing Utility Programs. *The Open Bioinformatics Journal*, 7(1), 1–8. <https://doi.org/10.2174/1875036201307010001>
- Ashton, D. T., Hilario, E., Jaksons, P., Ritchie, P. A., & Wellenreuther, M. (2019). Genetic diversity and heritability of economically important traits in captive Australasian snapper (*Chrysophrys auratus*). *Aquaculture*, 505, 190–198. <https://doi.org/10.1016/j.aquaculture.2019.02.034>
- Ashton, D. T., Ritchie, P. A., & Wellenreuther, M. (2017). Fifteen years of quantitative trait loci studies in fish: Challenges and future directions. *Molecular Ecology*, 26(6), 1465–1476. <https://doi.org/10.1111/mec.13965>
- Ashton, D. T., Ritchie, P. A., & Wellenreuther, M. (2019). High-Density Linkage Map and QTLs for Growth in Snapper (*Chrysophrys auratus*). *G3 Genes/Genomes/Genetics*, 9(4), 1027–1035. <https://doi.org/10.1534/g3.118.200905>
- Aubin-Horth, N., & Renn, S. C. P. (2009). Genomic reaction norms: Using integrative biology to understand molecular mechanisms of phenotypic plasticity. *Molecular Ecology*, 18(18), 3763–3780. <https://doi.org/10.1111/j.1365-294X.2009.04313.x>
- Auer, P. L., & Doerge, R. W. (2011). A two-stage Poisson model for testing RNA-seq data. *Statistical Applications in Genetics and Molecular Biology*, 10(1).
- Ayroles, J. F., Carbone, M. A., Stone, E. A., Jordan, K. W., Lyman, R. F., Magwire, M. M., Rollmann, S. M., Duncan, L. H., Lawrence, F., & Anholt, R. R. (2009). Systems genetics of complex traits in *Drosophila melanogaster*. *Nature Genetics*, 41(3), 299–307.
- Azzurro, E., Sbragaglia, V., Cerri, J., Bariche, M., Bolognini, L., Ben Souissi, J., Busoni, G., Coco, S., Chryssanthi, A., & Fanelli, E. (2019). Climate change, biological invasions, and the shifting distribution of Mediterranean fishes: A large-scale survey based on local ecological knowledge. *Global Change Biology*, 25(8), 2779–2792.
- Baranski, M., Moen, T., & Våge, D. I. (2010). Mapping of quantitative trait loci for flesh colour and growth traits in Atlantic salmon (*Salmo salar*). *Genetics Selection Evolution*, 42(1), 17. <https://doi.org/10.1186/1297-9686-42-17>
- Barton, N. H., & Turelli, M. (1989). Evolutionary quantitative genetics: How little do we know. *Annual Review of Genetics*, 23(1), 337–370.
- Bazakos, C., Hanemian, M., Trontin, C., Jiménez-Gómez, J. M., & Loudet, O. (2017). New strategies and tools in quantitative genetics: How to go from the phenotype to the genotype. *Annual Review of Plant Biology*, 68, 435–455.

- Beaman, J. E., White, C. R., & Seebacher, F. (2016). Evolution of Plasticity: Mechanistic Link between Development and Reversible Acclimation. *Trends in Ecology & Evolution*, 31(3), 237–249. <https://doi.org/10.1016/j.tree.2016.01.004>
- Bearham, D., Robert, M., Chaplin, J. A., Moore, G. I., Fairclough, D. V., Bertram, A., Bearham, D., Robert, M., Chaplin, J. A., Moore, G. I., Fairclough, D. V., & Bertram, A. (2020). Molecular evidence of three species in the *Pseudocaranx dentex* complex (Carangidae) in Australian waters. *Marine and Freshwater Research*, 71(4), 518–531. <https://doi.org/10.1071/MF18445>
- Beavis, W. D. (1994). The power and deceit of QTL experiments: Lessons from comparative QTL studies. *Proceedings of the Forty-Ninth Annual Corn and Sorghum Industry Research Conference*, 250, 266.
- Beissinger, T. M., Hirsch, C. N., Sekhon, R. S., Foerster, J. M., Johnson, J. M., Muttoni, G., Vaillancourt, B., Buell, C. R., Kaeppler, S. M., & de Leon, N. (2013). Marker density and read depth for genotyping populations using genotyping-by-sequencing. *Genetics*, 193(4), 1073–1081.
- Béné, C., Arthur, R., Norbury, H., Allison, E. H., Beveridge, M., Bush, S., Campling, L., Leschen, W., Little, D., & Squires, D. (2016). Contribution of fisheries and aquaculture to food security and poverty reduction: Assessing the current evidence. *World Development*, 79, 177–196.
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(1), 289–300.
- Bentsbn, H. B. (1994). Genetic effects of selection on polygenic traits with examples from Atlantic salmon, *Salmo salar* L. *Aquaculture Research*, 25(1), 89–102.
- Besnier, F., Solberg, M. F., Harvey, A. C., Carvalho, G. R., Bekkevold, D., Taylor, M. I., Creer, S., Nielsen, E. E., Skaala, Ø., Ayllon, F., Dahle, G., & Glover, K. A. (2020). Epistatic regulation of growth in Atlantic salmon revealed: A QTL study performed on the domesticated-wild interface. *BMC Genetics*, 21(1), 13. <https://doi.org/10.1186/s12863-020-0816-y>
- Blanco, A. M. (2020). Hypothalamic- and pituitary-derived growth and reproductive hormones and the control of energy balance in fish. *General and Comparative Endocrinology*, 287, 113322. <https://doi.org/10.1016/j.ygcen.2019.113322>

-
- Bolger, A., Lohse, M., & Usadel, B. (2014). *Trimmomatic: A flexible trimmer for Illumina sequence data* / *Bioinformatics* / Oxford Academic. <https://academic.oup.com/bioinformatics/article/30/15/2114/2390096>
- Borges, R. M. (2009). Phenotypic plasticity and longevity in plants and animals: Cause and effect? *Journal of Biosciences*, 34(4), 605–611.
- Bowler, K., & Terblanche, J. S. (2008). Insect thermal tolerance: What is the role of ontogeny, ageing and senescence? *Biological Reviews*, 83(3), 339–355. <https://doi.org/10.1111/j.1469-185X.2008.00046.x>
- Braasch, I., Peterson, S. M., Desvignes, T., McCluskey, B. M., Batzel, P., & Postlethwait, J. H. (2015). A new model army: Emerging fish models to study the genomics of vertebrate Evo-Devo. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, 324(4), 316–341.
- Brett, J. R., & Groves, T. D. D. (1979). Physiological energetics. *Fish Physiology*, 8(6).
- Brierley, A. S., & Kingsford, M. J. (2009). Impacts of climate change on marine organisms and ecosystems. *Current Biology*, 19(14), R602–R614.
- Broman, K. W., Wu, H., Sen, S., & Churchill, G. A. (2003). R/qtl: QTL mapping in experimental crosses. *Bioinformatics*, 19(7), 889–890. <https://doi.org/10.1093/bioinformatics/btg112>
- Brooks, A. (2020). *Using Ecological Niche Modelling to Predict Climate Change Responses of Ten Key Fishery Species in Aotearoa New Zealand*.
- Brown, E. J., Bruce, M., Pether, S., & Herbert, N. A. (2011). Do swimming fish always grow fast? Investigating the magnitude and physiological basis of exercise-induced growth in juvenile New Zealand yellowtail kingfish, *Seriola lalandi*. *Fish Physiology and Biochemistry*, 37(2), 327–336. <https://doi.org/10.1007/s10695-011-9500-5>
- Brun, P., Thuiller, W., Chauvier, Y., Pellissier, L., Wüest, R. O., Wang, Z., & Zimmermann, N. E. (2020). Model complexity affects species distribution projections under climate change. *Journal of Biogeography*, 47(1), 130–142.
- Bullard, J. H., Purdom, E., Hansen, K. D., & Dudoit, S. (2010). Evaluation of statistical methods for normalization and differential expression in mRNA-Seq experiments. *BMC Bioinformatics*, 11(1), 1–13.
- Byun, J., Logothetis, C. J., & Gorlov, I. P. (2009). Housekeeping genes in prostate tumorigenesis. *International Journal of Cancer*, 125(11), 2603–2608.

- Camara, M. D., & Symonds, J. E. (2014). Genetic improvement of New Zealand aquaculture species: Programmes, progress and prospects. *New Zealand Journal of Marine and Freshwater Research*, 48(3), 466–491. <https://doi.org/10.1080/00288330.2014.932291>
- Camara, M., & Symonds, J. (2014). Genetic improvement of New Zealand aquaculture species: Programmes, progress and prospects. *New Zealand Journal of Marine and Freshwater Research*, 48(3), 466–491. <https://doi.org/10.1080/00288330.2014.932291>
- Campitelli, B. E., Marais, D. L. D., & Juenger, T. E. (2016). Ecological interactions and the fitness effect of water-use efficiency: Competition and drought alter the impact of natural MPK12 alleles in Arabidopsis. *Ecology Letters*, 19(4), 424–434. <https://doi.org/10.1111/ele.12575>
- Canosa, L. F., Chang, J. P., & Peter, R. E. (2007). Neuroendocrine control of growth hormone in fish. *General and Comparative Endocrinology*, 151(1), 1–26.
- Carlborg, Ö., & Haley, C. S. (2004). Epistasis: Too often neglected in complex trait studies? *Nature Reviews Genetics*, 5(8), 618–625. <https://doi.org/10.1038/nrg1407>
- Carlson, M. (2019). *org. Mm. eg. db: Genome wide annotation for Mouse. R package version 3.2.3*. Bioconductor.
- Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: An analysis tool set for population genomics. *Molecular Ecology*, 22(11), 3124–3140. <https://doi.org/10.1111/mec.12354>
- Chapman, J. A., Ho, I., Sunkara, S., Luo, S., Schroth, G. P., & Rokhsar, D. S. (2011). Meraculous: De novo genome assembly with short paired-end reads. *PloS One*, 6(8), e23501.
- Chattopadhyay, & Chattoraj. (2017). A Review on the Reproduction and Development in Fish. *Biomedical Journal of Scientific & Technical Research*, 1(6). <https://doi.org/10.26717/BJSTR.2017.01.000529>
- Chavanne, H. (2016). A comprehensive survey on selective breeding programs and seed market in the European aquaculture fish industry. *Aquacult Int*, 21.
- Claussnitzer, M., Cho, J. H., Collins, R., Cox, N. J., Dermitzakis, E. T., Hurles, M. E., Kathiresan, S., Kenny, E. E., Lindgren, C. M., & MacArthur, D. G. (2020). A brief history of human disease genetics. *Nature*, 577(7789), 179–189.
- Cloonan, N., Forrest, A. R., Kolle, G., Gardiner, B. B., Faulkner, G. J., Brown, M. K., Taylor, D. F., Steptoe, A. L., Wani, S., & Bethel, G. (2008). Stem cell transcriptome profiling via massive-scale mRNA sequencing. *Nature Methods*, 5(7), 613–619.

-
- Concibido, V. C., Diers, B. W., & Arelli, P. R. (2004). A Decade of QTL Mapping for Cyst Nematode Resistance in Soybean. *Crop Science*, 44(4), 1121–1131. <https://doi.org/10.2135/cropsci2004.1121>
- Conesa, A., Madrigal, P., Tarazona, S., Gomez-Cabrero, D., Cervera, A., McPherson, A., Szcześniak, M. W., Gaffney, D. J., Elo, L. L., Zhang, X., & Mortazavi, A. (2016). A survey of best practices for RNA-seq data analysis. *Genome Biology*, 17(1), 13. <https://doi.org/10.1186/s13059-016-0881-8>
- Cooper, M., Tang, T., Ghosh, C., Hart, T., Hammer, G., & Messina, C. (2020). Integrating genetic gain and gap analysis to predict improvements in crop productivity. *Crop Science*, 60(2), 582–604.
- Coppieters, W., Riquet, J., Arranz, J.-J., Berzi, P., Cambisano, N., Grisart, B., Karim, L., Marcq, F., Moreau, L., & Nezer, C. (1998). A QTL with major effect on milk yield and composition maps to bovine chromosome 14. *Mammalian Genome*, 9(7), 540–544.
- Cowan, M., Azpeleta, C., & López-Olmeda, J. F. (2017). Rhythms in the endocrine system of fish: A review. *Journal of Comparative Physiology B*, 187(8), 1057–1089. <https://doi.org/10.1007/s00360-017-1094-5>
- Crabbe, J. C., Phillips, T. J., Kosobud, A., & Belknap, J. K. (1990). Estimation of genetic correlation: Interpretation of experiments using selectively bred and inbred animals. *Alcoholism: Clinical and Experimental Research*, 14(2), 141–151.
- Crossa, J., Campos, G. de los, Pérez, P., Gianola, D., Burgueño, J., Araus, J. L., Makumbi, D., Singh, R. P., Dreisigacker, S., & Yan, J. (2010). Prediction of genetic values of quantitative traits in plant breeding using pedigree and molecular markers. *Genetics*, 186(2), 713–724.
- Crosses, E. (2001). Review of statistical methods for QTL mapping in experimental crosses. *Lab Animal*, 30(7), 44–52.
- Crouch, D. J. M., & Bodmer, W. F. (2020). Polygenic inheritance, GWAS, polygenic risk scores, and the search for functional variants. *Proceedings of the National Academy of Sciences*, 117(32), 18924–18933. <https://doi.org/10.1073/pnas.2005634117>
- Cryer, P., & Cherring, A. D. (2001). *Handbook of Physiology: Section 7, The Endocrine System, Volume II, The Endocrine Pancreas and Regulation of Metabolism*.
- Dahlke, F. T., Wohlrab, S., Butzin, M., & Pörtner, H.-O. (2020). Thermal bottlenecks in the life cycle define climate vulnerability of fish. *Science*, 369(6499), 65–70.
- Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes

- Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, 27(15), 2156–2158. <https://doi.org/10.1093/bioinformatics/btr330>
- Darwin, C. (1858). *The Origin of Species by Means of Natural Selection or the Preservation of Favoured Races in the Struggle for Life*. Murray, London. r.
- Darwin, C., Wallace, A. R., Lyell, S. C., & Hooker, J. D. (1858). *On the tendency of species to form varieties: And on the perpetuation of varieties and species by natural means of selection*.
- Davies, I. P., Carranza, V., Froehlich, H. E., Gentry, R. R., Kareiva, P., & Halpern, B. S. (2019). Governance of marine aquaculture: Pitfalls, potential, and pathways forward. *Marine Policy*, 104, 29–36. <https://doi.org/10.1016/j.marpol.2019.02.054>
- Deane, E. E., & Woo, N. Y. S. (2009). Modulation of fish growth hormone levels by salinity, temperature, pollutants and aquaculture related stress: A review. *Reviews in Fish Biology and Fisheries*, 19(1), 97–120. <https://doi.org/10.1007/s11160-008-9091-0>
- del Pazo, F., Sánchez, S., Posner, V., Sciara, A. A., Arranz, S. E., & Villanova, G. V. (2021). Genetic diversity and structure of the commercially important native fish pacu (*Piaractus mesopotamicus*) from cultured and wild fish populations: Relevance for broodstock management. *Aquaculture International*, 29(1), 289–305. <https://doi.org/10.1007/s10499-020-00626-w>
- Deniel, C. (1990). Comparative study of growth of flatfishes on the west coast of Brittany. *Journal of Fish Biology*, 37(1), 149–166.
- De-Santis, C., & Jerry, D. R. (2007). Candidate growth genes in finfish—Where should we be looking? *Aquaculture*, 272(1), 22–38. <https://doi.org/10.1016/j.aquaculture.2007.08.036>
- Dettliff, P., Hernandez, E., Partridge, G., Lafarga-De la Cruz, F., & Martinez, V. (2020). Understanding the population structure and reproductive behavior of hatchery-produced yellowtail kingfish (*Seriola lalandi*). *Aquaculture*, 522, 734948. <https://doi.org/10.1016/j.aquaculture.2020.734948>
- Devlin, R. H., & Nagahama, Y. (2002). Sex determination and sex differentiation in fish: An overview of genetic, physiological, and environmental influences. *Aquaculture*, 208(3), 191–364. [https://doi.org/10.1016/S0044-8486\(02\)00057-1](https://doi.org/10.1016/S0044-8486(02)00057-1)
- DeWitt, T. J., Sih, A., & Wilson, D. S. (1998). Costs and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, 13(2), 77–81. [https://doi.org/10.1016/S0169-5347\(97\)01274-3](https://doi.org/10.1016/S0169-5347(97)01274-3)

-
- Dewitt, T., & Scheiner, S. (2004). Phenotypic plasticity: Functional and conceptual approaches. In *American Journal of Human Biology—AMER J HUM BIOL* (Vol. 17).
- Dmitriew, C. M. (2011). The evolution of growth trajectories: What limits growth rate? *Biological Reviews*, 86(1), 97–116. <https://doi.org/10.1111/j.1469-185X.2010.00136.x>
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, 29(1), 15–21.
- Doerge, R. W., & Churchill, G. A. (1996). Permutation Tests for Multiple Loci Affecting a Quantitative Character. *Genetics*, 142(1), 285–294. <https://doi.org/10.1093/genetics/142.1.285>
- Duan, C., Ren, H., & Gao, S. (2010). Insulin-like growth factors (IGFs), IGF receptors, and IGF-binding proteins: Roles in skeletal muscle growth and differentiation. *General and Comparative Endocrinology*, 167(3), 344–351. <https://doi.org/10.1016/j.ygcen.2010.04.009>
- Dwivedi, S. L., Stoddard, F. L., & Ortiz, R. (2019). Genomic-based root plasticity to enhance abiotic stress adaptation and edible yield in grain crops. *Plant Science*, 110365.
- Ellegren, H., & Sheldon, B. C. (2008). Genetic basis of fitness differences in natural populations. *Nature*, 452(7184), 169–175.
- Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler, E. S., & Mitchell, S. E. (2011). A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. *PLOS ONE*, 6(5), e19379. <https://doi.org/10.1371/journal.pone.0019379>
- Enberg, K., Jørgensen, C., Dunlop, E. S., Varpe, Ø., Boukal, D. S., Baulier, L., Eliassen, S., & Heino, M. (2012). Fishing-induced evolution of growth: Concepts, mechanisms and the empirical evidence. *Marine Ecology*, 33(1), 1–25. <https://doi.org/10.1111/j.1439-0485.2011.00460.x>
- Ern, R., Norin, T., Gamperl, A. K., & Esbaugh, A. J. (2016). Oxygen dependence of upper thermal limits in fishes. *Journal of Experimental Biology*, 219(21), 3376–3383. <https://doi.org/10.1242/jeb.143495>
- Essington, T. E., Kitchell, J. F., & Walters, C. J. (2011). The von Bertalanffy growth function, bioenergetics, and the consumption rates of fish. *Canadian Journal of Fisheries and Aquatic Sciences*. <https://doi.org/10.1139/f01-151>

- Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for multiple tools and samples in a single report. *Bioinformatics*, 32(19), 3047–3048.
- Falconer, D., & Mackay, T. (1996). *Introduction to quantitative genetics*. https://scholar.google.com/scholar_lookup?title=Introduction%20to%20Quantitative%20Genetics&publication_year=1996&author=D.%20Falconer&author=T.F.C.%20Mackay
- Falconer, D. S. (1960). Selection of mice for growth on high and low planes of nutrition. *Genetics Research*, 1(1), 91–113.
- Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics*. Essex. UK: Longman Group.
- FAO (Ed.). (2018). *Meeting the sustainable development goals*.
- Farrall, M. (2004). Quantitative genetic variation: A post-modern view. *Human Molecular Genetics*, 13(suppl_1), R1–R7. <https://doi.org/10.1093/hmg/ddh084>
- Feinberg, A. P. (2007). Phenotypic plasticity and the epigenetics of human disease. *Nature*, 447(7143), 433–440.
- Fernández-Montero, A., Caballero, M. J., Torrecillas, S., Tuset, V. M., Lombarte, A., Ginés, R. R., Izquierdo, M., Robaina, L., & Montero, D. (2018). Effect of temperature on growth performance of greater amberjack (*Seriola dumerili* Risso 1810) Juveniles. *Aquaculture Research*, 49(2), 908–918. <https://doi.org/10.1111/are.13537>
- Fernández-Montero, A., Torrecillas, S., Tort, L., Ginés, R., Acosta, F., Izquierdo, M. S., & Montero, D. (2020). Stress response and skin mucus production of greater amberjack (*Seriola dumerili*) under different rearing conditions. *Aquaculture*, 520, 735005. <https://doi.org/10.1016/j.aquaculture.2020.735005>
- Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate change on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, 17(4), 581–613. <https://doi.org/10.1007/s11160-007-9059-5>
- Fisher, Ronald A. (1918). XV.—The correlation between relatives on the supposition of Mendelian inheritance. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh*, 52(2), 399–433.
- Fisher, Ronald Aylmer. (1930). *The genetical theory of natural selection*. Oxford at the Clarendon Press.
- Fisheries New Zealand. (2019). *Fisheries Assessment Plenary, May 2019: Stock assessments and stock status*. Fisheries New Zealand Wellington, New Zealand.

-
- Frankham, R., Bradshaw, C. J., & Brook, B. W. (2014). Genetics in conservation management: Revised recommendations for the 50/500 rules, Red List criteria and population viability analyses. *Biological Conservation*, 170, 56–63.
- Fraslin, C., Quillet, E., Rochat, T., Dechamp, N., Bernardet, J.-F., Collet, B., Lallias, D., & Boudinot, P. (2020). Combining Multiple Approaches and Models to Dissect the Genetic Architecture of Resistance to Infections in Fish. *Frontiers in Genetics*, 11. <https://doi.org/10.3389/fgene.2020.00677>
- Froese, R., Thorson, J. T., & Reyes, R. B. (2014). A Bayesian approach for estimating length-weight relationships in fishes. *Journal of Applied Ichthyology*, 30(1), 78–85. <https://doi.org/10.1111/jai.12299>
- Fry, F. E. (1971). The effect of environmental factors on the physiology of fish. *Fish Physiology*, 1–98.
- Fu, B., Wang, X., Feng, X., Yu, X., & Tong, J. (2016). Comparative transcriptomic analyses of two bighead carp (*Hypophthalmichthys nobilis*) groups with different growth rates. *Comparative Biochemistry and Physiology Part D: Genomics and Proteomics*, 20, 111–117. <https://doi.org/10.1016/j.cbd.2016.08.006>
- Fu, B., Yu, X., Tong, J., Pang, M., Zhou, Y., Liu, Q., & Tao, W. (2019). Comparative transcriptomic analysis of hypothalamus-pituitary-liver axis in bighead carp (*Hypophthalmichthys nobilis*) with differential growth rate. *BMC Genomics*, 20(1), 328. <https://doi.org/10.1186/s12864-019-5691-4>
- Fuentes, E. N., Valdés, J. A., Molina, A., & Björnsson, B. T. (2013). Regulation of skeletal muscle growth in fish by the growth hormone–insulin-like growth factor system. *General and Comparative Endocrinology*, 192, 136–148.
- Fukusho, K. (1995). *Recent developments in aquaculture in Japan*. 8.
- GAA. (2021). *Global Aquaculture Alliance*. Global Aquaculture Alliance. <https://www.aquaculturealliance.org/>
- Galappaththi, E. K., Ichien, S. T., Hyman, A. A., Aubrac, C. J., & Ford, J. D. (2020). Climate change adaptation in aquaculture. *Reviews in Aquaculture*, 12(4), 2160–2176. <https://doi.org/10.1111/raq.12427>
- Galwey, N. W. (2014). *Introduction to mixed modelling: Beyond regression and analysis of variance*. John Wiley & Sons.
- Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read sequencing. *ArXiv:1207.3907 [q-Bio]*. <http://arxiv.org/abs/1207.3907>

- Gebhardt-Henrich, S., & Richner, H. (1998). Causes of growth variation and its consequences for fitness. *Oxford Ornithology Series*, 8, 324–339.
- Gentry, R. R., Froehlich, H. E., Grimm, D., Kareiva, P., Parke, M., Rust, M., Gaines, S. D., & Halpern, B. S. (2017). Mapping the global potential for marine aquaculture. *Nature Ecology & Evolution*, 1(9), 1317–1324. <https://doi.org/10.1038/s41559-017-0257-9>
- Georges, M., Nielsen, D., Mackinnon, M., Mishra, A., Okimoto, R., Pasquino, A. T., Sargeant, L. S., Sorensen, A., Steele, M. R., & Zhao, X. (1995). Mapping quantitative trait loci controlling milk production in dairy cattle by exploiting progeny testing. *Genetics*, 139(2), 907–920.
- Gilmour, A. R., Gogel, B. J., Cullis, B. R., Welham, S. J., & Thompson, R. (2015). ASReml user guide release 4.1 structural specification. *Hemel Hempstead: VSN International Ltd.*
- Gjedrem, T. (Ed.). (2005). *Selection and breeding programs in aquaculture*. Springer.
- Gjedrem, T., & Baranski, M. (2009). The success of selective breeding in aquaculture. In *Selective Breeding in Aquaculture: An Introduction* (pp. 13–23). Springer.
- Gjedrem, T., Robinson, N., & Rye, M. (2012). The importance of selective breeding in aquaculture to meet future demands for animal protein: A review. *Aquaculture*, 350–353, 117–129. <https://doi.org/10.1016/j.aquaculture.2012.04.008>
- Gjerde, B., Simianer, H., & Refstie, T. (1994). Estimates of genetic and phenotypic parameters for body weight, growth rate and sexual maturity in Atlantic salmon. *Livestock Production Science*, 38(2), 133–143.
- Gjuvslund, A. B., Hayes, B. J., Omholt, S. W., & Carlborg, O. (2007). Statistical epistasis is a generic feature of gene regulatory networks. *Genetics*, 175(1), 411–420.
- Glover, K. A., Solberg, M. F., McGinnity, P., Hindar, K., Verspoor, E., Coulson, M. W., Hansen, M. M., Araki, H., Skaala, Ø., & Svåsand, T. (2017). Half a century of genetic interaction between farmed and wild Atlantic salmon: Status of knowledge and unanswered questions. *Fish and Fisheries*, 18(5), 890–927. <https://doi.org/10.1111/faf.12214>
- Gobler, C. J., Merlo, L. R., Morrell, B. K., & Griffith, A. W. (2018). Temperature, Acidification, and Food Supply Interact to Negatively Affect the Growth and Survival of the Forage Fish, *Menidia beryllina* (Inland Silverside), and *Cyprinodon variegatus* (Sheepshead Minnow). *Frontiers in Marine Science*, 5. <https://doi.org/10.3389/fmars.2018.00086>
- Goddard, M. E., & Hayes, B. J. (2009). Mapping genes for complex traits in domestic animals and their use in breeding programmes. *Nature Reviews Genetics*, 10(6), 381–391.

-
- Gomon, M. F., Bray, D., & Kuiter, R. H. (2008). *Fishes of Australia's southern coast*. New Holland.
- Govender, R., Abrahamsén-Alami, S., Larsson, A., & Folestad, S. (2020). Therapy for the individual: Towards patient integration into the manufacturing and provision of pharmaceuticals. *European Journal of Pharmaceutics and Biopharmaceutics*, 149, 58–76.
- Gunter, H. M., Fan, S., Xiong, F., Franchini, P., Fruciano, C., & Meyer, A. (2013). Shaping development through mechanical strain: The transcriptional basis of diet-induced phenotypic plasticity in a cichlid fish. *Molecular Ecology*, 22(17), 4516–4531. <https://doi.org/10.1111/mec.12417>
- Gutierrez, A. P., Yáñez, J. M., Fukui, S., Swift, B., & Davidson, W. S. (2015). Genome-wide association study (GWAS) for growth rate and age at sexual maturation in Atlantic salmon (*Salmo salar*). *PloS One*, 10(3), e0119730.
- Han, J.-D. J. (2008). Understanding biological functions through molecular networks. *Cell Research*, 18(2), 224–237. <https://doi.org/10.1038/cr.2008.16>
- Hardcastle, T. J., & Kelly, K. A. (2010). baySeq: Empirical Bayesian methods for identifying differential expression in sequence count data. *BMC Bioinformatics*, 11(1), 1–14.
- Hayes, B. (2013). Overview of statistical methods for genome-wide association studies (GWAS). *Genome-Wide Association Studies and Genomic Prediction*, 149–169.
- He, X., Fuller, C. K., Song, Y., Meng, Q., Zhang, B., Yang, X., & Li, H. (2013). Sherlock: Detecting gene-disease associations by matching patterns of expression QTL and GWAS. *The American Journal of Human Genetics*, 92(5), 667–680.
- Hilario, E. (2015). The Restriction Enzyme Target Approach to Genotyping by Sequencing (GBS). In J. Batley (Ed.), *Plant Genotyping: Methods and Protocols* (pp. 271–279). Springer. https://doi.org/10.1007/978-1-4939-1966-6_20
- Hill, W. G., & Robertson, A. (1966). The effect of linkage on limits to artificial selection. *Genetics Research*, 8(3), 269–294.
- Hochachka, P. W., & Somero, G. N. (2002). *Biochemical adaptation: Mechanism and process in physiological evolution*. Oxford university press.
- Hodgins-Davis, A., & Townsend, J. P. (2009). Evolving gene expression: From G to E to G×E. *Trends in Ecology & Evolution*, 24(12), 649–658. <https://doi.org/10.1016/j.tree.2009.06.011>

- Hoekstra, L. A., Siddiq, M. A., & Montooth, K. L. (2013). Pleiotropic effects of a mitochondrial–nuclear incompatibility depend upon the accelerating effect of temperature in *Drosophila*. *Genetics*, *195*(3), 1129–1139.
- Hoff, K. J., Lomsadze, A., Stanke, M., & Borodovsky, M. (2018). BRAKER2: Incorporating protein homology information into gene prediction with GeneMark-EP and AUGUSTUS. *Plant and Animal Genomes XXVI*.
- Holm, J. Chr., Refstie, T., & Bø, S. (1990). The effect of fish density and feeding regimes on individual growth rate and mortality in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, *89*(3–4), 225–232. [https://doi.org/10.1016/0044-8486\(90\)90128-A](https://doi.org/10.1016/0044-8486(90)90128-A)
- Hong, E. P., & Park, J. W. (2012). Sample Size and Statistical Power Calculation in Genetic Association Studies. *Genomics & Informatics*, *10*(2), 117–122. <https://doi.org/10.5808/GI.2012.10.2.117>
- Houle, D., Govindaraju, D. R., & Omholt, S. (2010). Phenomics: The next challenge. *Nature Reviews Genetics*, *11*(12), 855–866.
- Howard, R., Carriquiry, A. L., & Beavis, W. D. (2014). Parametric and nonparametric statistical methods for genomic selection of traits with additive and epistatic genetic architectures. *G3: Genes, Genomes, Genetics*, *4*(6), 1027–1046.
- Huang, Y., & Agrawal, A. F. (2016). Experimental Evolution of Gene Expression and Plasticity in Alternative Selective Regimes. *PLOS Genetics*, *12*(9), e1006336. <https://doi.org/10.1371/journal.pgen.1006336>
- Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., Bravo, H. C., Davis, S., Gatto, L., & Girke, T. (2015). Orchestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*, *12*(2), 115.
- Huey, R. B., & Kingsolver, J. G. (1989). Evolution of thermal sensitivity of ectotherm performance. *Trends in Ecology & Evolution*, *4*(5), 131–135. [https://doi.org/10.1016/0169-5347\(89\)90211-5](https://doi.org/10.1016/0169-5347(89)90211-5)
- Huisman, J. (2017). Pedigree reconstruction from SNP data: Parentage assignment, sibship clustering and beyond. *Molecular Ecology Resources*, *17*(5), 1009–1024. <https://doi.org/10.1111/1755-0998.12665>
- Hutchings, J. A., Swain, D. P., Rowe, S., Eddington, J. D., Puvanendran, V., & Brown, J. A. (2007). Genetic variation in life-history reaction norms in a marine fish. *Proceedings of the Royal Society B: Biological Sciences*, *274*(1619), 1693–1699. <https://doi.org/10.1098/rspb.2007.0263>

-
- Imslund, A. K., Björnsson, B. T., Gunnarsson, S., Foss, A., & Stefansson, S. O. (2007). Temperature and salinity effects on plasma insulin-like growth factor-I concentrations and growth in juvenile turbot (*Scophthalmus maximus*). *Aquaculture*, 271(1), 546–552. <https://doi.org/10.1016/j.aquaculture.2007.07.007>
- Imslund, A. K., & Jonassen, T. M. (2003). Growth and age at first maturity in turbot and halibut reared under different photoperiods. *Aquaculture International*, 11(5), 463–475. <https://doi.org/10.1023/B:AQUI.00000004191.43885.b2>
- Imslund, A. K., & Jónsdóttir, Ó. D. B. (2002). Is there a genetic basis to growth in Atlantic cod? *Fish and Fisheries*, 3(1), 36–52. <https://doi.org/10.1046/j.1467-2979.2002.00070.x>
- IPCC, Hoegh-Guldberg, O., Jacob, D., Bindi, M., Brown, S., Camilloni, I., Diedhiou, A., Djalante, R., Ebi, K., Engelbrecht, F., & Guiot, J. (2018). Impacts of 1.5 C global warming on natural and human systems. *Global Warming of 1.5 C. An IPCC Special Report*.
- Irwin, S., O'Halloran, J., & FitzGerald, R. D. (1999). Stocking density, growth and growth variation in juvenile turbot, *Scophthalmus maximus* (Rafinesque). *Aquaculture*, 178(1), 77–88. [https://doi.org/10.1016/S0044-8486\(99\)00122-2](https://doi.org/10.1016/S0044-8486(99)00122-2)
- Jamieson, I. G., & Allendorf, F. W. (2012). How does the 50/500 rule apply to MVPs? *Trends in Ecology & Evolution*, 27(10), 578–584.
- Janssen, K., Chavanne, H., Berentsen, P., & Komen, H. (2017). Impact of selective breeding on European aquaculture. *Aquaculture*, 472, 8–16. <https://doi.org/10.1016/j.aquaculture.2016.03.012>
- Jones, G. P. (1986). Food availability affects growth in a coral reef fish. *Oecologia*, 70(1), 136–139.
- Jonsson, B., & Jonsson, N. (2019). Phenotypic plasticity and epigenetics of fish: Embryo temperature affects later-developing life-history traits. *Aquatic Biology*, 28, 21–32.
- Kamps, R., Brandão, R. D., Bosch, B. J., Paulussen, A. D., Xanthouleas, S., Blok, M. J., & Romano, A. (2017). Next-generation sequencing in oncology: Genetic diagnosis, risk prediction and cancer classification. *International Journal of Molecular Sciences*, 18(2), 308.
- Kanehisa, M., & Goto, S. (2000). KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Research*, 28(1), 27–30.

- Karås, P., & Klingsheim, V. (1997). Effects of temperature and salinity on embryonic development of turbot (*Scophthalmus maximus* L.) from the North Sea, and comparisons with Baltic populations. *Helgoländer Meeresuntersuchungen*, 5(2), 241.
- Kardos, M., Luikart, G., & Allendorf, F. W. (2015). Measuring individual inbreeding in the age of genomics: Marker-based measures are better than pedigrees. *Heredity*, 115(1), 63–72. <https://doi.org/10.1038/hdy.2015.17>
- Kasashima, K., & Endo, H. (2015). Interaction of human mitochondrial transcription factor A in mitochondria: Its involvement in the dynamics of mitochondrial DNA nucleoids. *Genes to Cells*, 20(12), 1017–1027.
- Kause, A., Ritola, O., Paananen, T., Wahlroos, H., & Mäntysaari, E. A. (2005). Genetic trends in growth, sexual maturity and skeletal deformations, and rate of inbreeding in a breeding programme for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 247(1), 177–187. <https://doi.org/10.1016/j.aquaculture.2005.02.023>
- Kemp, L. (2020). *A first investigation into the evolutionary relationships, population structure and demographic history of New Zealand Trevally (*Pseudocaranx georgianus*) and its implications for fisheries management*. <http://researcharchive.vuw.ac.nz/handle/10063/9011>
- Kent, J. T. (1983). Information gain and a general measure of correlation. *Biometrika*, 70(1), 163–173. <https://doi.org/10.1093/biomet/70.1.163>
- Kincaid, H. L. (1983). Inbreeding in fish populations used for aquaculture. *Aquaculture*, 33(1–4), 215–227. [https://doi.org/10.1016/0044-8486\(83\)90402-7](https://doi.org/10.1016/0044-8486(83)90402-7)
- Kitano, H. (2002). Systems biology: A brief overview. *Science*, 295(5560), 1662–1664.
- Klinger, D. H., Levin, S. A., & Watson, J. R. (2017). The growth of finfish in global open-ocean aquaculture under climate change. *Proceedings of the Royal Society B: Biological Sciences*, 284(1864), 20170834. <https://doi.org/10.1098/rspb.2017.0834>
- Ko, W.-Y., David, R. M., & Akashi, H. (2003). Molecular Phylogeny of the *Drosophila melanogaster* Species Subgroup. *Journal of Molecular Evolution*, 57(5), 562–573. <https://doi.org/10.1007/s00239-003-2510-x>
- Korte, A., & Farlow, A. (2013). The advantages and limitations of trait analysis with GWAS: A review. *Plant Methods*, 9(1), 1–9.
- Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D., Jones, S. J., & Marra, M. A. (2009). Circos: An information aesthetic for comparative genomics. *Genome Research*, 19(9), 1639–1645. <https://doi.org/10.1101/gr.092759.109>

-
- Last, P. R., White, W. T., Gledhill, D. C., Hobday, A. J., Brown, R., Edgar, G. J., & Pecl, G. (2011). Long-term shifts in abundance and distribution of a temperate fish fauna: A response to climate change and fishing practices. *Global Ecology and Biogeography*, 20(1), 58–72.
- Laurila, A., & Kujasalo, J. (1999). Habitat duration, predation risk and phenotypic plasticity in common frog (*Rana temporaria*) tadpoles. *Journal of Animal Ecology*, 68(6), 1123–1132.
- Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, 15(2), 1–17.
- Le Roith, D., Scavo, L., & Butler, A. (2001). What is the role of circulating IGF-I? *Trends in Endocrinology & Metabolism*, 12(2), 48–52.
- Leal, M. C., Seehausen, O., & Matthews, B. (2017). The Ecology and Evolution of Stoichiometric Phenotypes. *Trends in Ecology & Evolution*, 32(2), 108–117. <https://doi.org/10.1016/j.tree.2016.11.006>
- Lee, S., Lee, Y. M., Kim, K.-H., Kim, H. C., Park, C.-J., Park, J.-W., Noh, G. E., Kim, W.-J., & Hwang, H.-K. (2018). Effects of food availability on growth performance and immune-related gene expression of juvenile olive flounder (*Paralichthys olivaceus*). *Fish & Shellfish Immunology*, 80, 348–356. <https://doi.org/10.1016/j.fsi.2018.06.021>
- Legarra, A., Croiseau, P., Sanchez, M. P., Teyssèdre, S., Sallé, G., Allais, S., Fritz, S., Moreno, C. R., Ricard, A., & Elsen, J.-M. (2015). A comparison of methods for whole-genome QTL mapping using dense markers in four livestock species. *Genetics Selection Evolution*, 47(1), 1–10.
- Leung, P., Lee, C.-S., & O'Bryen, P. J. (2008). *Species and system selection for sustainable aquaculture*. John Wiley & Sons.
- Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*, 27(21), 2987–2993. <https://doi.org/10.1093/bioinformatics/btr509>
- Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, 25(14), 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
- Li, Heng, Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin, R. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, 25(16), 2078–2079.
- Li, J., & Li, M. J. (2012). *Package 'PoissonSeq.'*

- Li, N., Zhou, T., Geng, X., Jin, Y., Wang, X., Liu, S., Xu, X., Gao, D., Li, Q., & Liu, Z. (2018). Identification of novel genes significantly affecting growth in catfish through GWAS analysis. *Molecular Genetics and Genomics*, 293(3), 587–599. <https://doi.org/10.1007/s00438-017-1406-1>
- Lin, W.-Y., Huang, C.-C., Liu, Y.-L., Tsai, S.-J., & Kuo, P.-H. (2019). Genome-wide gene-environment interaction analysis using set-based association tests. *Frontiers in Genetics*, 9, 715.
- Liu, C. (2018). QTL Mapping of Molecular Traits for Studies of Human Complex Diseases. In *Applied Computational Genomics* (pp. 59–80). Springer.
- Liu, F., Sun, F., Xia, J. H., Li, J., Fu, G. H., Lin, G., Tu, R. J., Wan, Z. Y., Quek, D., & Yue, G. H. (2014). A genome scan revealed significant associations of growth traits with a major QTL and GHR2 in tilapia. *Scientific Reports*, 4(1), 1–9.
- Liu, J., Wu, B., Singh, R. P., & Velu, G. (2019). QTL mapping for micronutrients concentration and yield component traits in a hexaploid wheat mapping population. *Journal of Cereal Science*, 88, 57–64. <https://doi.org/10.1016/j.jcs.2019.05.008>
- Liu, X., Huang, M., Fan, B., Buckler, E. S., & Zhang, Z. (2016). Iterative Usage of Fixed and Random Effect Models for Powerful and Efficient Genome-Wide Association Studies. *PLOS Genetics*, 12(2), e1005767. <https://doi.org/10.1371/journal.pgen.1005767>
- Liu, Y., Wang, H., Wen, H., Shi, Y., Zhang, M., Qi, X., Zhang, K., Gong, Q., Li, J., He, F., Hu, Y., & Li, Y. (2020). First High-Density Linkage Map and QTL Fine Mapping for Growth-Related Traits of Spotted Sea bass (*Lateolabrax maculatus*). *Marine Biotechnology*, 22(4), 526–538. <https://doi.org/10.1007/s10126-020-09973-4>
- Loh, P.-R., Tucker, G., Bulik-Sullivan, B. K., Vilhjalmsen, B. J., Finucane, H. K., Salem, R. M., Chasman, D. I., Ridker, P. M., Neale, B. M., & Berger, B. (2015). Efficient Bayesian mixed-model analysis increases association power in large cohorts. *Nature Genetics*, 47(3), 284.
- Löhr, H., & Hammerschmidt, M. (2011). Zebrafish in endocrine systems: Recent advances and implications for human disease. *Annual Review of Physiology*, 73, 183–211.
- Lovatelli, A., Aguilar-Manjarrez, J., & Soto, D. (2013). *Expanding mariculture farther offshore. Technical, environmental, spatial and governance challenges. FAO Technical Workshop, Orbetello, Italy, 22-25 March 2010*. FAO Library.
- Love, M., Anders, S., & Huber, W. (2014). Differential analysis of count data—the DESeq2 package. *Genome Biol*, 15(550), 10–1186.

-
- Lu, J., Tomfohr, J. K., & Kepler, T. B. (2005). Identifying differential expression in multiple SAGE libraries: An overdispersed log-linear model approach. *BMC Bioinformatics*, 6(1), 165. <https://doi.org/10.1186/1471-2105-6-165>
- Lynch, M. (1999). Estimating genetic correlations in natural populations. *Genetical Research*, 74(3), 255–264. <https://doi.org/10.1017/S0016672399004243>
- Lynch, M., & Walsh, B. (1999). *Genetics and Analysis of Quantitative Traits* (Vol. 74). <https://www.journals.uchicago.edu/doi/10.1086/393101>
- Machluf, Y., Gutnick, A., & Levkowitz, G. (2011). Development of the zebrafish hypothalamus. *Annals of the New York Academy of Sciences*, 1220, 93–105. <https://doi.org/10.1111/j.1749-6632.2010.05945.x>
- Mackay, T. F., Stone, E. A., & Ayroles, J. F. (2009). The genetics of quantitative traits: Challenges and prospects. *Nature Reviews Genetics*, 10(8), 565–577.
- Maglott, D., Ostell, J., Pruitt, K. D., & Tatusova, T. (2010). Entrez Gene: Gene-centered information at NCBI. *Nucleic Acids Research*, 39(suppl_1), D52–D57.
- Mallya, Y. J. (2007). The effects of dissolved oxygen on fish growth in aquaculture. *The United Nations University Fisheries Training Programme, Final Project*.
- Marchant, T. A., & Peter, R. E. (1986). Seasonal variations in body growth rates and circulating levels of growth hormone in the goldfish, *Carassius auratus*. *Journal of Experimental Zoology*, 237(2), 231–239. <https://doi.org/10.1002/jez.1402370209>
- Marine, J.-C., Dawson, S.-J., & Dawson, M. A. (2020). Non-genetic mechanisms of therapeutic resistance in cancer. *Nature Reviews Cancer*, 20(12), 743–756.
- McCarthy, D. J., Chen, Y., & Smyth, G. K. (2012). Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research*, 40(10), 4288–4297.
- McGuffin, P., & Huckle, P. (1990). Simulation of Mendelism revisited: The recessive gene for attending medical school. *American Journal of Human Genetics*, 46(5), 994.
- McKinney, W. (2010). *Data Structures for Statistical Computing in Python*. 56–61. <https://doi.org/10.25080/Majora-92bf1922-00a>
- Mell, B., Cheng, X., & Joe, B. (2019). QTL mapping of rat blood pressure loci on RNO1 within a homologous region linked to human hypertension on HSA15. *PLOS ONE*, 14(8), e0221658. <https://doi.org/10.1371/journal.pone.0221658>
- Mendel, G. (1866). Versuche über pflanzenhybriden. Verhandlungen des naturforschenden vereines in brünn, bd. IV für das jahr 1865. *Abhandlungen*, 3–47.

- Metzger, D. C. H., & Schulte, P. M. (2018). Similarities in temperature-dependent gene expression plasticity across timescales in threespine stickleback (*Gasterosteus aculeatus*). *Molecular Ecology*, 27(10), 2381–2396. <https://doi.org/10.1111/mec.14591>
- Metzger, D. C., & Schulte, P. M. (2018). Similarities in temperature-dependent gene expression plasticity across timescales in threespine stickleback (*Gasterosteus aculeatus*). *Molecular Ecology*, 27(10), 2381–2396.
- Meuwissen, T. H. E., Hayes, B. J., & Goddard, M. E. (2001). Prediction of Total Genetic Value Using Genome-Wide Dense Marker Maps. *Genetics*, 157(4), 1819–1829. <https://doi.org/10.1093/genetics/157.4.1819>
- Miranda, A. C. (2017). Mechanisms of behavioural change in urban animals: The role of microevolution and phenotypic plasticity. In *Ecology and conservation of birds in urban environments* (pp. 113–132). Springer.
- Mitchell-Olds, T., Willis, J. H., & Goldstein, D. B. (2007). Which evolutionary processes influence natural genetic variation for phenotypic traits? *Nature Reviews Genetics*, 8(11), 845–856.
- Modig, C., Modesto, T., Canario, A., Cerdà, J., Hofsten, J. von, & Olsson, P.-E. (2006). Molecular Characterization and Expression Pattern of Zona Pellucida Proteins in Gilthead Seabream (*Sparus aurata*)1. *Biology of Reproduction*, 75(5), 717–725. <https://doi.org/10.1095/biolreprod.106.050757>
- Moran, D., Smith, C. K., Gara, B., & Poortenaar, C. W. (2007). Reproductive behaviour and early development in yellowtail kingfish (*Seriola lalandi* Valenciennes 1833). *Aquaculture*, 262(1), 95–104. <https://doi.org/10.1016/j.aquaculture.2006.10.005>
- Morrongiello, J. R., & Thresher, R. E. (2015). A statistical framework to explore ontogenetic growth variation among individuals and populations: A marine fish example. *Ecological Monographs*, 85(1), 93–115. <https://doi.org/10.1890/13-2355.1>
- MPI. (2021). *New Zealand Ministry for Primary Industries Website*. <https://fs.fish.govt.nz/Page.aspx?pk=7&tk=100&sc=TRE>
- Munday, P. L., Kingsford, M. J., O’Callaghan, M., & Donelson, J. M. (2008). Elevated temperature restricts growth potential of the coral reef fish *Acanthochromis polyacanthus*. *Coral Reefs*, 27(4), 927–931. <https://doi.org/10.1007/s00338-008-0393-4>
- Mundy, B. C. (2005). Checklist of the fishes of the Hawaiian Archipelago. *Bishop Mus. Bull. Zool.*, 6, 1–704.

-
- Nadeem, M. A., Nawaz, M. A., Shahid, M. Q., Doğan, Y., Comertpay, G., Yıldız, M., Hatipoğlu, R., Ahmad, F., Alsaleh, A., & Labhane, N. (2018). DNA molecular markers in plant breeding: Current status and recent advancements in genomic selection and genome editing. *Biotechnology & Biotechnological Equipment*, 32(2), 261–285.
- Nett, T. M., Turzillo, A. M., Baratta, M., & Rispoli, L. A. (2002). Pituitary effects of steroid hormones on secretion of follicle-stimulating hormone and luteinizing hormone. *Domestic Animal Endocrinology*, 23(1–2), 33–42.
- Newell, M., Cook, D., Tinker, N., & Jannink, J.-L. (2011). Population structure and linkage disequilibrium in oat (*Avena sativa* L.): Implications for genome-wide association studies. *Theoretical & Applied Genetics*, 122(3), 623–632. <https://doi.org/10.1007/s00122-010-1474-7>
- Nezer, C., Moreau, L., Brouwers, B., Coppieters, W., Dettloux, J., Hanset, R., Karim, L., Kvasz, A., Leroy, P., & Georges, M. (1999). An imprinted QTL with major effect on muscle mass and fat deposition maps to the IGF2 locus in pigs. *Nature Genetics*, 21(2), 155–156.
- Nguyen, N. H., Rastas, P. M. A., Premachandra, H. K. A., & Knibb, W. (2018). First High-Density Linkage Map and Single Nucleotide Polymorphisms Significantly Associated With Traits of Economic Importance in Yellowtail Kingfish *Seriola lalandi*. *Frontiers in Genetics*, 9. <https://doi.org/10.3389/fgene.2018.00127>
- Nguyen, T. T., Bowman, P. J., Haile-Mariam, M., Nieuwhof, G. J., Hayes, B. J., & Pryce, J. E. (2017). Implementation of a breeding value for heat tolerance in Australian dairy cattle. *Journal of Dairy Science*, 100(9), 7362–7367.
- Norbeck, L. A., Kittilson, J. D., & Sheridan, M. A. (2007). Resolving the growth-promoting and metabolic effects of growth hormone: Differential regulation of GH–IGF-I system components. *General and Comparative Endocrinology*, 151(3), 332–341. <https://doi.org/10.1016/j.ygcen.2007.01.039>
- O'Connor, L. J. (2020). The distribution of common-variant effect sizes. *BioRxiv*, 2020.09.19.304097. <https://doi.org/10.1101/2020.09.19.304097>
- O'Connor, L. J., Schoech, A. P., Hormozdiari, F., Gazal, S., Patterson, N., & Price, A. L. (2019). Extreme Polygenicity of Complex Traits Is Explained by Negative Selection. *The American Journal of Human Genetics*, 105(3), 456–476. <https://doi.org/10.1016/j.ajhg.2019.07.003>

- Oppedal, F., Taranger, G. L., Juell, J.-E., Fosseidengen, J. E., & Hansen, T. (1997). Light intensity affects growth and sexual maturation of Atlantic salmon (*Salmo salar*) postsmolts in sea cages. *Aquatic Living Resources*, 10(6), 351–357.
- Orellana, J., Waller, U., & Wecker, B. (2014). Culture of yellowtail kingfish (*Seriola lalandi*) in a marine recirculating aquaculture system (RAS) with artificial seawater. *Aquacultural Engineering*, 58, 20–28.
- Orr, H. A. (1998). The population genetics of adaptation: The distribution of factors fixed during adaptive evolution. *Evolution*, 52(4), 935–949.
- Oshlack, A., Robinson, M. D., & Young, M. D. (2010). From RNA-seq reads to differential expression results. *Genome Biology*, 11(12), 1–10.
- Oshlack, A., & Wakefield, M. J. (2009). Transcript length bias in RNA-seq data confounds systems biology. *Biology Direct*, 4(1), 1–10.
- Pachauri, R. K., Allen, M. R., Barros, V. R., Broome, J., Cramer, W., Christ, R., Church, J. A., Clarke, L., Dahe, Q., Dasgupta, P., Dubash, N. K., Edenhofer, O., Elgizouli, I., Field, C. B., Forster, P., Friedlingstein, P., Fuglestvedt, J., Gomez-Echeverri, L., Hallegatte, S., ... van Ypersele, J.-P. (2014). Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. In R. K. Pachauri & L. Meyer (Eds.), *EPIC3 Geneva, Switzerland, IPCC, 151 p., pp. 151, ISBN: 978-92-9169-143-2* (p. 151). IPCC. <https://epic.awi.de/id/eprint/37530/>
- Palaiokostas, C., Cariou, S., Bestin, A., Bruant, J.-S., Haffray, P., Morin, T., Cabon, J., Allal, F., Vandeputte, M., & Houston, R. D. (2018a). Genome-wide association and genomic prediction of resistance to viral nervous necrosis in European sea bass (*Dicentrarchus labrax*) using RAD sequencing. *Genetics Selection Evolution*, 50(1), 1–11.
- Palaiokostas, C., Cariou, S., Bestin, A., Bruant, J.-S., Haffray, P., Morin, T., Cabon, J., Allal, F., Vandeputte, M., & Houston, R. D. (2018b). Genome-wide association and genomic prediction of resistance to viral nervous necrosis in European sea bass (*Dicentrarchus labrax*) using RAD sequencing. *Genetics Selection Evolution*, 50(1), 30. <https://doi.org/10.1186/s12711-018-0401-2>
- Parisi, M. A., Cramp, R. L., Gordos, M. A., & Franklin, C. E. (2020). Can the impacts of cold-water pollution on fish be mitigated by thermal plasticity? *Conservation Physiology*, 8(coaa005). <https://doi.org/10.1093/conphys/coaa005>
- Parks, B. W., Nam, E., Org, E., Kostem, E., Norheim, F., Hui, S. T., Pan, C., Civelek, M., Rau, C. D., Bennett, B. J., Mehrabian, M., Ursell, L. K., He, A., Castellani, L. W., Zinker,

-
- B., Kirby, M., Drake, T. A., Drevon, C. A., Knight, R., ... Lusi, A. J. (2013). Genetic Control of Obesity and Gut Microbiota Composition in Response to High-Fat, High-Sucrose Diet in Mice. *Cell Metabolism*, 17(1), 141–152. <https://doi.org/10.1016/j.cmet.2012.12.007>
- Parmesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annu. Rev. Ecol. Evol. Syst.*, 37, 637–669.
- Pauly, D. (1980). On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES Journal of Marine Science*, 39(2), 175–192. <https://doi.org/10.1093/icesjms/39.2.175>
- Pedersen, T., & Jobling, M. (1989). Growth rates of large, sexually mature cod *Gadus morhua*, in relation to condition and temperature during an annual cycle. *Aquaculture*, 81(2), 161–168.
- Peltier, E., Sharma, V., Martí Raga, M., Roncoroni, M., Bernard, M., Jiranek, V., Gibon, Y., & Marullo, P. (2018). Dissection of the molecular bases of genotype x environment interactions: A study of phenotypic plasticity of *Saccharomyces cerevisiae* in grape juices. *BMC Genomics*, 19(1), 772. <https://doi.org/10.1186/s12864-018-5145-4>
- Pérez-Sánchez, J., Simó-Mirabet, P., Naya-Català, F., Martos-Sitcha, J. A., Perera, E., Bermejo-Nogales, A., Benedito-Palos, L., & Caldach-Giner, J. A. (2018). Somatotropic Axis Regulation Unravels the Differential Effects of Nutritional and Environmental Factors in Growth Performance of Marine Farmed Fishes. *Frontiers in Endocrinology*, 9. <https://doi.org/10.3389/fendo.2018.00687>
- Perretant, M. R., Cadalen, T., Charmet, G., Sourdille, P., Nicolas, P., Boeuf, C., Tixier, M. H., Branlard, G., & Bernard, S. (2000). QTL analysis of bread-making quality in wheat using a doubled haploid population. *Theoretical and Applied Genetics*, 100(8), 1167–1175.
- Perry, R. W., Plumb, J. M., & Huntington, C. W. (2015). Using a Laboratory-Based Growth Model to Estimate Mass- and Temperature-Dependent Growth Parameters across Populations of Juvenile Chinook Salmon. *Transactions of the American Fisheries Society*, 144(2), 331–336. <https://doi.org/10.1080/00028487.2014.996667>
- Picha, M. E., Turano, M. J., Beckman, B. R., & Borski, R. J. (2008). Endocrine biomarkers of growth and applications to aquaculture: A minireview of growth hormone, insulin-like growth factor (IGF)-I, and IGF-binding proteins as potential growth indicators in fish. *North American Journal of Aquaculture*, 70(2), 196–211.

- Pirozzi, I., & Booth, M. A. (2009). The routine metabolic rate of mulloay (*Argyrosomus japonicus*: Sciaenidae) and yellowtail kingfish (*Seriola lalandi*: Carangidae) acclimated to six different temperatures. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 152(4), 586–592. <https://doi.org/10.1016/j.cbpa.2009.01.008>
- Poortenaar, C. W., Hooker, S. H., & Sharp, N. (2001). Assessment of yellowtail kingfish (*Seriola lalandi lalandi*) reproductive physiology, as a basis for aquaculture development. *Aquaculture*, 201(3–4), 271–286.
- Pörtner, H. O., & Farrell, A. P. (2008). Physiology and climate change. *Science*, 690–692.
- Posbergh, C. J., & Huson, H. J. (2021). All sheeps and sizes: A genetic investigation of mature body size across sheep breeds reveals a polygenic nature. *Animal Genetics*, 52(1), 99–107. <https://doi.org/10.1111/age.13016>
- Premachandra, H. K. A., Nguyen, N. H., Miller, A., D’Antignana, T., & Knibb, W. (2017). Genetic parameter estimates for growth and non-growth traits and comparison of growth performance in sea cages vs land tanks for yellowtail kingfish *Seriola lalandi*. *Aquaculture*, 479, 169–175. <https://doi.org/10.1016/j.aquaculture.2017.05.043>
- Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*, 81(3), 559–575. <https://doi.org/10.1086/519795>
- R. Core Team. (2013). *R: A language and environment for statistical computing*.
- Rashwan, A. M. A. (2016). Comparative study in fifteen genotypes of tomato for heat tolerance under Upper Egypt conditions. *Journal of American Science*, 12(6), 68–76.
- Rastas, P. (2017). Lep-MAP3: Robust linkage mapping even for low-coverage whole genome sequencing data. *Bioinformatics*, 33(23), 3726–3732. <https://doi.org/10.1093/bioinformatics/btx494>
- Réale, D., McAdam, A. G., Boutin, S., & Berteaux, D. (2003). Genetic and plastic responses of a northern mammal to climate change. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 270(1515), 591–596.
- Reinecke, M. (2010). Influences of the environment on the endocrine and paracrine fish growth hormone–insulin-like growth factor-I system. *Journal of Fish Biology*, 76(6), 1233–1254. <https://doi.org/10.1111/j.1095-8649.2010.02605.x>
- Risch, N., & Merikangas, K. (1996). The future of genetic studies of complex human diseases. *Science*, 273(5281), 1516–1517.

-
- Risch, N., Spiker, D., Lotspeich, L., Nouri, N., Hinds, D., Hallmayer, J., Kalaydjieva, L., McCague, P., Dimiceli, S., & Pitts, T. (1999). A genomic screen of autism: Evidence for a multilocus etiology. *The American Journal of Human Genetics*, 65(2), 493–507.
- Roberts, C., Stewart, A. L., & Struthers, C. D. (2015). *The Fishes of New Zealand*. Te Papa Press.
- Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, 26(1), 139–140.
- Robinson, M. D., & Oshlack, A. (2010). A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biology*, 11(3), 1–9.
- Robinson, M. D., & Smyth, G. K. (2007). Moderated statistical tests for assessing differences in tag abundance. *Bioinformatics*, 23(21), 2881–2887.
- Robinson, M. R., Wray, N. R., & Visscher, P. M. (2014). Explaining additional genetic variation in complex traits. *Trends in Genetics*, 30(4), 124–132. <https://doi.org/10.1016/j.tig.2014.02.003>
- Rockman, M. V. (2012). The QTN program and the alleles that matter for evolution: All that's gold does not glitter. *Evolution: International Journal of Organic Evolution*, 66(1), 1–17.
- Rockman, M. V., & Kruglyak, L. (2006). Genetics of global gene expression. *Nature Reviews Genetics*, 7(11), 862–872.
- Roff, D. A., & Fairbairn, D. J. (2007). The evolution of trade-offs: Where are we? *Journal of Evolutionary Biology*, 20(2), 433–447.
- Rousseau, K., & Dufour, S. (2007). Comparative aspects of GH and metabolic regulation in lower vertebrates. *Neuroendocrinology*, 86(3), 165–174.
- Ruigrok, M., Catanach, A., Bowatte, D., Davey, M., Storey, R., Valenza-Troubat, N., Lopez-Girona, E., Hilario, E., Wylie, M. J., Chagne, D., & Wellenreuther, M. (2021). The genome of New Zealand trevally (Carangidae: *Pseudocaranx georgianus*) uncovers a XY sex determination locus. *BioRxiv*, 2021.04.25.441282. <https://doi.org/10.1101/2021.04.25.441282>
- Sae-Lim, P., Gjerde, B., Nielsen, H. M., Mulder, H., & Kause, A. (2016). A review of genotype-by-environment interaction and micro-environmental sensitivity in aquaculture species. *Reviews in Aquaculture*, 8(4), 369–393.

- Sae-Lim, P., Mulder, H., Gjerde, B., Koskinen, H., Lillehammer, M., & Kause, A. (2015). Genetics of Growth Reaction Norms in Farmed Rainbow Trout. *PLOS ONE*, 10(8), e0135133. <https://doi.org/10.1371/journal.pone.0135133>
- Salles, O. C., Pujol, B., Maynard, J. A., Almany, G. R., Berumen, M. L., Jones, G. P., Saenz-Agudelo, P., Srinivasan, M., Thorrold, S. R., & Planes, S. (2016). First genealogy for a wild marine fish population reveals multigenerational philopatry. *Proceedings of the National Academy of Sciences*, 113(46), 13245–13250. <https://doi.org/10.1073/pnas.1611797113>
- Sandblom, E., Gräns, A., Axelsson, M., & Seth, H. (2014). Temperature acclimation rate of aerobic scope and feeding metabolism in fishes: Implications in a thermally extreme future. *Proceedings of the Royal Society B: Biological Sciences*, 281(1794), 20141490.
- Santure, A. W., & Garant, D. (2018). Wild GWAS—association mapping in natural populations. *Molecular Ecology Resources*, 18(4), 729–738. <https://doi.org/10.1111/1755-0998.12901>
- Schaal, S. M., & Lotterhos, K. E. (2021). Comparative thermal performance among four young-of-the-year temperate reef fish species. *ICES Journal of Marine Science*. <https://doi.org/10.1093/icesjms/fsab072>
- Schally, A. V., Arimura, A., Baba, Y., Nair, R. M. G., Matsuo, H., Redding, T. W., Debeljuk, L., & White, W. F. (1971). Isolation and properties of the FSH and LH-releasing hormone. *Biochemical and Biophysical Research Communications*, 43(2), 393–399.
- Scheiner, S. M. (1993). Genetics and Evolution of Phenotypic Plasticity. *Annual Review of Ecology and Systematics*, 24(1), 35–68. <https://doi.org/10.1146/annurev.es.24.110193.000343>
- Scheiner, S. M., & Goodnight, C. J. (1984). The Comparison of Phenotypic Plasticity and Genetic Variation in Populations of the Grass *Danthonia spicata*. *Evolution*, 38(4), 845–855. <https://doi.org/10.2307/2408395>
- Schielzeth, H., & Husby, A. (2014). *Challenges and prospects in genome wide QTL mapping of standing genetic variation in natural populations*.
- Schlichting, C. D., & Pigliucci, M. (1993). Control of Phenotypic Plasticity Via Regulatory Genes. *The American Naturalist*, 142(2), 366–370. <https://doi.org/10.1086/285543>
- Schulte, P. M., Healy, T. M., & Fangue, N. A. (2011). Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology*, 51(5), 691–702.

-
- Sebens, K. P. (1987). The Ecology of Indeterminate Growth in Animals. *Annual Review of Ecology and Systematics*, 18(1), 371–407.
<https://doi.org/10.1146/annurev.es.18.110187.002103>
- Seibel, B. A., & Drazen, J. C. (2007). The rate of metabolism in marine animals: Environmental constraints, ecological demands and energetic opportunities. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 362(1487), 2061–2078.
- Shen, Y., & Yue, G. (2019). Current status of research on aquaculture genetics and genomics-information from ISGA 2018. *Aquaculture and Fisheries*, 4(2), 43–47.
<https://doi.org/10.1016/j.aaf.2018.11.001>
- Shi, L., Yang, Y., Xie, Q., Miao, H., Bo, K., Song, Z., Wang, Y., Xie, B., Zhang, S., & Gu, X. (2018). Inheritance and QTL mapping of cucumber mosaic virus resistance in cucumber (*Cucumis Sativus* L.). *PloS One*, 13(7), e0200571.
- Shi, Y., Zhou, Z., Liu, B., Kong, S., Chen, B., Bai, H., Li, L., Pu, F., & Xu, P. (2020). Construction of a High-Density Genetic Linkage Map and QTL Mapping for Growth-Related Traits in *Takifugu bimaculatus*. *Marine Biotechnology*, 22(1), 130–144.
<https://doi.org/10.1007/s10126-019-09938-2>
- Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, 31(19), 3210–3212.
- Sinnott-Armstrong, N., Naqvi, S., Rivas, M., & Pritchard, J. K. (2021). GWAS of three molecular traits highlights core genes and pathways alongside a highly polygenic background. *ELife*, 10, e58615. <https://doi.org/10.7554/eLife.58615>
- Skaramuca, B., Kožul, V., Teskeredžić, Z., Bolotin, J., & Onofri, V. (2001). Growth rate of tank-reared Mediterranean amberjack, *Seriola dumerili* (Risso 1810) fed on three different diets. *Journal of Applied Ichthyology*, 17(3), 130–133.
- Slate, J. (2005). INVITED REVIEW: Quantitative trait locus mapping in natural populations: progress, caveats and future directions. *Molecular Ecology*, 14(2), 363–379.
<https://doi.org/10.1111/j.1365-294X.2004.02378.x>
- Slepecky, R. A., & Starmer, W. T. (2009). Phenotypic plasticity in fungi: A review with observations on *Aureobasidium pullulans*. *Mycologia*, 101(6), 823–832.
- Smit, A. F. (2004). Repeat-Masker Open-3.0. [Http://Www. Repeatmasker. Org](http://www.Repeatmasker.Org).
- Smit, A. F. A., Hubley, R., & Green, P. (2015). RepeatModeler Open-1.0. 2008–2015. *Seattle, USA: Institute for Systems Biology*. Available from: [Httpwww. Repeatmasker. Org](http://www.Repeatmasker.Org), Last Accessed May, 1, 2018.

- Smith, S., Bernatchez, L., & Beheregaray, L. B. (2013). RNA-seq analysis reveals extensive transcriptional plasticity to temperature stress in a freshwater fish species. *BMC Genomics*, 14(1), 375. <https://doi.org/10.1186/1471-2164-14-375>
- Smith-Vaniz, W. F., & Jelks, H. L. (2006). Australian trevallies of the genus *Pseudocaranx* (Teleostei: Carangidae), with description of a new species from Western Australia. *Memoirs of Museum Victoria*, 63(1), 97–106. <https://doi.org/10.24199/j.mmv.2006.63.12>
- Smyth, G. K. (2004). Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, 3(1).
- Smyth, G. K., & Verbyla, A. P. (1996). A conditional likelihood approach to residual maximum likelihood estimation in generalized linear models. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(3), 565–572.
- Sonah, H., O'Donoghue, L., Cober, E., Rajcan, I., & Belzile, F. (2015). Identification of loci governing eight agronomic traits using a GBS-GWAS approach and validation by QTL mapping in soya bean. *Plant Biotechnology Journal*, 13(2), 211–221.
- Su, S., Li, H., Du, F., Zhang, C., Li, X., Jing, X., Liu, L., Li, Z., Yang, X., Xu, P., Yuan, X., Zhu, J., & Bouzoualegh, R. (2018). Combined QTL and Genome Scan Analyses With the Help of 2b-RAD Identify Growth-Associated Genetic Markers in a New Fast-Growing Carp Strain. *Frontiers in Genetics*, 9. <https://doi.org/10.3389/fgene.2018.00592>
- Sultan, S. E. (2000). Phenotypic plasticity for plant development, function and life history. *Trends in Plant Science*, 5(12), 537–542. [https://doi.org/10.1016/S1360-1385\(00\)01797-0](https://doi.org/10.1016/S1360-1385(00)01797-0)
- Sun, C., Niu, Y., Ye, X., Dong, J., Hu, W., Zeng, Q., Chen, Z., Tian, Y., Zhang, J., & Lu, M. (2017). Construction of a high-density linkage map and mapping of sex determination and growth-related loci in the mandarin fish (*Siniperca chuatsi*). *BMC Genomics*, 18(1), 446. <https://doi.org/10.1186/s12864-017-3830-3>
- Supek, F., Bošnjak, M., Škunca, N., & Šmuc, T. (2011). REVIGO summarizes and visualizes long lists of gene ontology terms. *PloS One*, 6(7), e21800.
- Swindell, W. R., Huebner, M., & Weber, A. P. (2007). Plastic and adaptive gene expression patterns associated with temperature stress in *Arabidopsis thaliana*. *Heredity*, 99(2), 143–150.

-
- Tan, K., Zhang, H., & Zheng, H. (2020). Selective breeding of edible bivalves and its implication of global climate change. *Reviews in Aquaculture*, 12(4), 2559–2572.
- Thoday, J. M., & Thompson, J. N. (1976). The number of segregating genes implied by continuous variation. *Genetica*, 46(3), 335–344.
- Thompson, R., Crump, R., Juga, J., & Visscher, P. (1995). Estimating variances and covariances for bivariate animal models using scaling and transformation. *Genetics Selection Evolution*, 27(1), 33–42.
- Thorland, I., Thodesen, J., Refstie, T., Folkedal, O., Stien, L. H., Nilsson, J., Seim, R. R., Kristiansen, T. S., & Rye, M. (2020). Genetic variation in growth pattern within a population of farmed Atlantic salmon (*Salmo salar*) during a standard production cycle. *Aquaculture*, 518, 734735. <https://doi.org/10.1016/j.aquaculture.2019.734735>
- Thorpe, J. E. (1994). Performance Thresholds and Life-History Flexibility in Salmonids. *Conservation Biology*, 8(3), 877–879.
- Tollrian, R., & Heibl, C. (2004). Phenotypic plasticity in pigmentation in *Daphnia* induced by UV radiation and fish kairomones. *Functional Ecology*, 18(4), 497–502.
- Toolkit, P. (2015). *Broad Institute, GitHub. Repository. 2019.*
- Torkamaneh, D., Boyle, B., St-Cyr, J., Légaré, G., Pomerleau, S., & Belzile, F. (2020). NanoGBS: A Miniaturized Procedure for GBS Library Preparation. *Frontiers in Genetics*, 11. <https://doi.org/10.3389/fgene.2020.00067>
- Torres-Contreras, A. M., Senés-Guerrero, C., Pacheco, A., González-Agüero, M., Ramos-Parra, P. A., Cisneros-Zevallos, L., & Jacobo-Velázquez, D. A. (2018). Genes differentially expressed in broccoli as an early and late response to wounding stress. *Postharvest Biology and Technology*, 145, 172–182.
- Trapnell, C., Hendrickson, D. G., Sauvageau, M., Goff, L., Rinn, J. L., & Pachter, L. (2013). Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nature Biotechnology*, 31(1), 46–53.
- Triantaphyllopoulos, K. A., Cartas, D., & Miliou, H. (2020). Factors influencing GH and IGF-I gene expression on growth in teleost fish: How can aquaculture industry benefit? *Reviews in Aquaculture*, 12(3), 1637–1662. <https://doi.org/10.1111/raq.12402>
- Tsai, H. Y., Hamilton, A., Guy, D. R., Tinch, A. E., Bishop, S. C., & Houston, R. D. (2015). The genetic architecture of growth and fillet traits in farmed Atlantic salmon (*Salmo salar*). *BMC Genetics*, 16(1), 51. <https://doi.org/10.1186/s12863-015-0215-y>
- Van Laere, A.-S., Nguyen, M., Braunschweig, M., Nezer, C., Collette, C., Moreau, L., Archibald, A. L., Haley, C. S., Buys, N., & Tally, M. (2003). A regulatory mutation in

- IGF2 causes a major QTL effect on muscle growth in the pig. *Nature*, 425(6960), 832–836.
- Veilleux, H. D., Ryu, T., Donelson, J. M., van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen, M. L., Leggat, W., Ravasi, T., & Munday, P. L. (2015). Molecular processes of transgenerational acclimation to a warming ocean. *Nature Climate Change*, 5(12), 1074–1078. <https://doi.org/10.1038/nclimate2724>
- Vieira, C., Pasyukova, E. G., Zeng, Z.-B., Hackett, J. B., Lyman, R. F., & Mackay, T. F. C. (2000). Genotype-Environment Interaction for Quantitative Trait Loci Affecting Life Span in *Drosophila melanogaster*. *Genetics*, 154(1), 213–227. <https://doi.org/10.1093/genetics/154.1.213>
- Visser, P., Hill, W., & Wray, N. (2008). Heritability in the genomics era-concepts and misconceptions. *Nat Rev Genet* 9: 255-266. *Nature Reviews. Genetics*, 9, 255–266. <https://doi.org/10.1038/nrg2322>
- Visser, P. M., & Goddard, M. E. (2019). From RA Fisher's 1918 paper to GWAS a century later. *Genetics*, 211(4), 1125–1130.
- Voorrips, R. E. (2002). MapChart: Software for the Graphical Presentation of Linkage Maps and QTLs. *Journal of Heredity*, 93(1), 77–78. <https://doi.org/10.1093/jhered/93.1.77>
- Waldvogel, A.-M., Feldmeyer, B., Rolshausen, G., Exposito-Alonso, M., Rellstab, C., Kofler, R., Mock, T., Schmid, K., Schmitt, I., & Bataillon, T. (2020). Evolutionary genomics can improve prediction of species' responses to climate change. *Evolution Letters*, 4(1), 4–18.
- Wang, J., & Zhang, Z. (2020). *GAPIT Version 3: Boosting Power and Accuracy for Genomic Association and Prediction* [Preprint]. Bioinformatics. <https://doi.org/10.1101/2020.11.29.403170>
- Wang, M., Wang, Q., & Zhang, B. (2013). Evaluation and selection of reliable reference genes for gene expression under abiotic stress in cotton (*Gossypium hirsutum* L.). *Gene*, 530(1), 44–50.
- Wang, S., Hard, J. J., & Utter, F. (2002). Salmonid inbreeding: A review. *Reviews in Fish Biology and Fisheries*, 11(4), 301–319. <https://doi.org/10.1023/A:1021330500365>
- Wang, X., Jiang, S., Zheng, L., Xiao, L., Zhang, X., Wang, D., Li, S., Shi, Q., Wu, S., Lin, H., You, X., & Zhang, Y. (2019). An SNP-Based Genetic Map and QTL Mapping for Growth Traits in the Red-Spotted Grouper (*Epinephelus akaara*). *Genes*, 10(10), 793. <https://doi.org/10.3390/genes10100793>

-
- Wang, Z., Gerstein, M., & Snyder, M. (2009). RNA-Seq: A revolutionary tool for transcriptomics. *Nature Reviews Genetics*, 10(1), 57–63.
- Wang, Z., Liu, T., Lin, Z., Hegarty, J., Koltun, W. A., & Wu, R. (2010). A general model for multilocus epistatic interactions in case-control studies. *PLoS One*, 5(8), e11384.
- Waterhouse, R. M., Seppey, M., Simão, F. A., Manni, M., Ioannidis, P., Klioutchnikov, G., Kriventseva, E. V., & Zdobnov, E. M. (2018). BUSCO applications from quality assessments to gene prediction and phylogenomics. *Molecular Biology and Evolution*, 35(3), 543–548.
- Watson, J. D. (1970). Molecular biology of the gene. *Molecular Biology of the Gene.*, 2nd edn.
- Weatherley, A. H. (1972). *Growth and Ecology of Fish Populations* (Academic Press Inc.).
- Weedon, M. N., Lango, H., Lindgren, C. M., Wallace, C., Evans, D. M., Mangino, M., Freathy, R. M., Perry, J. R. B., Stevens, S., Hall, A. S., Samani, N. J., Shields, B., Prokopenko, I., Farrall, M., Dominiczak, A., Johnson, T., Bergmann, S., Beckmann, J. S., Vollenweider, P., ... Frayling, T. M. (2008). Genome-wide association analysis identifies 20 loci that influence adult height. *Nature Genetics*, 40(5), 575–583. <https://doi.org/10.1038/ng.121>
- Wei, X., & Zhang, J. (2017). The Genomic Architecture of Interactions Between Natural Genetic Polymorphisms and Environments in Yeast Growth. *Genetics*, 205(2), 925–937. <https://doi.org/10.1534/genetics.116.195487>
- Wellenreuther, M., & Hansson, B. (2016). Detecting Polygenic Evolution: Problems, Pitfalls, and Promises. *Trends in Genetics*, 32(3), 155–164. <https://doi.org/10.1016/j.tig.2015.12.004>
- Wellenreuther, M., Le Luyer, J., Cook, D., Ritchie, P. A., & Bernatchez, L. (2019). Domestication and Temperature Modulate Gene Expression Signatures and Growth in the Australasian Snapper *Chrysophrys auratus*. *G3 Genes/Genomes/Genetics*, 9(1), 105–116. <https://doi.org/10.1534/g3.118.200647>
- West-Eberhard, M. J. (1989). Phenotypic Plasticity and the Origins of Diversity. *Annual Review of Ecology and Systematics*, 20(1), 249–278. <https://doi.org/10.1146/annurev.es.20.110189.001341>
- Whatmore, P., Nguyen, N. H., Miller, A., Lamont, R., Powell, D., D'Antignana, T., Bubner, E., Elizur, A., & Knibb, W. (2013). Genetic parameters for economically important traits in yellowtail kingfish *Seriola lalandi*. *Aquaculture*, 400–401, 77–84. <https://doi.org/10.1016/j.aquaculture.2013.03.002>

- Wieters, B., Steige, K. A., He, F., Koch, E. M., Ramos-Onsins, S. E., Gu, H., Guo, Y.-L., Sunyaev, S., & Meaux, J. de. (2021). Polygenic adaptation of rosette growth in *Arabidopsis thaliana*. *PLOS Genetics*, 17(1), e1008748. <https://doi.org/10.1371/journal.pgen.1008748>
- Wilson, A. J., Réale, D., Clements, M. N., Morrissey, M. M., Postma, E., Walling, C. A., Kruuk, L. E. B., & Nussey, D. H. (2010). An ecologist's guide to the animal model. *Journal of Animal Ecology*, 79(1), 13–26. <https://doi.org/10.1111/j.1365-2656.2009.01639.x>
- Wolak, M. E., & Reid, J. M. (2017). Accounting for genetic differences among unknown parents in microevolutionary studies: How to include genetic groups in quantitative genetic animal models. *Journal of Animal Ecology*, 86(1), 7–20.
- Woodin, S. A., Hilbish, T. J., Helmuth, B., Jones, S. J., & Wetthey, D. S. (2013). Climate change, species distribution models, and physiological performance metrics: Predicting when biogeographic models are likely to fail. *Ecology and Evolution*, 3(10), 3334–3346.
- Wray, N., & Visscher, P. (2008). *Estimating trait heritability*. https://scholar.google.com/scholar_lookup?title=Estimating%20trait%20heritability&publication_year=2008&author=N.%20Wray&author=P.M.%20Visscher
- Wu, H., Wang, C., & Wu, Z. (2013). A new shrinkage estimator for dispersion improves differential expression detection in RNA-seq data. *Biostatistics*, 14(2), 232–243.
- Wyrick, J. J., & Young, R. A. (2002). Deciphering gene expression regulatory networks. *Current Opinion in Genetics & Development*, 12(2), 130–136. [https://doi.org/10.1016/S0959-437X\(02\)00277-0](https://doi.org/10.1016/S0959-437X(02)00277-0)
- Yamahira, K., & Conover, D. O. (2002). Intra-vs. interspecific latitudinal variation in growth: Adaptation to temperature or seasonality? *Ecology*, 83(5), 1252–1262.
- Yang J, Manolio TA, Pasquale LR, Boerwinkle E, Caporaso N, Cunningham JM, de Andrade M, Feenstra B, Feingold E, Hayes MG, Hill WG, Landi MT, Alonso A, Lettre G, Lin P, Ling H, Lowe W, Mathias RA, Melbye M, Pugh E, Cornelis MC, Weir BS, Goddard ME, Visscher PM. (2011). Genome partitioning of genetic variation for complex traits using common SNPs. Yang, Jian et al. “Genome partitioning of genetic variation for complex traits using common SNPs.” *Nature genetics* vol. 43,6 (2011): 519-25. doi:10.1038/ng.823.

-
- Yang, D., Liu, Y., Cheng, H., Chang, L., Chen, J., Chai, S., & Li, M. (2016). Genetic dissection of flag leaf morphology in wheat (*Triticum aestivum* L.) under diverse water regimes. *BMC Genetics*, 17(1), 94. <https://doi.org/10.1186/s12863-016-0399-9>
- Yang, Y., Wu, L., Wu, X., Li, B., Huang, W., Weng, Z., Lin, Z., Song, L., Guo, Y., Meng, Z., Liu, X., & Xia, J. (2020). Identification of Candidate Growth-Related SNPs and Genes Using GWAS in Brown-Marbled Grouper (*Epinephelus fuscoguttatus*). *Marine Biotechnology*, 22(2), 153–166. <https://doi.org/10.1007/s10126-019-09940-8>
- Yasumoto, T. I., Nakatsukasa, M., Nagano, A. J., Yasugi, M., Yoshimura, T., & Shinomiya, A. (2020). Genetic analysis of body weight in wild populations of medaka fish from different latitudes. *Plos One*, 15(6), e0234803.
- Ye, B., Wan, Z., Wang, L., Pang, H., Wen, Y., Liu, H., Liang, B., Lim, H. S., Jiang, J., & Yue, G. (2017). Heritability of growth traits in the Asian seabass (*Lates calcarifer*). *Aquaculture and Fisheries*, 2(3), 112–118. <https://doi.org/10.1016/j.aaf.2017.06.001>
- Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology analysis for RNA-seq: Accounting for selection bias. *Genome Biology*, 11(2), 1–12.
- Yuan, Y., Peng, D., Gu, X., Gong, Y., Sheng, Z., & Hu, X. (2018). Polygenic Basis and Variable Genetic Architectures Contribute to the Complex Nature of Body Weight—A Genome-Wide Study in Four Chinese Indigenous Chicken Breeds. *Frontiers in Genetics*, 9. <https://doi.org/10.3389/fgene.2018.00229>
- Zenger, K. R., Khatkar, M. S., Jerry, D. R., & Raadsma, H. W. (2017). *THE NEXT WAVE IN SELECTIVE BREEDING: IMPLEMENTING GENOMIC SELECTION IN AQUACULTURE*. 8.
- Zhang, Yan, Qi, G., Park, J.-H., & Chatterjee, N. (2018). Estimation of complex effect-size distributions using summary-level statistics from genome-wide association studies across 32 complex traits. *Nature Genetics*, 50(9), 1318–1326. <https://doi.org/10.1038/s41588-018-0193-x>
- Zhang, Yangfan, Timmerhaus, G., Anttila, K., Mauduit, F., Jørgensen, S. M., Kristensen, T., Claireaux, G., Takle, H., & Farrell, A. P. (2016). Domestication compromises athleticism and respiratory plasticity in response to aerobic exercise training in Atlantic salmon (*Salmo salar*). *Aquaculture*, 463, 79–88.
- Zhang, Yifeng, Zhang, J., Gong, H., Cui, L., Zhang, W., Ma, J., Chen, C., Ai, H., Xiao, S., & Huang, L. (2019). Genetic correlation of fatty acid composition with growth, carcass, fat deposition and meat quality traits based on GWAS data in six pig populations. *Meat Science*, 150, 47–55.

- Zhou, Z., Li, Y., Zhang, G., Ye, H., & Luo, J. (2020). Effects of temperature on the transcriptomes of pituitary and liver in Golden Pompano *Trachinotus blochii*. *Fish Physiology and Biochemistry*, 46(1), 63–73. <https://doi.org/10.1007/s10695-019-00695-6>
- Zhu, C., Liu, H., Pan, Z., Chang, G., Wang, H., Wu, N., Ding, H., & Yu, X. (2019). Construction of a high-density genetic linkage map and QTL mapping for growth traits in *Pseudobagrus ussuriensis*. *Aquaculture*, 511, 734213. <https://doi.org/10.1016/j.aquaculture.2019.734213>
- Zuk, O., Hechter, E., Sunyaev, S. R., & Lander, E. S. (2012). The mystery of missing heritability: Genetic interactions create phantom heritability. *Proceedings of the National Academy of Sciences*, 109(4), 1193–1198.