Hoki (*Macruronus novaezelandiae*) diet variability and
associated middle-depth demersal fish species depth
distribution in the ecosystem on the
Chatham Rise, New Zealand.

Amelia MacLeod Connell

A thesis submitted to the Victoria University of Wellington in fulfilment of the
requirements for the degree of Masters of Science in Marine Biology.

Victoria University of Wellington

2009

ABSTRACT

Fisheries management in New Zealand is mostly on a single species basis. Globally there is a shift towards multispecies or ecosystem based fisheries management. For this to happen an understanding of how the ecosystem is organised and functions is needed. Trophic food web and diet studies have been used effectively to begin to understand the functioning of marine ecosystems. Who eats whom, however, is not the full extent of ecosystem function. Understanding of species distribution patterns, of both predators and prey species are also needed to begin to understand the full function of the marine ecosystem.

The first part of this study investigated the diet of hoki (*Macruronus novaezelandiae*) over the Chatham Rise, New Zealand, between 200-800m. It characterised the diet of hoki as well as investigated potential sources of diet variability. Hoki diet was found to consist largely of mesopelagic teleosts, mainly of the family Myctophidae, natant decapods and euphausiids, suggesting a pelagic feeding strategy, as other studies have also found. Differences were found in diet composition between this study and other studies on hoki diet, potentially suggesting differences in prey distribution between study areas. Differences in diet were found between fish from different depths and different sized fish from the same depth. No consistent pattern of diet differences was found between the different areas studied, suggesting that the patterns found may be aliasing depth and size patterns as well as reflecting differences in hoki size class distribution. The distribution of hoki was not homogeneous over the study area, with small fish found mainly in the western part of the study area in shallower water, while large fish were predominately found at greater depths over the whole study area.

The second part of this study looked at the overall species distribution of 30 demersal fish species over the Chatham Rise, specifically examining for evidence of the mid-domain effect. The study also investigated body-size depth trends between these species, and split by class Osteichthyes and Chondrichthyes. The mid-domain effect predicts species richness, and thus distribution, is due to geometric constraints with the greatest species richness to be found at the centre of a geographically constrained domain. The overall

species distribution was found to be explained by the mid-domain effect. The distribution pattern of larger individuals being found in deeper water, with smaller individuals found in shallower water has often been seen in marine systems. We found no interspecific pattern for body-size depth distribution with the entire species assemblage, nor when the assemblage was split by class into Osteichthyes and Chondrichthyes, which supports our findings of the mid-domain effect. At a species level patterns of positive, negative and no trend were found with body-size depth relationships. At a community level species distribution over the studied depth range was largely explained by the geometric constraints of the mid-domain effect, while at a species level distribution over depth was often a reflection of body size. Some species had large individuals deep while other species had small. Overall this supports the hypothesis that competition or adaptation works more strongly at a population or species level, than on the overall community who's species distribution can more often be attributed to random chance.

This study begins to explain predator species distribution over the Chatham Rise and looks at the diet of one dominant species in the Chatham Rise ecosystem. This provides some of the basic knowledge needed for fisheries management to move towards a more ecosystem based approach. Further research should include investigation into prey species distribution and abundance to clarify some of the questions raised in the diet part of this study about the cause of diet variability and whether it was related to either prey abundance or patchy prey distribution. Research into the diet of other fish would be useful to ascertain which species compete with hoki for food and would provide fisheries managers with a list of species that may be affected indirectly through changes in hoki quota of abundance.

ACKNOWLEDGEMENTS

First and foremost this project would not have gone ahead without the funding and support of the Ministry of Fisheries and the National Institute of Water and Atmospheric Research Ltd. (NIWA). The data used for this study were collected as part of a Ministry of Fisheries funded project, ZBD2004-02 *Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise*, carried out by NIWA.

A big thank you goes to my supervisors, Drs Matthew Dunn (NIWA), Phil Lester (VUW) and KC Burns (VUW), who have provided me with guidance, support and constructive comments throughout this study.

- Thank you Matt for having an open door to answer even my most trivial of questions, as well as having patience when explaining statistical concepts to me and giving me a huge amount of help with R programming.
- Phil and KC, thank you for forcing me to think, to push myself, and ultimately go further with the theoretical side of this project than I ever thought possible.

Thank you to all the wonderful friendly and helpful people I have met and worked with at NIWA you have really made my time here great. Specifically I need to thank:

- Darren Stevens who helped teach me the ropes with fish gut dissection, and helped with many species identifications.
- Peter McMillian, your expert skills of identification when presented with semi digested fish have been invaluable,
- Neil Bagley your treasure trove of historical research and help with fish species identification has been most helpful,
- Brent Wood, you have answered many of my questions and solved many of my problems with the database used to hold the stomach content information, thank you,
- Graham Rickard, you provided me with the current and temperature maps of the Chatham Rise,
- Rick Webber (Te Papa), Yves Cherel (CEBC – CNRS, France), Graham Fenwick, Kareen Schnabel, Shane Ayhoun, and Kelly Merrin for all your crustacean identification skills, thank you,
- Paul Grimes and Don Robertson your help with Myctophid identification and distribution has been particularly valuable, and
- The Masters and crew of *RV Tangaroa* without your support the at sea part of this project would not have been as successful,

And finally, my family, my good friends and my partner deserve an enormous amount of thanks. You have all given me never ending support, particularly when things have been tough, and been understanding during my periods of absence. Without you I would not have gotten through.

Table of Contents

1 GENERAL INTRODUCTION	12
1.1 Community interactions.....	12
1.2 Ecological theory and mid-domain effects	14
1.3 Hoki.....	16
1.4 Study area – Chatham Rise	16
1.5 Aims and outline of this study	17
1.6 LITERATURE CITED	19
2 The diet and diet variability of hoki (<i>Macruronus novaezelandiae</i>) on the Chatham Rise, New Zealand.	25
2.1 ABSTRACT.....	25
2.2 INTRODUCTION	26
2.3 MATERIALS AND METHODS.....	31
2.3.1 Sampling	31
2.3.2 Laboratory analysis	32
2.3.3 Hoki distribution	32
2.3.4 Overall diet.....	33
2.3.5 Diet variability	35
2.4 RESULTS	38
2.4.1 Hoki distribution	38
2.4.2 Overall Diet.....	41
2.4.3 Diet variability	45
2.4.3.1 Size based diet variation	45
2.4.3.1.1 Size based diet variation within depth ranges	48
2.4.3.2 Depth based diet variation	52
2.4.3.2.1 Depth based diet variation within size classes	54
2.4.3.3 Area based diet variation	58
2.4.3.3.1 Area based diet variation within set depth ranges and size classes	62
2.4.3.4 Sex based diet variation	66
2.4.4 Overview of hoki diet variability	66
2.5 DISCUSSION	67

2.5.1	Summary	73
3	The Mid-Domain Effect and Body-Size Depth Distribution in a Demersal Marine Fish Assemblage in New Zealand.	99
3.1	ABSTRACT	99
3.2	INTRODUCTION	100
3.3	MATERIALS AND METHODS.....	103
3.3.1	Study Area	103
3.3.2	Sampling	105
3.3.3	Mid-Domain Effect	106
3.3.4	Body-size depth relationship.....	107
3.4	RESULTS	107
3.4.1	Mid-domain effect	107
3.4.2	Body-size depth relationship.....	109
3.5	DISCUSSION	113
3.6	LITERATURE CITED	118
4	GENERAL DISCUSSION	127
4.1	Hoki diet.....	128
4.2	Body-size depth relationship.....	129
4.3	Mid-domain effect	131
4.4	Conclusions.....	133
4.5	Future directions	134
4.6	LITERATURE CITED	136

Table of figures

Figure 2.1: New Zealand with 1000 m isobath. Hoki spawning locations A = Hokitika canyon, and B = Cook Strait canyon (Top panel). The Chatham Rise from Banks Peninsula on the left to the Chatham Islands on the right (Bottom panel).	28
Figure 2.2: Number of samples collected in each area by size class and depth range.....	34
Figure 2.3: Relationship between hoki length (total length in cm) and weight (g) for all fish sampled, showing a near cubic relationship between length and weight ($y = 0.039x^{2.9357}$).....	39
Figure 2.4: Hoki length (cm) plotted against depth, for all fish sampled showing a significant relationship of body size increasing over depth $R^2 = .03724$ ($p < 0.001$).....	39
Figure 2.5: Hoki length plotted against sex (Male = 1, Female = 2), median (thick line), 25% and 75% quartiles (box) and 95% intervals (whiskers). The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.	40
Figure 2.6: Hoki sex (Male = 1, Female = 2) plotted against depth, median (thick line), 25% and 75% quartiles (box) and 95% intervals (whiskers). The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.	40
Figure 2.7: Proportions of stomachs in each digestive state plotted against the time of day that the fish was caught. The bars are drawn with widths proportional to the square-roots of the number of observations in the groups.	42
Figure 2.8: Proportions of stomach fullness plotted against time of day that fish were caught. The bars are drawn with widths proportional to the square-roots of the number of observations in the groups.....	42
Figure 2.9: The total number of prey categories (prey count) found in each stomach against hoki length. The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.	43

Figure 2.10: Prey accumulation curve for all 77 prey taxon identified, estimated from 999 random repeats, standard deviation plotted for every 20 th record. Horizontal lines show final five percent of total no of prey taxon identified.	44
Figure 2.11: Prey accumulation curve for the 11 prey categories used for multivariate analysis, estimated from 999 random repeats, standard deviation plotted for every 20 th record.....	44
Figure 2.12: Three dimensional MDS ordination of samples, categorised by size class.....	47
Figure 2.13: Similarity percentage contributions given by the prey groups that contributed for 90% of the diet similarity for each size group.	47
Figure 2.14: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each size group within the 200-400m depth range.	50
Figure 2.15: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each size group within the 400-600m depth range.	50
Figure 2.16: Three dimensional MDS ordination of samples, categorised by depth range.....	53
Figure 2.17: Similarity percentage contributions given by the prey groups that contribute to 90% of diet similarity for each depth range.	53
Figure 2.18: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 2+ size class.	56
Figure 2.19: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 3+ size class.	56
Figure 2.20: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 4+ size class.	57

Figure 2.21: Three dimensional MDS ordination of samples, categorised by area. Areas are North West (NW), South West (SW), North East (NE), and South East (SE).	60
Figure 2.22: Similarity percentage contributions given by the prey groups that contributed to 90% of diet similarity for each area.....	60
Figure 2.23: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each area group within the 400-600m depth range and 2+ size class.	64
Figure 2.24: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each area group within the 600-800m depth range and 4+ size class.	64
Figure 3.1: New Zealand with 1000 m depth contour, in relation to the Chatham Rise, dark rectangle (Top). The Chatham Rise from Banks Peninsula on the left to the Chatham Islands on the right (Bottom).	105
Figure 3.2: Output of the RangeModel showing the depth ranges of the 30 study species displayed as horizontal lines centred around their depth midpoints.	108
Figure 3.3: (A) Observed species richness (circles) over the depth domain, plotted with the mean expected species richness (solid line), as predicted by Colwell's Range Model, and the upper and lower 95% confidence intervals (broken lines). (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel. Line of best fit fitted through the data. $R^2 = 0.7732$ ($p < 0.001$).	108
Figure 3.4: (A) Observed species richness (circles) over the depth domain, plotted with mean expected species richness (solid line) as predicted by the RangeModel, and the upper and lower 95% confidence intervals (broken lines) for species with ranges set at one standard deviation of the observed depth ranges. (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel for species with depth ranges set at one standard deviation of observed depth ranges. $R^2 = 0.6667$ ($p < 0.001$).	110
Figure 3.5: (A) Observed species richness (circles) over the depth domain, plotted with mean expected species richness (solid line) as predicted by the	

RangeModel, and the upper and lower 95% confidence intervals (broken lines) for species with ranges set at two standard deviations of the observed depth ranges. (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel for species with depth ranges set at two standard deviations of observed depth ranges. $R^2 = 0.5949$ ($p < 0.001$). 110

Figure 3.6: Interspecific relationship of average body size and average depth caught at for (A) all 30 species studied, (B) Osteichthyes only, and (C) Chondrichthyes only. 111

Figure 3.7: Intraspecific relationship of body size and depth caught at for each of the 23 species studied. For ease of interpretation the regression lines are shown over the entire length range, but were only fitted over the length and depth of the actual species data. No regression line was plotted for non-significant results. Species codes are given in APPENDIX 3.1. 112

Table of tables

Table 2.1: Number of samples from each size class, depth range and area.....	34
Table 2.2: Summary table of the 11 groups prey was categorised into – full species list for each category given in APPENDIX 2.4.....	35
Table 2.3: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups.	48
Table 2.4: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups within 200-400m depth range.....	51
Table 2.5: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups within 400-600m depth range.....	51
Table 2.6: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups.....	54
Table 2.7: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 2+ size class.	57
Table 2.8: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 3+ size class.	58
Table 2.9: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 4+ size class.	58
Table 2.10: Pairwise test for significance showing R values and significance levels for the four areas compared.	59
Table 2.11: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups.	61
Table 2.12: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups within 400-600m depth range and 2+ size class.....	65
Table 2.13: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups within 600-800m depth range and 4+ size class.....	66
Table 3.1: Number of species, total and split by Class, and the type of body-size depth relationship shown. Significance at 0.05 level.....	111

Table of appendices

APPENDIX 2.1: Temperature maps.....	83
APPENDIX 2.2: Current maps.	85
APPENDIX 2.3: Stomach removal protocol.	87
APPENDIX 2.4: Categorised prey groupings.	89
APPENDIX 2.5: All prey %IRI table.	92
APPENDIX 2.6: Categorised data %IRI table.	95
APPENDIX 2.7: Prey species codes.....	96
APPENDIX 3.1: Species codes.	126

1 GENERAL INTRODUCTION

Understanding marine fish species interactions is important for both ecology and fisheries management. Commercial fishing is the removal of fish species in large amounts from the marine environment (Annala et al. 2004). Fishing generally targets the larger individuals of a population and given the amount/volume of removal is a strong top-down force that is artificial to the structure of the naturally functioning ecosystem (Pauly et al. 1998). Fisheries management sets limits on the amount of fish believed to be able to be removed sustainably (Ministry of Fisheries 2008). This estimation is usually based on an assessment of the target species with little or no regard to the potential flow-on effects of the target species removal (Ministry of Fisheries 2008). This disregard is often due to a lack of knowledge on the function of the ecosystem and the species interactions within it. Understanding the ecosystem is not only beneficial from a science perspective, but it can also have a useful purpose for fisheries management. Diet studies to determine prey consumption by predators is crucial for the development of the information base to model and understand food web interactions (Pimm et al. 1991). Information on diet composition is, however, not sufficient to quantify the functional relationships between predators and prey. Changes in ocean climate can cause spatial and temporal changes in predator and prey distributions that in turn lead to changes in trophodynamics (Marasco et al. 2007). A combination of food web investigation, the understanding of predator and prey distribution and research into environmental conditions are needed to understand the functioning of an ecosystem, and in turn lead to better information for fisheries management decisions.

1.1 Community interactions

One component of community organisation is “who-eats-whom”. The transfer of food energy from its source in plants through herbivores to carnivores is referred to as the food chain. It has long been recognised that these food chains are not isolated units, but linked together to form food webs (Krebs 2001). Food webs can form a useful starting point for theoretical analysis of community organisation (Pimm et al. 1991). In marine systems analysis of stomach contents has been shown to be among the most efficient and effective

General Introduction

means of understanding trophic interactions affecting ecosystem-scale function (Link 1999) and is an important step towards understanding wider trophic relationships (Pinnegar et al. 2003). Quantitative data (e.g. percent by weight for each prey item in the predator diet) are superior to qualitative data (presence-absence in diet) for generating estimates of carbon and/or energy flows through food webs, the fundamental basis of trophic modelling.

Morphology of predators and prey change with ontogeny as predators gape size increases and morphological defences of prey become more robust. The relationship between predator and prey sizes has been recognised as important in the likelihood of interspecific interactions. Often with fishes, the size of prey able to be consumed generally increases with increasing predator size (Juanes and Conover 1994, Keast and Webb 1966, Nielsen 1980, Persson et al. 1996, Scharf et al. 2000). Also the range of prey sizes eaten often increased with increasing predator size, with maximum prey size increasing while minimum prey size changes very little (Scharf et al. 2000). This prey size increase with predator size increase is seen in Garicía-Berthou and Moreno-Amich's (2000) study on pumpkinseed sunfish where not only does prey size increase but the diet also shifts with ontogeny. In some species there is a gradual change in diet with ontogeny, with larger fish able to eat larger prey and therefore a greater range of prey is available to them. In other species, like the pumpkinseed sun fish (Garicía-Berthou and Moreno-Amich 2000) and tuna (Graham et al. 2007) an abrupt change in diet is found once fish reach a certain size. One of the goals of this work was to look for such changes in hoki diet.

Along with size, diet can also change with time and space. Prey species are not distributed evenly over the environment and therefore predators in different places are forced to eat different things in the absence of other prey. Montgomery et al. (1993) found evidence of this in Antarctica with fish of the same species caught in different locations having very different diet composition. Seasonality may also affect prey distribution with environmental conditions being favourable to prey growth at certain times while at other times water temperature or food availability may be outside the prey species tolerance limits. Dempson et al. (2002) found evidence of diet changes with

different times of year. Prey species populations may also fluctuate on an annual (or longer) basis (i.e. sardine and anchovy population fluctuations (Lluch-Belda et al. 1992)) and predator species may have to adapt their diet to take advantage of the abundant prey species.

In New Zealand, feeding of exploited fish species has seldom been considered in evaluating their population dynamics in relation to fisheries management. In general, virtually nothing is known about trophic interactions between exploited fish species and other organisms in the ecosystem, or the factors determining predator-prey and competitive interactions (Annala et al. 2004).

1.2 Ecological theory and mid-domain effects

Two contrasting paradigms exist in community ecology, that of niche theory and neutrality.

Niche theory looks at how the differences between species allow them to co-exist (Whitfield 2002). There are two types of niche: the fundamental niche, the range of environmental conditions within which a species can live indefinitely in the absence of negative interactions (i.e. competition, predation, parasitism); and the realised niche, the environmental conditions/resources that the species actually exists in/uses in the presence of negative interactions (Krebs 2001). The niche concept implicitly assumes that neighbouring species have negative impacts (or niche shrinking affect) on one another and is firmly bound to the notion of competitive exclusion (that no two species can occupy the same niche) (Krebs 2001). Niche theory considers communities to be structured/organised based on species assemblages of those species which can best utilise the available resources (Nybakken 2001).

The second paradigm is neutral theory, which was developed by Hubbell (1979, 2001, 2003, 2005). The theory essentially considers the ecological properties of every individual in the population to be identical. Individuals compete for the same, fully

General Introduction

exploited pool of resources, but the identity of the winner of any single contest is left to chance (Whitfield 2002). Hubbell (2001) has shown that neutral theory is capable of summarizing a surprising array of empirical patterns of species abundances. Neutral theory is considered by some ecologists as a radical shift from established niche theories (Chave 2004).

Niche and neutral theories do not need to be conflicting, but can rather be viewed together as complementary (Leibold and McPeck 2006). Theories of coexistence by niche differentiation are mostly concerned with deterministic processes, and a small number of species that interact through fixed rules, as prescribed in the Lotka-Volterra equations. By contrast, the neutral theory is primarily concerned with species-rich communities (tropical forests, coral reefs) with many rare species, where the role of stochasticity at an individual scale becomes unavoidable (Chave 2004).

The cause or driving mechanism for the species richness gradients observed in the world is still unanswered. One hypothesis for species richness gradients is predicted through the mid-domain effect. The mid-domain effect was developed by Colwell and Hurtt (1994) and Willig and Lyons (1998). This model or effect is the resulting species richness pattern produced when latitudinal species ranges are shuffled randomly within the boundaries of a biogeographic domain, with the requirement that species ranges do not extend beyond the outer limits of the domain. The pattern of species richness produced is a parabolic curve with the peak of species richness in the centre of the domain, and declining towards the edges. Pure mid-domain effect models attempt to model species richness gradients without the consideration of external factors such as environmental or evolutionary events, and as a result have been considered a null model by some (Colwell et al. 2004, Kendall and Haedrich 2006). However, there are arguments about its suitability as a null model as it does not meet all the requirements of neutrality (Hawkins and Diniz-Filho 2002, Zapata et al. 2005), as it is not truly independent of external factors. A major goal of this work was to test for the mid-domain effect on the depth distribution of 30 abundant demersal fish species.

1.3 Hoki

Hoki (*Macruronus novaezelandiae*) is New Zealand's most important commercial fishery, worth \$693 million in 2007 (Ministry of Fisheries 2008). When the fishery started in 1970s and 80s the fish stock was thought to be able to sustain an annual total allowable commercial catch (TACC) of 250,000 tonnes (Ministry of Fisheries 2008). Since 2000 the TACC has steadily decreased to its present 90,000 tonnes. This decrease is in reaction to a falling biomass. The cause for this biomass drop is not totally understood, there is speculation that it may be due to heavy fishing pressure, a result of poor recruitment or a combination of both (Ministry of Fisheries 2007).

Hoki is distributed widely around New Zealand from depths of 10m to 900m, with greatest abundance between 200-600m (Annala et al. 2004). They are moderately – highly fecund with a female greater than 90cm total length releasing more than 1 million eggs in one spawning season. Hoki spawn during the winter months on two main spawning grounds. Over the first year planktonic eggs and larvae are moved inshore by advection and upwelling (Murdoch et al. 1990), and juveniles are distributed through out most coastal waters. The Chatham Rise is a nursery ground for 2-4 year old fish, with larger / older fish migrating from the Chatham Rise to the Subantarctic as they reach maturity (Livingston et al. 2002). Previous diet studies on hoki suggest a pelagic feeding strategy. These studies found hoki to feed on pelagic myctophid fish, euphausiids, prawns and amphipods (Bulman and Blaber 1986, Clark 1985b, Kerstan and Sahrhage 1980, Kuo and Tanaka 1984). Clark's (1985a) temporal and spatial analysis found hoki in different places and at different times to feed on different prey.

1.4 Study area – Chatham Rise

The Chatham Rise is a prominent bathymetric ridge that projects about 500 nautical miles (926 km) east from Banks Peninsula, on the east coast of the South Island, New Zealand, to around the Chatham Islands. It has a depth varying from about 250 m down to about 3000 m. Pacific deep water is found at the depths below about 2000 m, and has a temperature of 1–2°C with no appreciable seasonal variation. Above this, the Chatham

General Introduction

Rise is an area where the subtropical convergence forms, as warmer and more saline subtropical water from the north meets subantarctic water from the south (Sutton 2001). At 300m depth the average temperature on the Northern Rise is 10-11°C and at 800m it is 7-8°C, while the average temperature on the Southern Rise at 300m is 8-9°C and 4-6°C at 800m (Rickard 2008b). The subtropical convergence is the meeting point of warm pacific water from the North and cold Antarctic water from the south. It coincides with the location of the Chatham Rise, however it oscillates over the year being further south during the austral summer and further north during winter (Butler et al. 1992). Currents on the Chatham Rise move from west to east, they come down the eastern coast of the North Island from the north, and up the eastern coast of the South Island from the south, meet at the Rise and move east, oscillating north and south as it does so on either side of the Rise (Rickard 2008a).

Primary productivity is relatively high on the Chatham Rise compared with other offshore areas around New Zealand (Bradford-Grieve et al. 1999, Bradford-Grieve et al. 1997, Murphy et al. 2001). Zooplankton productivity may also be relatively high (McClatchie et al. 2004) and mesopelagic fish biomass has been reported as relatively high (McClatchie and Dunford 2003). This high productivity is widely believed to be the underlying reason why the Chatham Rise supports major fisheries such as hoki (Annala et al. 2004). When compared to the Subantarctic, the Chatham Rise has higher species diversity and richness (McClatchie et al. 1997).

1.5 Aims and outline of this study

The purpose of the first part of this study was to investigate the effects of several sources of variation on the feeding ecology of hoki (*Macruronus novaezelandiae*). The specific objectives were to investigate the distribution of hoki on the Chatham Rise by size, depth, and sex; characterise the diet of hoki on the Chatham Rise, New Zealand, between 200-800m; and test for the effects of fish size, depth, location and sex on diet.

General Introduction

The second part of this study tested for evidence of the mid-domain effect on 30 of the abundant species associated with hoki on the Chatham Rise. It also investigated whether overall, larger bodied fish were found in deeper water, whether there was a difference between Osteichthyes (bony fish, e.g. barracouta) and Chondrichthyes (cartilaginous fish, e.g. sharks) in body-size depth trends, and on an individual species basis, to characterise these trends, if present.

This thesis is structured into four chapters, beginning with this introduction. Chapter Two presents the results of the diet characterisation for hoki and the investigation of diet variability with fish size, depth and area. Chapter Three looks at a 30 species fish assemblage and explains their depth distribution over the Chatham Rise by way of the mid-domain effect, as well as investigating the individual species body-size depth relationships. The general conclusions of this study are presented in Chapter Four. Chapters Two and Three are written in the format of scientific papers each with separate abstracts, results, discussions and references. Because of this style there is some repetition in the introduction and methods sections.

1.6 LITERATURE CITED

Annala, J.H., Sullivan, K.J., Smith, N.W.M., Griffiths, M.H., Todd, P.R., Mace, P.M., and Connell, A.M. 2004. Report from the Fisheries Assessment Plenary, May 2004: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 690.

Bradford-Grieve, J., Boyd, P., Chang, F., Chiswell, S., Hadfield, M., Hall, J., James, M., Nodder, S., and Shushkina, E. 1999. Pelagic ecosystem structure and functioning in the subtropical front region east of New Zealand in austral winter and spring 1993. *Journal of Plankton Research* **21**(3): 405-428.

Bradford-Grieve, J.M., Chang, F.H., Gall, M., Pickmere, S., and Richards, F. 1997. Size-fractionated phytoplankton standing stocks and primary production during austral winter and spring 1993 in the Subtropical Convergence region near New Zealand. *New Zealand Journal of Marine and Freshwater Research* **31**: 201-224.

Bulman, C.M., and Blaber, S.J.M. 1986. Feeding ecology of *Macruronus novaezelandiae* (Hector) (Teleostei: Merlucciidae) in south-east Australia. *Australian Journal of Marine and Freshwater Research* **37**(5): 621-639.

Butler, E.C.V., Butt, J.A., Lindstrom, E.J., Teldesley, P.C., Pickmere, S., and Vincent, W.F. 1992. Oceanography of the subtropical convergence zone around southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* **26**: 131-154.

Chave, J. 2004. Neutral theory and community ecology. *Ecology Letters* **7**(3): 241-253.

Clark, M. 1985a. Feeding relationships of seven fish species from the Campbell Plateau, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **19**: 365-374.

Clark, M. 1985b. The food and feeding of seven fish species from the Campbell Plateau, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **19**: 339-363.

General Introduction – Literature Cited

- Colwell, R.K., and Hurtt, G.C. 1994. Nonbiological Gradients in Species Richness and a Spurious Rapoport Effect. *The American Naturalist* **144**(4): 570-595.
- Colwell, R.K., Rahbek, C., and Gotelli, N.J. 2004. The mid-domain effect and species richness patterns: what have we learned so far. *The American Naturalist* **163**(3).
- Dempson, J.B., Shears, M., and Bloom, M. 2002. Spatial and temporal variability in the diet of anadromous Arctic charr, *Salvelinus alpinus*, in northern Labrador. *Environmental Biology of Fishes* **64**(1): 49-62.
- Garicía-Berthou, E., and Moreno-Amich, R. 2000. Food of introduced pumpkinseed sunfish: ontogenetic diet shift and seasonal variation. *Journal of Fish Biology* **57**(1): 29-40.
- Graham, B.S., Grubbs, D., Holland, K., and Popp, B.N. 2007. A rapid ontogenetic shift in the diet of juvenile yellowfin tuna from Hawaii. *Marine Biology* **150**(4): 647-658.
- Hawkins, B.A., and Diniz-Filho, J.A.F. 2002. The mid-domain effect cannot explain the diversity gradient of Nearctic birds. *Global Ecology & Biogeography* **11**(5): 419-426.
- Hubbell, S.P. 1979. Tree Dispersion, Abundance, and Diversity in a Tropical Dry Forest. *Science* **203**(4387): 1299-1309.
- Hubbell, S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press.
- Hubbell, S.P. 2003. Modes of speciation and the lifespans of species under neutrality: a response to the comment of Robert E. Ricklefs. *Oikos* **100**(1): 193-199.
- Hubbell, S.P. 2005. Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional Ecology* **19**(1): 166-172.
- Juanes, F., and Conover, D.O. 1994. Piscivory and prey size selection in young-of-the-year bluefish: predator preference or size-dependent capture success? *Marine Ecology Progress Series* **114**: 59-59.

General Introduction – Literature Cited

- Keast, A., and Webb, D. 1966. Mouth and body form relative to feeding ecology in the fish fauna of a small lake, Lake Opinicon, Ontario. *Journal of the Fisheries Research Board of Canada* **23**: 1845 - 1874.
- Kendall, V.J., and Haedrich, R.L. 2006. Species richness in Atlantic deep-sea fishes assessed in terms of the mid-domain effect and Rapoport's rule. *Deep Sea Research Part I: Oceanographic Research Papers* **53**(3): 506-515.
- Kerstan, M., and Sahrhage, D. 1980. Biological investigations on fish stocks in the waters off New Zealand. Bundesforschungsanstalt für Fischerei, Hamburg. *Mitteilungen aus dem Institut für Seefischerei*: 29.
- Krebs, C.J. 2001. *Ecology: the Experimental Analysis of Distribution and Abundance*. Benjamin Cummings, San Francisco.
- Kuo, C., and Tanaka, S. 1984. Feeding habit of hoki *Macruronus novaezelandiae* (Hector) in waters around New Zealand. *Bulletin of the Japanese Society of Scientific Fisheries/Nissuishi* **50**(5): 783-786.
- Leibold, M.A., and McPeck, M.A. 2006. Coexistence of the niche and neutral perspectives in community ecology. *Ecology* **87**(6): 1399-1410.
- Link, J. 1999. Reconstructing food webs and managing fisheries. *In* *Proceedings of the 16th Lowell Wakefield Fisheries Symposium*. pp. 99-01.
- Livingston, M.E., Bull, B., Stevens, D.W., and Bagley, N.W. 2002. A review of hoki and middle depth trawl surveys of the Chatham Rise, January 1992-2001. NIWA Technical Report **113**.
- Lluch-Belda, D., Schwartzlose, R.A., Serra, R., Parrish, R., Kawasaki, T., Hedgecock, D., and Crawford, R.J.M. 1992. Sardine and anchovy regime fluctuations of abundance in four regions of the world oceans: a workshop report. *Fisheries Oceanography* **1**(4): 339-347.

General Introduction – Literature Cited

- Marasco, R.J., Goodman, D., Grimes, C.B., Lawson, P.W., Punt, A.E., and Quinn, T.J., II. 2007. Ecosystem-based fisheries management: some practical suggestions. *Canadian Journal of Fisheries and Aquatic Sciences* **64**(6): 928-912).
- McClatchie, S., and Dunford, A. 2003. Estimated biomass of vertically migrating mesopelagic fish off New Zealand. *Deep Sea Research Part I: Oceanographic Research Papers* **50**(10-11): 1263-1281.
- McClatchie, S., Macaulay, G.J., and Coombs, R.F. 2004. Acoustic backscatter and copepod secondary production across the Subtropical Front to the east of New Zealand. *Journal of Geophysical Research* **109**.
- McClatchie, S., Millar, R.B., Webster, F., Lester, P.J., Hurst, R., and Bagley, N. 1997. Demersal fish community diversity off New Zealand: Is it related to depth, latitude and regional surface phytoplankton? *Deep Sea Research Part I: Oceanographic Research Papers* **44**(4): 647-667.
- Ministry of Fisheries. 2007. Report from the Fisheries Assessment Plenary, May 2007: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 344-370.
- Ministry of Fisheries. 2008. Report from the Fisheries Assessment Plenary, May 2008: stock assessments and yield estimates. 347-371.
- Montgomery, J.C., Foster, B.A., Milton, R.C., and Carr, E. 1993. Spatial and temporal variations in the diet of nototheniid fish in McMurdo Sound, Antarctica. *Polar Biology* **13**(6): 429-431.
- Murdoch, R.C., Guo, R., and McCrone, A. 1990. Distribution of hoki (*Macruronus novaezelandiae*) eggs and larvae in relation to hydrography in eastern Cook Strait, September 1987. *New Zealand Journal of Marine and Freshwater Research* **24**(4): 529-539.
- Murphy, R.J., Pinkerton, M.H., Richardson, K.M., Bradford-Grieve, J.M., and Boyd, P.W. 2001. Phytoplankton distributions around New Zealand derived from SeaWiFS

General Introduction – Literature Cited

remotely-sensed ocean colour data. New Zealand Journal of Marine and Freshwater Research **35**: 343-362.

Nielsen, L.A. 1980. Effect of walleye (*Stizostedion vitreum vitreum*) predation on juvenile mortality and recruitment of yellow perch (*Perca flavescens*) in Oneida Lake, New York. Canadian Journal of Fisheries and Aquatic Sciences **37**(1): 11-19.

Nybakken, J.W. 2001. Marine biology : an ecological approach. Benjamin Cummings, San Francisco.

Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., and Torres Jr, F. 1998. Fishing down marine food webs. Science **279**(5352): 860.

Persson, L., Andersson, J., Wahlstrom, E., and Eklov, P. 1996. Size-specific interactions in lake systems: predator gape limitation and prey growth rate and mortality. Ecology: 900-911.

Pimm, S.L., Lawton, J.H., and Cohen, J.E. 1991. Food web patterns and their consequences. Nature **350**(6320): 669-674.

Pinnegar, J.K., Trenkel, V.M., Tidd, A.N., Dawson, W.A., and buit, M.H.D. 2003. Does diet in Celtic Sea fishes reflect prey availability? Journal of Fish Biology **63**(s1): 197-212.

Rickard, G. 2008a. Current maps of the Chatham Rise at depths. NIWA. p. 2.

Rickard, G. 2008b. Temperature maps of Chatham Rise at depths. NIWA. p. 2.

Scharf, F.S., Juanes, F., and Rountree, R.A. 2000. Predator size-prey size relationships of marine fish predators: interspecific variation and effects of ontogeny and body size on trophic-niche breadth. Marine Ecology Progress Series **208**: 229-248.

Sutton, P. 2001. Detailed structure of the Subtropical Front over Chatham Rise, east of New Zealand. Journal of Geophysical Research. C. Oceans **106**: 31045-31056.

Whitfield, J. 2002. Neutrality versus the niche. Nature **417**(6888): 480-481.

General Introduction – Literature Cited

Willig, M.R., and Lyons, S.K. 1998. An Analytical Model of Latitudinal Gradients of Species Richness with an Empirical Test for Marsupials and Bats in the New World. *Oikos* **81**(1): 93-98.

Zapata, F.A., Gaston, K.J., and Chown, S.L. 2005. The mid-domain effect revisited. *The American Naturalist* **166**: 144-148.

2 The diet and diet variability of hoki (*Macruronus novaezelandiae*) on the Chatham Rise, New Zealand.

2.1 ABSTRACT

The Chatham Rise, New Zealand, is a highly productive ecosystem, supporting commercial fisheries on several species, one of which is hoki (*Macruronus novaezelandiae*). Fisheries management in New Zealand is mainly on a single species basis. In order to shift towards a multispecies or ecosystem based management strategy, as is the global trend, ecosystem function, organisation and species interactions, such as predator prey relationships or competition must be understood. Diet studies are an effective method of investigating marine food webs. Hoki were collected from known locations over the Chatham Rise on three sampling occasions over the period 2005 to 2007 and their stomachs dissected to quantify diet. Hoki diet was found to consist largely of mesopelagic teleosts, mainly of the family Myctophidae, natant decapods and euphausiids, suggesting a pelagic feeding strategy. Differences were found in diet composition between this study and other studies on hoki diet, potentially suggesting differences in prey distribution between study areas. Differences in diet were found between fish from different depths and between different sized fish. The difference in diet between depths was independent of changes in size. The difference in diet between different sized fish was also still found independent of changes in depth. The maintenance of the independent size and depth diet differences indicated that there were true differences in diet with these variables, rather than one aliasing for the other or the differences in diet found being due to hoki size distribution over depth. The distribution of hoki was not homogeneous over the study area, with small (35-50cm) fish found mainly in the western part of the study area in shallower water, while large fish were predominately found at deeper depths over the whole study area. No consistent pattern of diet differences was found between the different areas within this study, suggesting that the patterns found may be aliasing depth and size patterns as well as reflecting differences in hoki distribution.

2.2 INTRODUCTION

One useful way to investigate the ecological role of marine animals and their place within food webs is through dietary studies (Cortés 1997, Pauly et al. 1998, Scharf et al. 2000). Both inter and intraspecific interactions affect ecosystem dynamics, and the population dynamics of the species involved.

Fisheries management has a vested interest in understanding population dynamics of target species. Stock management decisions are made based on the best understanding of the population dynamics of target species. As the global understanding of marine ecosystem function grows, management is moving away from the often failed single stock / species management approach (Francis et al. 2007) towards an ecosystem based management structure (Marasco et al. 2007). In order to successfully manage on an ecosystem scale, the ecosystem being managed must be understood. Ecosystem conservation is also growing in popularity, and in order to be able to conserve things one needs to have some form of baseline knowledge. Not only do we need to know what is there, and what used to be there, but also be able to monitor the effects of conservation efforts to see whether they are effective or not. A better understanding of ecosystem processes should also lead to a better understanding of target species population dynamics.

Natural systems vary with space and time, they fluctuate around a long-term equilibrium (Paine 1988). With this natural variation, diet will vary over space and time meaning that the interactions between predators and their prey will also change. Both predator size and location will affect the predator – prey relationship, as well as the further trophic interactions within the ecosystem. Changes in time and space may change predator – prey interactions. Ontogenetic changes in depth and area distributions may contribute to some changes in predator – prey relationships with adult fish often feeding on completely different prey from juvenile fish (Montgomery et al. 1993). Other studies have reported marked seasonal, regional and ontogenetic differences in diets (Braccini et al. 2005, Kuo

Hoki Diet – Introduction

and Tanaka 1984b). Historical studies have characterised diet and begun to make inferences about feeding ecology but have, in some cases, lacked sufficient data to make clear conclusions or to determine diet variability. This study attempts to quantitatively look at the intraspecific feeding variability of hoki.

Hoki (*Macruronus novaezelandiae*) is the most valuable commercial fish species in New Zealand, based on commercial quota values for 2007 (Ministry of Fisheries 2008). This fish stock has recently experienced a period of stock decline, so now is a pertinent time to look at hoki diet and try and further understand its population dynamics and place in the Chatham Rise ecosystem (Livingston et al. 2002, Ministry of Fisheries 2008). The reason for this decline is poorly understood, as we still do not understand what drives recruitment (Bull and Livingston 2001, Francis et al. 2006, Livingston 2000). One can speculate that diet and productivity may have changed or that this stock decline may be driven by poor recruitment, and / or heavy fishing mortality.

Hoki are found all around New Zealand, with the majority of commercial catch being taken from around the South Island. Hoki range in depth from 10-900m and have been found in highest abundance between 200-600m (Anderson et al. 1998, Annala et al. 2004). Females reach a larger maximum size than males (130cm total length (TL) for females and 115cm TL for males) (Annala et al. 2004). Maturity is reached at 3-5 years, maximum age is about 25 years (Horn 2007, Horn and Sullivan 1996, Ministry of Fisheries 2008). In the late 1980s and 1990s, hoki were very abundant, considered able to sustain an annual total allowable commercial catch (TACC) of between 200-250,000 tonnes. The TACC has decreased since 2000/01 from 250,000 tonnes to the current 90,000 tonnes for 2008/09. The drop in TACC is a reflection of the decrease in hoki biomass and a move towards catch rates believed to be sustainable over the long-term. Of the 90,000 TACC in the 2006/07 fishing year (1 October 2006 – 30 September 2007), 38,000 tonnes were caught on the Chatham Rise (Ministry of Fisheries 2008). Despite the recent drop in abundance, hoki are still a dominant benthic-pelagic predator on the Chatham Rise (Livingston et al. 2002, Stevens and O'Driscoll 2007) and are likely to play an important part in structuring that ecosystem.

The Chatham Rise (Figure 2.1) is a prominent bathymetric ridge that projects about 500 nautical miles (n. miles) (926 km) east from Banks Peninsula, on the east coast of the South Island, to around the Chatham Islands. It has a depth varying from about 250 m down to about 3000 m. Pacific deep water is found at the depths below about 2000 m, and has a temperature of 1–2° C with no appreciable seasonal variation.

Figure 2.1: New Zealand with 1000 m isobath. Hoki spawning locations A = Hokitika canyon, and B = Cook Strait canyon (Top panel). The Chatham Rise from Banks Peninsula on the left to the Chatham Islands on the right (Bottom panel).

Above this, the Chatham Rise is an area where the subtropical convergence forms, as warmer and more saline subtropical water from the north converge with subantarctic water from the south (Sutton 2001). The surface water temperatures on the Chatham Rise recorded during trawl surveys have consequently been about 2–3° C warmer on the northern side, and vary seasonally by about 3–4° C. The bottom water temperatures at

depths greater than 700 m, recorded during trawl surveys have been much less variable, being warmer on the northern side, but by $<1^{\circ}\text{C}$, and varying seasonally by $<1^{\circ}\text{C}$. APPENDIX 2.1 shows plots of temperatures on the Chatham Rise at various depths (Rickard 2008b). At 300m depth the average temperature on the Northern Rise is $10\text{--}11^{\circ}\text{C}$ and at 800m it is $7\text{--}8^{\circ}\text{C}$, while the average temperature on the Southern Rise at 300m is $8\text{--}9^{\circ}\text{C}$ and $4\text{--}6^{\circ}\text{C}$ at 800m. The subtropical convergence runs from about 45°S on the west coast of the South Island, around the bottom of the South Island, up the east coast, then extends (to a variable extent) through the Mernoo Gap at the western end of the Chatham Rise, before turning southwards and running eastwards along the Chatham Rise. Currents on the Chatham Rise move from west to east, they come down the eastern coast of the North Island from the north, and up the eastern coast of the South Island from the south, meet at the Rise and move east, oscillating north and south as it does so on either side of the Rise. APPENDIX 2.2 shows currents in detail at various depths (Rickard 2008a).

Primary productivity is relatively high on the Chatham Rise compared with other offshore areas around New Zealand (Bradford-Grieve et al. 1999, Bradford-Grieve et al. 1997, Murphy et al. 2001). Zooplankton productivity may also be relatively high (McClatchie et al. 2004) and mesopelagic fish biomass has also been reported as relatively high (McClatchie and Dunford 2003). This high productivity is widely believed to be the underlying reason why the Chatham Rise supports major fisheries such as hoki (Annala et al. 2004). When compared to the Subantarctic, the Chatham Rise has higher species diversity and richness (McClatchie et al. 1997). Roberston et al. (1978) found evidence of location specificity by species of mesopelagic fauna to the north and south of the Chatham Rise, giving a different species assemblage on the north and south. In the area directly over the rise there was lower species diversity, suggesting a relative paucity of species adapted to the mixed hydrological conditions found there. Hot spots of mesopelagic fish are suggested to exist over two banks at the western end of the rise (McClatchie et al. 2005). Benthic invertebrate biomass on the northern side of the Chatham Rise displays a logarithmic decline with water depth, comparable to other bathyal regions, but biomass estimates from the south side show no such relationship and

Hoki Diet – Introduction

are higher at depth than would be expected (Nodder et al. 2003, Probert and McKnight 1993). Benthic invertebrate patterns are important to not when investigating fish diet, as some diet variation can be a reflection of invertebrate prey distribution (Montgomery et al. 1993).

Hoki migrate over the course of their life. The main hoki spawning grounds are on the west coast of the South Island, in the Hokitika Canyon, and in the Cook Strait Canyon, northern South Island. The planktonic eggs and larvae are moved inshore by advection or upwelling (Murdoch et al. 1990) and are dispersed widely to the north and south. The resulting 0+ and one year old fish can be found in most coastal area of the South Island, and parts of the North Island. The major nursery ground for juvenile hoki (2-4 years) is on the Chatham Rise, in depths between 200 and 600m (Annala et al. 2004). The older fish disperse to deeper water, and are distributed widely through the Sub-Antarctic, Southern Plateau and Chatham Rise. From analysis of trawl survey (1991-2002) data, it is suggested that a significant proportion of hoki move off the Chatham Rise to the Sub-Antarctic as they reach maturity. Most movement was found to occur between 3 and 7 years (Livingston et al. 2002).

Several studies have been previously conducted on the feeding of hoki around New Zealand. Kerstan and Sahrhage (1980) gave some general notes on the feeding of hoki. Kuo and Tanaka (1984b) provided qualitative information on diet composition by area, feeding periodicity, ontogenetic diet change, and seasonality. Clark (1985) looked more quantitatively at the diet of hoki on the Campbell Plateau using the Index of Relative Importance (IRI) to describe diet, however this study was limited to species lists and graphs for comparison, lacking a true quantitative approach to investigate seasonal feeding and diet changes with ontogeny.

A quantitative approach is needed in order to determine the true variability of hoki diet with fish size, depth and location, as well as being to determine the trophic status of hoki and likely role in the Chatham Rise ecosystem. The purpose of this study is to investigate the effects of several sources of variation in the feeding ecology of hoki. The specific

objectives are to investigate the distribution of hoki on the Chatham Rise by size, depth, and sex; characterise the diet of hoki on the Chatham Rise, New Zealand, between 200-800m; and test for the effects of fish size, depth, location and sex on diet variability of hoki.

2.3 MATERIALS AND METHODS

2.3.1 Sampling

Stomach sampling took place on Ministry of Fisheries funded annual trawl surveys, that have been ongoing since 1992, carried out by the National Institute of Water and Atmospheric Research Ltd (NIWA) for hoki and middle depth species on the Chatham Rise, New Zealand (Figure 2.1). Survey details and results can be found in Stevens and O'Driscoll (2006, 2007). Briefly, sampling was conducted for roughly four weeks from late December to late January each year. Samples for this study were collected from the surveys fishing in January 2005, 2006, and 2007. Fish were captured in bottom trawls, the net was an 8-seam hoki bottom trawl with 100m sweeps, 50m bridles, 12m backstops, 58.8m ground rope, 45m headline and 60mm cod end mesh (see Hurst and Bagley (1994) for net plan and rigging details).

The sample design was a 2-phase stratified random, after Francis (1984). The same 26 strata were consistently used. Phase-1 trawl stations were allocated from simulations based on catch rates from the previous three trawl surveys, after Bull et al. (2000). At each station, the trawl was towed for approximately 3 nautical miles at a speed over ground of 3.5 knots (Hurst et al. 1992). The catch was sorted by species and either the whole catch or a representative sub sample of the fishes were weighed, to the nearest gram (on electronic, calibrated, motion compensated balances), and measured to the nearest millimetre. Fish length, weight and sex were recorded with the latitudinal and longitudinal location of each trawl station.

2.3.2 Laboratory analysis

Sub samples of fishes were randomly selected for analysis of stomach contents – full sampling protocol is given in APPENDIX 2.3 and Dunn (2005, 2006, 2007). The stomachs of the sampled fish were removed at sea, frozen and returned to land (full stomach removal protocol is given in APPENDIX 2.3). Once on land the stomachs were defrosted, weighed full and empty to get the total content weight, and prey (stomach content) was identified to the lowest taxonomic level possible, usually to species or genus, using existing guides and guides developed as part of the project from reference material collected on the trawl surveys. The number of individuals, weight, and digestive state of the prey was recorded in an electronic PostgreSQL database.

Stomach fullness, number of prey, prey weight and digestive state were recorded for all samples (stomachs) processed. Stomach fullness and digestive state were recorded in a number of different categories. Stomach fullness was recorded on a four level scale, 0 = empty, 1 = trace – empty to $\frac{1}{4}$ full, 2 = part full – $\frac{1}{4}$ to $\frac{3}{4}$ full, and 3 = full – over $\frac{3}{4}$ full. Digestive state was measured on a five level scale 1 = Fresh – Specimen entire and as ingested (may be broken from chewing), 2 = Slightly digested – Outer/exposed tissues starting to digest, otherwise specimen appears fresh, 3 = Partially digested – Outer/exposed tissues breaking down, still mostly complete with some soft tissue attached, 4 = Largely digested – Soft tissues mostly digested, some hardparts (vertebrae, exoskeleton etc.) still joined together, and 5 = Digested – Soup of digested hardparts (eyeballs, bones, beaks, etc.) with no soft tissue remaining.

2.3.3 Hoki distribution

Hoki distribution and characteristics over the Chatham Rise and were investigated using plots of fish length against weight, fish size against depth and sex, and fish sex against depth. A linear regression analysis was used to test for significance of the relationship between body size and depth. The effect of sex (male or female) upon body length and depth distribution were tested separately using independent sample t-tests with unequal variance.

2.3.4 Overall diet

Stomach fullness and prey digestive state were plotted against time to test whether evidence of feeding periodicity was present. The number of prey categories present per stomach was plotted against fish size and sex to look for evidence of the number of prey types consumed increasing with ontogeny. This provided a basic description of the data set and is important for understanding the potential relationships between proposed predictors of diet variability (e.g. fish size) associated with the data.

To assess whether the number of samples analysed was adequate, the cumulative number of new prey types against the cumulative number of stomachs sampled were plotted (Castriota et al. 2005, Ferry and Caillet 1996). If the resulting curve flattened out or reached an asymptote, as no further increases in prey species are found with additional stomachs investigated, then diet can be considered described. The PRIMER v6 software (Clarke and Gorley 2006) was used to create a prey species accumulation plot based on an average from 999 curves created by random reordering of the stomach samples. There is no definitive method for defining the asymptote of the curve. One definition of when the asymptote is reached is when the goodness of fit coefficient R^2 for the logistic regression is larger than the R^2 for the linear regression for the same curve (Campo et al. 2006). Another method, used by Braccini et al. (2005), was to consider the asymptote reached when at least the two previous values to the total sample diversity were in the range of asymptotic diversity ± 0.05 (Koen Alonso et al. 2002). Both methods were used in this study to test for adequacy of sampling.

No one basic statistic can completely describe fish diet (Hyslop 1980), therefore a combination of statistics were used. The Index of Relative Importance (IRI) is a statistic often used to evaluate the importance of prey items in diet. In this study, the percent weight (%W), percent number (%N), and percent frequency of occurrence (%O), were used to estimate the % IRI (Cortés 1997, Pinkas et al. 1971). Bootstrap methods (999 repeats) were used to estimate confidence intervals (95% or 2.5th% and 97.5th%) around the dietary statistics (%W, %N, %O and %IRI) (Haddon 2001).

Hoki Diet – Materials & Methods

The number of stomachs collected are shown by size, depth and are in Table 2.1 and Figure 2.2.

Table 2.1: Number of samples from each size class, depth range and area.

Size class	1+	2+	3+	4+
	125	432	220	511
Depth range	200-400m	400-600m	600-800m	
	344	707	237	
Area	NW	NE	SW	SE
	323	338	325	302

NW	1+	2+	3+	4+	NE	1+	2+	3+	4+
200-400m	61	59	4	9	200-400m	3	15	7	11
400-600m	10	71	15	17	400-600m	1	82	41	111
600-800m	0	4	9	64	600-800m	0	1	5	61

SW	1+	2+	3+	4+	SE	1+	2+	3+	4+
200-400m	39	36	6	9	200-400m	5	33	20	27
400-600m	2	54	56	60	400-600m	4	74	44	65
600-800m	0	0	9	54	600-800m	0	3	4	23

Figure 2.2: Number of samples collected in each area by size class and depth range.

2.3.5 Diet variability

Prey were categorised into 11 groups based on morphological and phylogenetic similarities (CRU, EUP, FIS, LAN, LAN 1, MES 1, OTH, PRA, RAT, SQX, UNF) (Table 2.2). A full species list and grouping is given in APPENDIX 2.4.

Table 2.2: Summary table of the 11 groups prey was categorised into – full species list for each category given in APPENDIX 2.4.

Group code	Contents
CRU	Other crustaceans, eg Amphipods and Copepods, NOT crustacean contained in groups EUP or PRA.
EUP	Euphausiids.
FIS	Unknown teleosts and benthic teleost.
LAN	Small bodied Myctophidae and similar sized mid water teleosts.
LAN 1	Large bodied Myctophidae and similar sized mid water teleosts.
MES 1	Other mid water teleosts.
OTH	Salps, shell fragments.
PRA	All Prawns and Mysids.
RAT	All rat tails and Javelin fish.
SQX	All squid.
UNF	All unidentified prey.

Size- related diet variations were investigated by dividing the collected sample into four size classes, roughly equivalent to age classes (Horn and Sullivan 1996, Livingston and Stevens 2005, Livingston et al. 2004, Stevens and O'Driscoll 2006): 1+ class: 35.6–49.9cm, 2+ class: 50.0–64.9cm, 3+ class: 65.0–72.9cm, and 4+ class: 73.0–113.8cm. Depth related diet variations were investigated by dividing the sample into three depth classes: 200–400m, 400–600m, and 600–800m. Area related diet variations were also investigated by dividing the sample into four areas: North East (NE) (east of 179.5W and north of 43.5S), North West (NW) (west of 179.5W and north of 43.5S), South East (SE) (east of 179.5W and south of 43.5S), and South West (SW) (west of 179.5W and south of 43.5S). Analysis of similarity (ANOSIM) based on Bray-Curtis dissimilarities from fourth-root transformed prey weight, was used to test the *a priori* hypothesis that hoki diet would differ significantly between the different variables (Clarke and Warwick 2001). Non-metric multidimensional scaling (MDS) ordination was used to visually display the differences in hoki diet. Prey weight was standardised to sum to one for each

stomach to remove the influence of differing stomach size. The fourth-root transformation was used, to change the relative weight of prey categories, as it is an intermediately severe transformation (Clarke and Warwick 2001), which has been used on previous diet studies (Braccini et al. 2005).

Analysis of similarity (ANOSIM) implemented in PRIMER was used to test for differences between different variable groups (depth range, size class, area and sex). ANOSIM creates a value (R value) or measure of difference that is between zero and one. A large value, near unity is indicative of complete separation of groups, while a small value, close to zero, implies little or no separation (Clarke and Warwick 2001). A small R value will indicate that groups do not have exactly the same diet (the hypothesis $R = 0$ can be rejected) but the small R value also tells us that the diets are strongly overlapping (R is close to zero). Small R values can still be significant because just as in ordinary univariate statistics, when the number of replicates is large in the groups being compared, giving rise to a very large number of possible permutations, small differences can still be statistically different. Generally the larger the sample size is, the more likely the R value will be small (Clarke and Gorley 2006).

A multidimensional scaling (MDS) ordination was used in PRIMER to visually represent the differences between the samples. In the MDS ordination plots each point is plotted in relation to all other points, with the most dissimilar points being placed furthest from each other. These ordinations can be plotted in two or three dimensions. When taking these multidimensional ordinations and reducing them to two or three dimensions a certain degree of distortion between the points will occur this is termed stress. The principle of the MDS algorithm is to choose a configuration of points that minimises this degree of stress while maintaining the true ordination of points. The level of stress is an indication of how true the point placement in the diagram is to what is represented in the data set. If stress is less than 0.05 it indicated a very good representation of the data. If the stress is less than 0.1, in two dimensions, then this is an adequate representation of the data and there is no need to view the ordination in 3D. Should the stress be less than 0.2 in two dimensions, while still useful, it is better to look at a 3D ordination. If stress is

greater than 0.3, in two dimensions, it is considered an inadequate representation and 3D should be used (Clarke and Warwick 2001).

A multivariate multiple permutation test, SIMPER (Similarity Percentages, PRIMER) (Clarke and Warwick 2001) was used to determine which prey categories, within each variable group (e.g. depth range), were responsible for the differences that were found, in terms of Bray-Curtis similarity. The SIMPER routine calculates the average dissimilarity between all pairs of inter-group samples and then breaks this average down into the separate contributions of each species to the overall dissimilarity between the two groups in the pairwise test. Typically there are many pairs of samples contributing to the average dissimilarity and a useful measure of how consistently a species contributes to the average dissimilarity across all pairs is the standard deviation values. If the average dissimilarity is large and the standard deviation small (and thus the ratio of average dissimilarity/standard deviation of average dissimilarity is large) then the species producing this not only contributes much to the dissimilarity between the two groups but it does so consistently in the inter-group comparisons of all samples in the two groups; it would thus be a good discriminating species (Clarke and Warwick 2001). A ratio value of 1.3 is considered a good cut off point for discriminating species, if the value is greater than 1.3 then it is considered a large ratio value, and therefore a discriminating species (Kröger 2003).

The more abundant a species is within a group, the more it will contribute to the intra-group similarities. It can characterise that group if it is found at a consistent abundance throughout, so the standard deviation of its abundance of its contribution is low, and average dissimilarity/standard deviation of average dissimilarity ratio is high. This, however, says nothing about whether the species is a good discriminator of one group from another, as it may be typical of a number of groups (Clarke and Warwick 2001).

The effect of each of the dietary variables (e.g. depth range) given the effect of other variables was tested by holding one variable constant (e.g. depth range) and testing the other (e.g. size) using pairwise tests. Tests for the independence of area within specific

depth ranges and size classes were restricted to two combinations only, as tests were not performed where less than 20 samples were collected per stratum, as small sample sizes will not fully demonstrate the diet and so may be biased.

2.4 RESULTS

2.4.1 Hoki distribution

A length weight relationship plot for hoki, showed a near cubic relationship between increasing fish length and weight ($y = 0.0039x^{2.9357}$) (Figure 2.3), and was similar to the length at weight parameter used by Francis (2003) in hoki stock assessments for the Chatham Rise ($y = 0.00479x^{2.89}$).

A noticeable absence of fish smaller than about 50cm, deeper than 500m, was found when fish size was plotted against depth (Figure 2.4). Fish larger than 50cm were found through out the studied depth range. There was a distinct group of small fish (~35-50cm) separated from the main bulk of the sample. This group was visible in other length weight relationships, and likely corresponds to the newly recruited one year old year class (Stevens and O'Driscoll 2006, 2007). A significant relationship of increasing body size with increasing depth was found $R^2 = 0.3724$ ($p = < 0.001$).

Results from an independent samples t-test with unequal variance showed that females were significantly larger than males ($p < 0.001$). Male and female hoki plotted against length in a box plot (Figure 2.5) that showed that the median female fish length was larger than the 75% quartile of the male length and that maximum female length was larger than maximum male length. The median male length was about the same size as the 25% quartile for females (~60cm).

The median depth of females was found towards the 75% quartile of the depth of males, with the median male depth found at the 25% quartile of female depth, indicating that

Hoki Diet – Results

more females were found in deeper water than males (Figure 2.6). Results from an independent samples t-test with unequal variance showed a significant difference between the average depths of male and female fish ($p = < 0.001$).

Female hoki are on average larger and found deeper than male hoki.

Figure 2.3: Relationship between hoki length (total length in cm) and weight (g) for all fish sampled, showing a near cubic relationship between length and weight ($y = 0.039x^{2.9357}$).

Figure 2.4: Hoki length (cm) plotted against depth, for all fish sampled showing a significant relationship of body size increasing over depth $R^2 = .03724$ ($p = < 0.001$)

Hoki Diet – Results

Figure 2.5: Hoki length plotted against sex (Male = 1, Female = 2), median (thick line), 25% and 75% quartiles (box) and 95% intervals (whiskers). The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.

Figure 2.6: Hoki sex (Male = 1, Female = 2) plotted against depth, median (thick line), 25% and 75% quartiles (box) and 95% intervals (whiskers). The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.

2.4.2 Overall Diet

Of the 1288 stomachs examined, 2 were empty, and of the remaining stomachs 37% contained only a single prey type. The number of prey types ranged from one to nine, with the majority having four or less prey types.

The proportion of stomachs in each digestive state was plotted against time of day (Figure 2.7), to investigate feeding periodicity. An increasing and then decreasing trend of the proportion of stomachs containing fresh prey was found, which reached its maximum between 1200-1600h. Stomach fullness was also plotted against time of day to see whether it coincided with digestive state (Figure 2.8). A decreasing trend of the proportion of full stomachs was found, with the minimum at 1600-2000h. Part full stomach proportions displayed an increasing and then decreasing trend, with the highest proportion of part full stomachs found at 1600-2000h. While not coinciding exactly with digestive state, stomach fullness did coincide with the hypotheses that hoki return to the bottom from night time mid-water feeding at dawn (McClatchie and Dunford 2003).

The number of different prey categories found in each stomach was plotted against hoki length. Fish of all sizes were found to consume four or less types of prey at one time, while five or more types of prey were only found in fish greater than 50cm in length (Figure 2.9). Of the 1288 stomachs analysed only 30 had five or more types of prey present.

The stomachs contained 77 taxonomically different prey items: 36 fish, 23 crustaceans, 10 molluscs, and other items such as polychaetes, salps, foraminifera and two unidentified categories (APPENDIX 2.5). The prey accumulation curve for the uncategorised data (Figure 2.10), while not completely flattening out at 1000 stomachs, was nearing an asymptote, to increase the number of prey taxon by 4 items (or 5% of the total number) an additional 120 stomachs were required. According to the method employed by Campo et al. (2006) the sample size of 1288 stomachs was adequate as the cumulative prey curve (Figure 2.10) fitted better with a logistic model ($R^2 = 0.9745$, $p < 0.001$) than with a linear model ($R^2 = 0.8652$, $p < 0.001$). Using Braccini's (2005)

Hoki Diet – Results

method, the asymptote was reached after 22 stomachs, and at 91 stomachs using ± 0.01 , so a sample size of 1288 was easily large enough to describe the diet of hoki. For the data grouped into the 11 categories for multivariate analysis, the prey accumulation curve completely flattened off at about 250 stomachs (Figure 2.11).

Figure 2.7: Proportions of stomachs in each digestive state plotted against the time of day that the fish was caught. The bars are drawn with widths proportional to the square-roots of the number of observations in the groups.

Figure 2.8: Proportions of stomach fullness plotted against time of day that fish were caught. The bars are drawn with widths proportional to the square-roots of the number of observations in the groups.

Hoki Diet – Results

Figure 2.9: The total number of prey categories (prey count) found in each stomach against hoki length. The boxes are drawn with widths proportional to the square-roots of the number of observations in the groups.

Euphausiids (family Euphausiacea) were the dominant prey item, contributing to the highest values of percent Index of Relative Importance (%IRI) (43.46%), percent number (%N) (40.25%) and second highest value of percent occurrence (%O) (29.47%); they did not feature in the top five dominant prey items by percent weight (%W). Lantern fish (family Myctophidae) were the second most important prey by %IRI (26.89%), % N (14.67%), and %O (33.52%) but only the third most important by %W (7.94%). The pasiphaeid prawn, *Pasiphaea spp.*, was the third most important prey item by %IRI (10.32%), %N (9.88%) and %O (19.29%) but also didn't feature in the top five important prey items by weight. The full %IRI table is presented in APPENDIX 2.5, with a full list of all species names, common names and codes used presented in APPENDIX 2.7. No evidence of cannibalism was found from my diet analysis of hoki stomachs.

Hoki Diet – Results

Figure 2.10: Prey accumulation curve for all 77 prey taxa identified, estimated from 999 random repeats, standard deviation plotted for every 20th record. Horizontal lines show final five percent of total no of prey taxa identified.

Figure 2.11: Prey accumulation curve for the 11 prey categories used for multivariate analysis, estimated from 999 random repeats, standard deviation plotted for every 20th record.

2.4.3 Diet variability

Based on the 11 category classification used for the multivariate analysis, LAN (small bodied lantern fish) was the most important prey item by % IRI (38.43%) and % O (48.08%), the second most important item by % N (25.72%) and the third most important item by % W (15.24%). The second most dominant prey group was EUP (Euphausiids, family Euphausiacea) by % IRI (23.91%), it was the most dominant prey group by % N (40.25%), the third most dominant by % O (29.47%) and did not feature in the top five most dominant by % W. PRA (prawns, order Decapoda and mysids, order Mysidae) were the third most dominant prey group by % IRI (22.67%) and by % N (19.38%), it was the second most dominant prey group by % O (41.37%) and the fourth most important by % W (8.69%). Complete %IRI table for the categorised prey can be found APPENDIX 2.6.

2.4.3.1 Size based diet variation

Multidimensional scaling (MDS) (Figure 2.12) showed some separation between the four size classes, however there was also a large amount of overlap observed. From the analysis of similarity (ANOSIM) the Global R (or sample statistic) was 0.031 with a significance level of 0.1% ($p = 0.001$) from 999 random permutations, indicating that there was a significant difference between the size classes. The pairwise tests resulted in significant differences being found between group 1+ and group 4+ ($R = 0.097$, $p = 0.001$), group 1+ and group 3+ ($R = 0.066$, $p = 0.001$), and between group 2+ and group 4+ ($R = 0.053$, $p = 0.001$), adjacent size classes were non significant. Figure 2.13 shows the proportions of the prey groups that contributed to 90% of the within group similarity.

Table 2.3 shows the dissimilarity contributions and average abundances for each prey group for the pairwise test groups that were significantly different. Four main prey groups (LAN, PRA, EUP and FIS) made up the majority of the diet in the four size class groups. The differences between the size class groups were largely based on the differing proportions of the four prey groups and the differing proportions that they contributed to the diet of the particular size class group. There were some other prey groups (RAT, MES 1 and CRU) that contributed to diet composition and fluctuated largely, they had

Hoki Diet – Results

quite different abundances between the size class groups, however their overall contribution to diet was small.

Few LAN were found in the large (4+) size group, with more FIS and PRA found instead. No PRA or FIS were found in the smallest (1+) size group, only LAN and EUP were found.

Hoki Diet – Results

Figure 2.12: Three dimensional MDS ordination of samples, categorised by size class.

Figure 2.13: Similarity percentage contributions given by the prey groups that contributed for 90% of the diet similarity for each size group.

Hoki Diet – Results

Table 2.3: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups.

Groups 1+ & 3+ Average dissimilarity = 69.15					
Species	Group 1+ Av.Abund	Group 3+ Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.59	1.6	18.21	1.06	26.34
EUP	1.41	0.65	16.31	1.06	23.58
PRA	0.42	1.23	15.06	0.91	21.78
FIS	0.46	0.53	10.15	0.62	14.68
CRU	0.21	0.21	4.07	0.45	5.89

Groups 1+ & 4+ Average dissimilarity = 79.84					
Species	Group 1+ Av.Abund	Group 4+ Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.59	0.84	18.57	1.05	23.26
EUP	1.41	0.29	16.52	1.01	20.69
PRA	0.42	1.2	14.83	0.9	18.57
FIS	0.46	0.93	13.67	0.78	17.12
CRU	0.21	0.17	3.73	0.42	4.67
RAT	0	0.3	3.58	0.33	4.49
MES 1	0.03	0.24	3.09	0.3	3.87

Groups 2+ & 4+ Average dissimilarity = 75.84					
Species	Group 2+ Av.Abund	Group 4+ Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.53	0.84	17.7	1.05	23.34
PRA	0.88	1.2	15.7	0.98	20.7
FIS	0.68	0.93	14.46	0.83	19.07
EUP	0.8	0.29	10.12	0.74	13.34
CRU	0.23	0.17	3.96	0.44	5.22
RAT	0.01	0.3	3.61	0.33	4.76
MES 1	0.04	0.24	3.08	0.31	4.06

2.4.3.1.1 Size based diet variation within depth ranges

Very few cases were found where the diets of adjacent hoki size classes were significantly different from each other, this supports the findings from the whole of the Chatham Rise that there is a gradual change in diet with increasing fish size. Large fish, while eating similar things to small fish also consumed other fish and an increased weight of prawns. Smaller hoki tended to eat more euphausiids than larger fish. LAN and FIS had no clear increasing or decreasing trend with fish size, and were eaten by all sizes of fish. Size class effect was found at 200-400m and 400-600m, and showed a similar pattern in diet change with hoki size. This may indicate that a diet change with fish size was not aliasing for depth changes (fish length and depth being correlated).

200-400m

A significant difference was found between the 4+ size class and both 1+ ($R = 0.125$, $p = 0.001$) and 2+ ($R = 0.072$, $p = 0.001$) size classes. No difference was found between the other size classes. The SIMPER pairwise test (Table 2.4) between 2+ and 4+ found 2+ to have a higher average abundance of LAN and EUP, 4+ to have higher abundances in FIS, PRA and RAT. The pairwise test (Table 2.4) between 1+ and 4+ found 1+ to have higher average abundances in LAN and EUP, and higher abundances in 4+ of FIS, PRA and RAT. Figure 2.14 and Figure 2.15 show the contributions of species groups that accounted for 90% of intra group similarity. The 4+ group had comparatively more PRA and FIS than both 1+ and 2+ in the 200-400m depth range. There was a decrease in the percentage contribution that EUP made as fish size increased.

400-600m

The 4+ size class was found to be significantly different to both the 2+ ($R = 0.054$, $p = 0.001$) and 3+ ($R = 0.021$, $p = 0.028$) size classes. No other significant differences were found between the different size classes. Results from the 2+ and 4+ pairwise test (Table 2.5) found 2+ with higher average abundances of LAN and EUP and 4+ with higher average abundances of PRA, FIS and RAT. The 3+ and 4+ pairwise test (Table 2.5) resulted in LAN and EUP being in higher abundance in 3+ and 4+ having higher abundances of PRA FIS and RAT. In the 400-600m depth range, 4+ has considerably less LAN than both 2+ and 3+ these two groups on the other hand have less PRA than 4+, as well as a smaller proportion of FIS.

600-800m

No significant differences were found between the diets of hoki at different size classes.

Hoki Diet – Results

Figure 2.14: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each size group within the 200-400m depth range.

Figure 2.15: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each size group within the 400-600m depth range.

Hoki Diet – Results

Table 2.4: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups within 200-400m depth range.

Groups 2+ & 4+		Average dissimilarity = 72.06				
Species	Group 2+ Av.Abund	Group 4+ Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.38	1.34	16.95	1.07	23.53	23.53
EUP	1.3	0.5	15.01	0.97	20.83	44.35
FIS	0.73	0.96	14.34	0.84	19.89	64.25
PRA	0.62	1.01	12.68	0.92	17.59	81.84
CRU	0.32	0.16	4.45	0.51	6.17	88.02
RAT	0.02	0.29	3.19	0.34	4.42	92.44

Groups 1+ & 4+
Average dissimilarity = 73.26

Species	Group 4+ Av.Abund	Group 1+ Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.34	1.48	17.81	1.06	24.31	24.31
EUP	0.5	1.5	17.07	1.05	23.3	47.62
FIS	0.96	0.47	13.58	0.78	18.53	66.15
PRA	1.01	0.43	12.33	0.87	16.83	82.98
CRU	0.16	0.23	3.69	0.45	5.04	88.02
RAT	0.29	0	3.06	0.33	4.18	92.2

Table 2.5: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different size groups within 400-600m depth range.

Groups 2+ & 4+		Average dissimilarity = 73.66				
Species	Group 2+ Av.Abund	Group 4+ Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.62	0.91	17.83	1.06	24.21	24.21
PRA	1	1.38	16.47	1.05	22.36	46.57
FIS	0.65	0.82	13.17	0.8	17.88	64.45
EUP	0.55	0.36	7.9	0.7	10.73	75.18
CRU	0.19	0.24	4.09	0.45	5.55	80.72
RAT	0.01	0.35	4.03	0.36	5.47	86.2
LAN 1	0.17	0.06	2.52	0.28	3.42	89.61
UNF	0.07	0.15	2.11	0.3	2.86	92.47

Groups 3+ & 4+ Average dissimilarity = 72.21

Species	Group 4+ Av.Abund	Group 3+ Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	0.91	1.67	17.73	1.09	24.55	24.55
PRA	1.38	1.24	16.74	1.06	23.18	47.73
FIS	0.82	0.47	11.94	0.75	16.53	64.26
EUP	0.36	0.57	7.96	0.71	11.02	75.28
RAT	0.35	0.01	4.03	0.36	5.58	80.86
CRU	0.24	0.19	3.98	0.45	5.51	86.37
MES 1	0.13	0.1	2.56	0.28	3.55	89.93
OTH	0.14	0.07	2.15	0.27	2.98	92.91

2.4.3.2 Depth based diet variation

Some separation was shown in the MDS ordination (Figure 2.16) for the three depth ranges, however there was still a lot of overlap. From the analysis of similarity (ANOSIM) the Global R (or sample statistic) was 0.079 with a significance level of 0.1% ($p = 0.001$) from 999 random permutations, indicating that there was a significant difference between the depth ranges. The pairwise tests found significant (0.1% significance level) differences between all three depth ranges. Figure 2.17 shows the proportions of the prey groups that contributed to 90% of the within group similarity.

Table 2.6 shows the dissimilarity contributions and average abundances for each prey group for the pairwise test groups that were significantly different. Four main prey groups (LAN, PRA, EUP and FIS) made up the majority of the diet in the three depth range groups. The differences between the depth ranges was largely based on the differing proportions of the four prey groups and the differing proportions that they contributed to the diet of the particular depth range. There were some other prey groups (RAT, MES 1, SQX and CRU) that contribute to diet composition and fluctuate largely, but they had quite different abundances between the depth ranges, however their overall contribution to diet was small. EUP was found only in the shallow, 200-400m, depth. The presence of LAN was found to decrease with depth, while the presence of FIS and PRA increased with the increasing depth.

Hoki Diet – Results

Figure 2.16: Three dimensional MDS ordination of samples, categorised by depth range.

Figure 2.17: Similarity percentage contributions given by the prey groups that contribute to 90% of diet similarity for each depth range.

Hoki Diet – Results

Table 2.6: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups.

Groups 400-600m & 600-800m Average dissimilarity = 76.56					
Species	400-600m Av.Abund	600-800m Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.39	0.66	17.31	0.99	22.61
PRA	1.18	1.08	16.99	1	22.19
FIS	0.66	1.07	15.79	0.87	20.62
EUP	0.49	0.14	6.42	0.57	8.39
MES 1	0.09	0.38	5.18	0.4	6.76
RAT	0.13	0.22	3.94	0.34	5.15
CRU	0.2	0.11	3.24	0.37	4.23
SQX	0.09	0.17	2.67	0.33	3.49

Groups 200-400m & 600-800m Average dissimilarity = 80.21					
Species	200-400m Av.Abund	600-800m Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.46	0.66	17.62	1.03	21.97
FIS	0.64	1.07	15.66	0.86	19.52
EUP	1.24	0.14	14.85	0.9	18.51
PRA	0.67	1.08	14.77	0.9	18.42
MES 1	0.02	0.38	4.58	0.38	5.71
CRU	0.27	0.11	3.86	0.42	4.81
RAT	0.07	0.22	3.23	0.31	4.03

Groups 200-400m & 400-600m Average dissimilarity = 71.83					
Species	200-400m Av.Abund	400-600m Av.Abund	Av.Diss	Diss/SD	Contrib%
LAN	1.46	1.39	17.43	1.06	24.27
PRA	0.67	1.18	14.53	0.95	20.22
EUP	1.24	0.49	14.52	0.97	20.22
FIS	0.64	0.66	12.16	0.73	16.92
CRU	0.27	0.2	4.43	0.49	6.16
RAT	0.07	0.13	2.11	0.26	2.94

2.4.3.2.1 Depth based diet variation within size classes

EUP consistently decreased with the increase of depth across different size classes. In general, PRA increased with the increase of depth. FIS showed a weak trend of increasing with depth increases, however it is not consistent across all sizes classes. The group MES 1 featured only in the deepest depth and the largest size class. These findings are similar to the overall picture of depth (Figure 2.24, Table 2.6)

1+

No significant differences in the diet of hoki were found between the different depths.

2+

A significant difference was found between the 200-400m and 400-600m depth ranges ($R = 0.041$, $p = 0.003$). No other differences were found. The results from the pairwise SIMPER test (Table 2.7) between 200-400m and 400-600m found higher average abundances of EUP and FIS in 200-400m while 400-600m had higher average abundances of LAN and PRA, Figure 2.18 shows the contributions of species groups that account for 90% of intra group similarity

3+

The 600-800m depth range was significantly different to the other two depth ranges, 200-400m ($R = 0.218$, $p = 0.001$) and 400-600m ($R = 0.122$, $p = 0.009$). No difference was found between the other depth ranges. From the pairwise test (Table 2.8) between 400-600m and 600-800m it was found that 400-600m had higher average abundances in LAN and EUP, while 600-800m had higher average abundances of PRA, FIS and MES 1. The pairwise test (Table 2.8) between 200-400m and 600-800m resulted in LAN and EUP being found in higher average abundances in 200-400m, while PRA and FIS were found in higher average abundances in 600-800m. Figure 2.19 shows the contributions of species groups that account for 90% of intra group similarity.

4+

A significant difference was found between the 400-600m and 600-800m depth ranges ($R = 0.033$, $p = 0.001$), no other differences were found between the depth ranges. The pairwise test between 400-600m and 600-800m found higher average abundances of PRA, LAN, RAT and EUP in 400-600m, and 600-800m had higher average abundances of FIS and MES 1. Figure 2.20 shows the contributions of species groups that account for 90% of intra group similarity.

Hoki Diet – Results

Figure 2.18: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 2+ size class.

Figure 2.19: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 3+ size class.

Hoki Diet – Results

Figure 2.20: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each depth group within the 4+ size class.

Table 2.7: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 2+ size class.

Groups 200-400m & 400-600m			Average dissimilarity = 70.26			
Species	400-600m Av.Abund	200-400m Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.62	1.38	17.9	1.08	25.47	25.47
EUP	0.55	1.3	15.36	1	21.86	47.33
PRA	1	0.62	13.25	0.89	18.86	66.19
FIS	0.65	0.73	12.94	0.75	18.42	84.61
CRU	0.19	0.32	4.87	0.5	6.94	91.55

Hoki Diet – Results

Table 2.8: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 3+ size class.

Groups 400-600m & 600-800m			Average dissimilarity = 72.04			
Species	400-600m Av.Abund	600-800m Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.67	0.77	19.47	1.09	27.03	27.03
PRA	1.24	1.39	18.46	1.05	25.63	52.66
FIS	0.47	1.13	15.66	0.83	21.73	74.39
EUP	0.57	0.13	7.04	0.62	9.77	84.16
CRU	0.19	0.18	4.07	0.39	5.64	89.8
MES 1	0.1	0.11	2.28	0.26	3.17	92.97

Groups 200-400m & 600-800m			Average dissimilarity = 74.57			
Species	200-400m Av.Abund	600-800m Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	1.91	0.77	18.8	1.27	25.21	25.21
PRA	1.1	1.39	16.22	1.11	21.76	46.97
EUP	1.36	0.13	14.78	1.11	19.82	66.78
FIS	0.34	1.13	13.86	0.82	18.59	85.37
CRU	0.34	0.18	5.19	0.52	6.96	92.33

Table 2.9: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different depth groups 4+ size class.

Groups 400-600m & 600-800m			Average dissimilarity = 76.64			
Species	600-800m Av.Abund	400-600m Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
PRA	1.03	1.38	17.43	1.04	22.75	22.75
FIS	1.07	0.82	15.98	0.92	20.85	43.59
LAN	0.62	0.91	13.39	0.82	17.48	61.07
RAT	0.25	0.35	6.49	0.46	8.47	69.54
MES 1	0.44	0.13	6.09	0.45	7.95	77.49
EUP	0.13	0.36	4.9	0.49	6.39	83.88
CRU	0.09	0.24	3.36	0.38	4.39	88.27
SQX	0.19	0.13	3.18	0.37	4.15	92.42

2.4.3.3 Area based diet variation

Multidimensional scaling (MDS) of the four areas showed some separation (Figure 2.21), however there was also lot of overlap. From the analysis of similarity (ANOSIM) the Global R (or sample statistic) is 0.01 with a significance level of 0.1% ($p = 0.001$) from 999 random permutations, indicating that there was a significant difference found between the areas. The pairwise tests showed significant differences (Table 2.10) between all but one pairwise combination, North West and North East comparisons were non-significant. Figure 2.22 shows the proportions of the prey groups that contributed to 90% of the within group similarity. Table 2.11 shows the dissimilarity contributions and

Hoki Diet – Results

average abundances for each prey group for the pairwise test groups that were significantly different. Four main prey groups (LAN, PRA, EUP and FIS) made up the majority of the diet in the four area groups. The differences between the areas was largely based on the differing proportions that the four main prey groups contributed to the diet of the particular area. There were some other prey groups (RAT, MES 1, SQX and CRU) that contributed to diet composition and fluctuated largely, while they had quite different abundances between the depth ranges, however their overall contribution to diet was small.

Table 2.10: Pairwise test for significance showing R values and significance levels for the four areas compared.

Groups	R Statistic	Significance Level %
NW, SW	0.013	0.1
NW, NE	0.002	11.3
NW, SE	0.011	0.4
SW, NE	0.019	0.1
SW, SE	0.008	1.4
NE, SE	0.006	3.5

LAN was found in higher abundances in the south, while PRA and FIS were found in higher abundances in the north, no obvious trend of prey distribution was apparent though.

Hoki Diet – Results

Figure 2.21: Three dimensional MDS ordination of samples, categorised by area. Areas are North West (NW), South West (SW), North East (NE), and South East (SE).

Figure 2.22: Similarity percentage contributions given by the prey groups that contributed to 90% of diet similarity for each area.

Hoki Diet – Results

Table 2.11: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups.

Groups NW & SW Average dissimilarity = 73.73

Species	Group NW	Group SW	Av.Diss	Diss/SD	Contrib%
	Av.Abund	Av.Abund			
LAN	1.09	1.33	17.15	1.02	23.26
PRA	1.18	0.87	15.84	0.96	21.49
FIS	0.76	0.64	13.44	0.75	18.23
EUP	0.62	0.88	12.84	0.84	17.41
CRU	0.17	0.25	4.13	0.43	5.6
MES 1	0.13	0.14	2.89	0.3	3.92
RAT	0.16	0.06	2.5	0.26	3.38

Groups SW & NE Average dissimilarity = 74.10

Species	Group SW	Group NE	Av.Diss	Diss/SD	Contrib%
	Av.Abund	Av.Abund			
LAN	1.33	1.23	17.26	1.03	23.29
PRA	0.87	1.17	15.52	0.97	20.94
FIS	0.64	0.7	12.71	0.74	17.15
EUP	0.88	0.43	11.75	0.79	15.85
CRU	0.25	0.21	4.4	0.46	5.94
MES 1	0.14	0.14	3.12	0.3	4.22
RAT	0.06	0.17	2.47	0.28	3.33

Groups NW & SE Average dissimilarity = 72.85

Species	Group NW	Group SE	Av.Diss	Diss/SD	Contrib%
	Av.Abund	Av.Abund			
LAN	1.09	1.48	17.65	1.03	24.23
PRA	1.18	0.86	15.59	0.97	21.39
FIS	0.76	0.84	14.25	0.83	19.55
EUP	0.62	0.59	10.25	0.78	14.07
CRU	0.17	0.19	3.46	0.43	4.75
RAT	0.16	0.14	3.25	0.31	4.47
MES 1	0.13	0.09	2.3	0.27	3.16

Groups SW & SE Average dissimilarity = 72.23

Species	Group SW	Group SE	Av.Diss	Diss/SD	Contrib%
	Av.Abund	Av.Abund			
LAN	1.33	1.48	17.85	1.04	24.71
PRA	0.87	0.86	13.75	0.9	19.04
FIS	0.64	0.84	13.62	0.78	18.85
EUP	0.88	0.59	12.31	0.84	17.05
CRU	0.25	0.19	4.25	0.45	5.88
MES 1	0.14	0.09	2.53	0.27	3.5
RAT	0.06	0.14	2.17	0.26	3

Groups NE & SE Average dissimilarity = 72.64

Species	Group NE	Group SE	Av.Diss	Diss/SD	Contrib%
	Av.Abund	Av.Abund			
LAN	1.23	1.48	17.56	1.04	24.18
PRA	1.17	0.86	15.26	0.97	21
FIS	0.7	0.84	13.62	0.82	18.76
EUP	0.43	0.59	8.79	0.72	12.1
CRU	0.21	0.19	3.75	0.46	5.16
RAT	0.17	0.14	3.22	0.32	4.43
UNF	0.14	0.13	2.59	0.32	3.57
MES 1	0.14	0.09	2.55	0.27	3.51

2.4.3.3.1 Area based diet variation within set depth ranges and size classes

Results from the investigations between the different size classes and different areas reflect some of the findings from tests performed on all samples (Figure 2.13, Table 2.3, Figure 2.22 and Table 2.11). Like the earlier results using all the samples, no clear pattern was exhibited between the different areas.

400-600m

1+

Sample sizes were too small for robust analysis.

2+

North West was found to be significantly different from all the other areas, South West ($R = 0.153$, $p = 0.001$), South East ($R = 0.152$, $p = 0.001$) North East ($R = 0.159$, $p = 0.001$). No differences were found between any of the other areas. The SIMPER pairwise test (Table 2.12) between North West and North East found PRA and FIS to be in higher average abundance in stomachs in North West, with North East having higher average abundances in LAN and EUP in stomachs. North West was found to have higher average abundances of PRA and FIS when compared with South West, which has higher average abundances of LAN and EUP. The pairwise test (Table 2.12) between North West and South East found PRA and FIS in higher average abundance in North West, and South East to have higher average abundance of LAN and EUP. Figure 2.14 shows the contributions of species groups that accounted for 90% of intra group similarity.

3+

Sample sizes were too small for robust analysis.

4+

Sample sizes were too small for robust analysis.

600-800m

1+

No samples collected.

2+

Sample sizes were too small for robust analysis.

3+

Sample sizes were too small for robust analysis.

4+

South East was significantly different from North West ($R = 0.077$, $p = 0.017$) and North East ($R = 0.078$, $p = 0.016$). No differences were found between the other areas. The pairwise test (Table 2.13) between South East and North West found higher average abundances of FIS, PRA and RAT in North West and South East to have higher average abundances of LAN and MES 1. From the pairwise test (Table 2.13) between South East and North East it was found that North East had higher average abundances of FIS, PRA and RAT, while South East had higher average abundances of LAN and MES 1. Figure 2.24 shows the contributions of species groups that account for 90% of intra group similarity.

Hoki Diet – Results

Figure 2.23: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each area group within the 400-600m depth range and 2+ size class.

Figure 2.24: Similarity percentage contribution for the main prey types contributing to at least 90% of the similarity within each area group within the 600-800m depth range and 4+ size class.

Hoki Diet – Results

Table 2.12: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups within 400-600m depth range and 2+ size class.

Groups NW & SW		Average dissimilarity = 73.32				
Species	Group NW Av.Abund	Group SW Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
PRA	1.88	0.65	20.17	1.21	27.51	27.51
LAN	0.87	1.71	18.61	1.1	25.37	52.89
FIS	0.86	0.73	14.84	0.81	20.24	73.13
EUP	0.3	0.9	10.98	0.83	14.97	88.1
CRU	0.21	0.14	3.6	0.38	4.9	93

Groups NW & NE		Average dissimilarity = 73.59				
Species	Group NW Av.Abund	Group NE Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	0.87	1.89	19.57	1.15	26.59	26.59
PRA	1.88	0.75	19.24	1.21	26.14	52.73
FIS	0.86	0.4	12.61	0.74	17.14	69.87
EUP	0.3	0.57	7.54	0.73	10.25	80.12
LAN 1	0.13	0.41	5.51	0.44	7.49	87.61
CRU	0.21	0.27	4.67	0.47	6.35	93.96

Groups NW & SE		Average dissimilarity = 71.86				
Species	Group NW Av.Abund	Group SE Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
LAN	0.87	1.96	20.48	1.17	28.5	28.5
PRA	1.88	0.7	20.03	1.22	27.88	56.38
FIS	0.86	0.65	14.02	0.81	19.51	75.89
EUP	0.3	0.52	7.49	0.67	10.43	86.31
CRU	0.21	0.12	3.28	0.4	4.56	90.88

Hoki Diet – Results

Table 2.13: Pairwise similarity percentage (SIMPER) tests for dissimilarity between significantly different area groups within 600-800m depth range and 4+ size class.

Groups NW & SE		Average dissimilarity = 81.43				
Species	Group NW Av.Abund	Group SE Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
FIS	1.23	0.82	18.9	0.92	23.21	23.21
LAN	0.5	0.92	14.02	0.78	17.22	40.42
PRA	0.92	0.48	13.33	0.83	16.37	56.8
MES 1	0.33	0.65	9.95	0.6	12.22	69.02
RAT	0.55	0.14	8.41	0.5	10.33	79.35
SQX	0.17	0.27	5.39	0.4	6.62	85.97
CRU	0.04	0.29	3.66	0.36	4.49	90.46

Groups NE & SE		Average dissimilarity = 80.65				
Species	Group NE Av.Abund	Group SE Av.Abund	Av.Diss	Diss/SD	Contrib%	Cum.%
FIS	1.03	0.82	17.46	0.89	21.65	21.65
PRA	1.21	0.48	16.3	0.93	20.21	41.86
LAN	0.67	0.92	15.03	0.82	18.64	60.5
MES 1	0.4	0.65	10.77	0.62	13.36	73.86
SQX	0.26	0.27	6.48	0.45	8.03	81.89
CRU	0.04	0.29	3.68	0.35	4.57	86.46
RAT	0.15	0.14	3.67	0.31	4.56	91.01

2.4.3.4 Sex based diet variation

From the analysis of similarity (ANOSIM) the Global R (or sample statistic) is 0.007 with a significance level of 12.7% ($p = 0.127$) from 999 random permutations, indicating that there is no significant difference in diet found between the sexes.

2.4.4 Overview of hoki diet variability

Differences in diet were found between fish of different sizes, however adjacent size classes were not found to be different from each other, indicating a gradual change in diet with increasing size. Fish from different depths were also found to have different diets. This could be a result of smaller fish being found consistently in the shallower depths only, however, when only the shallow depth was investigated the co-occurring larger fish were eating different prey from the smaller fish. When only large fish were investigated a difference in diet was also found between the different depths. Differences were found between the different areas investigated, however no consistent pattern in prey difference was found. No significant difference in diet was found between the sexes. In all the

pairwise SIMPER tests performed, no one prey species was found that could be considered to be a discriminating species, nor were there any species that characterised any of the groups tested. Four prey species groups consistently ranked in the top four contributors to the dissimilarities between the variable groups tested. The relative contribution of each of the four prey groups was the main difference between the variable groups tested.

2.5 DISCUSSION

The diet of hoki from the Chatham Rise included a range of teleosts, predominately from the family Myctophidae, Sub-order Crustacea, which were largely euphausids and natant decapods, and Cephalopods (mainly squid). This result suggests a predominantly pelagic feeding strategy, as also found by Bulman and Blaber (1986), Clark (1985), Kerstan and Sahrhage (1980), and Kuo and Tanaka (1984b). The most important prey by percent Index of Relative Importance (% IRI) in the overall diet of hoki was euphausids. When examined by the 11 prey groupings, the group LAN was found to be most important, with euphausids (EUP) second most important. This difference is due to the way the data has been categorised. EUP is a single taxon grouping, where as LAN contains multiple taxa and species; for EUP to come out as second most important prey in the categorised data highlights its importance to the diet of hoki. LAN is made up of unidentified lantern fish (Family Myctophidae), which ranked second most important by %IRI in the uncategorised data, as well as the myctophid LHE (*Lampanyctodes hectoris*), which ranked fourth in the uncategorised data. Given that the category LAN (in the categorised data) contains both the second and fourth most important taxa in the diet of hoki it comes as little surprise that it ranks more important in the categorised data, than EUP.

Sampling effort was spread fairly evenly by area. By depth range, about twice as many fish were sampled in 400-600m as the shallower and deeper depth categories. When split by size class, the 4+ (largest) size class had the most samples, followed by the 2+ size class, then 3+, with the fewest samples coming from the 1+ (smallest) size class. Even

Hoki Diet – Discussion

though 1288 stomachs were studied, there were still limitations to the data set. When broken down by depth, size class, and area there are still some groups with zero or very few samples (Table 2.1 and Figure 2.2). Some of this is due to fish of each criteria not being found in the specific places, for example 1+ fish not being found in 600-800m depth, other reasons for this may be due to restricted sampling effort due to limits in time and funding. That being said, the samples collected are adequate to describe the overall diet of hoki and look at general patterns over broad categories.

Two methods (Braccini et al. 2005, Campo et al. 2006) were used to determine whether adequate number of samples had been collected and processed, based on species accumulation curves reaching the asymptote. Using Braccini et al. (2005) method, the hoki sample was considered adequate while the species accumulation curve was still steeply rising. This method was originally also used on species diversity which creates a curve that flattens earlier than an equivalent curve based on number of prey categories accumulation. While this study did not assess at what point the samples would be considered adequate using the method from Campo et al. (2006), the criteria seems rather easy to satisfy. The quantitative criteria used to determine whether a prey species or diversity accumulation curve was nearing an asymptote was arbitrary, and clearly differs between studies and authors.

Soft bodied prey, or prey that digests easily, may be under represented if remaining hard parts, (e.g. otoliths, scales, beaks and bones) are included in further analysis (MacDonald et al. 1982). Hard parts (e.g. otoliths and beaks) may also remain in the stomach for several days and feeding episodes and if used in further analysis bias the results towards prey that contain these hard parts. Soft and slippery prey may also be more easily regurgitated prior to stomach removal, while hard and or spiky prey may remain in the stomach. The amount of bias introduced is hard to determine and has not been investigated as part of this study.

Clark (1985) found natant decapods, mainly the undescribed caridean shrimp, *Pasiphaea* sp., to be the most important prey overall, followed by myctophid fishes (notably

Hoki Diet – Discussion

Lampanyctodes hectoris and *Gymnoscopelus piabilis*), followed by amphipods. While the order of important prey is not the same as this study, two of the top three important prey are the same, lantern fish and the prawn *Pasiphaea* sp. The big difference between the two studies is the dominance of amphipods found in the diet from the Campbell Plateau, which was not found in this study on the Chatham Rise, which may be due to patchy prey distribution but not enough is known about amphipod distribution to draw a clear conclusion.

The samples used in this study created a snapshot of the diet of hoki, studied at only one time during the year. This limited sampling time allowed for no investigation of the changes in diet with changes in season. There may be fluctuations in the abundance of prey, and therefore shifts in diet reflecting this variation. Clark (1985) found differences in the importance of prey species between different regions, both within and between seasons. Myctophids declined in importance and the myctophid *Lampanyctodes hectoris* was not recorded during spring in the vicinity of the Auckland Islands on the Campbell Plateau. Clark (1985) suggested this decrease in myctophids and absence of *L. hectoris* may have been due to changes in the position of the subtropical convergence which may not have extended as far south in spring 1979 as it did in autumn. Kuo and Tanaka (1984b) found evidence of increased feeding during autumn and postulated that this was in anticipation for the upcoming winter spawning season. Diet may also change between sampling years, however this was not investigated in my study. Prey identification improved over the course of the three years, with more prey being identified to a lower taxonomic level in the third year than in the earlier ones. Keeping the three years separate might therefore introduce some further bias into the analysis as the three years would not be truly comparable. Amalgamating the samples from the three years likely reduced this bias, and produces a larger sample for a more robust analysis.

Kerstan and Sahrhage (1980) found hoki mainly ate fish, squid, natant decapods and euphausiids. They also observed that hoki less than 60 cm preferred depths shallower than 500m, whereas larger and therefore older fish occurred most frequently deeper than 500m. This study found this to be true, however 50 cm was found to be a clearer cut off

Hoki Diet – Discussion

point in the data. Kerstan and Sahrhage (1980) found evidence, with the use of echo sounders, of patterns characteristic of diel vertical migration. No description of species was given for fish or natant decapods, nor were proportions or frequencies of occurrence reported.

Kuo and Tanaka (1984b) found the dominant food, by rate of occurrence, to be small shrimp, myctophids, euphausiids, synodontidae (lizard fish) and galatheids. They gave no location or area description but refer to three different areas, however one might assume that they are referring to the same areas described in Kuo and Tanaka (1984a). Using that assumption, their “East” area covers the Chatham Rise and is essentially comparable. Their findings of dominant prey are similar in some respects, in that small shrimps, myctophids and euphausiids feature highly, as they do in this study, but in a different order of importance. However the two studies findings differ largely in the findings of lizard fish and galatheids occurring frequently. No lizard fish were found in this study and only a single occurrence of a galatheid was found. Kuo and Tanaka (1984b) also found cannibalism occurring on the Chatham Rise, again this was not found to be the case in this study despite a sample 1288 fish. Kuo and Tanaka (1984b) found the occurrence of fish and myctophids decreased with increasing fish size, while the occurrence of small shrimps (no species given) increased with fish size increase. In this study it was also found that the importance of prawns increased with fish size and that the importance of euphausiids decreased with fish size.

Bulman and Blaber (1986) used bomb calorimetry to determine the energy values of major prey of hoki found around South Eastern Australia. While not directly comparable, as this study does not estimate or evaluate the energy intake or components of hoki diet, the prey composition in itself is interesting to note that there are large similarities in species composition. The biggest difference between the two studies is the evidence of cannibalism found by Bulman and Blaber (1986). Differential distribution of adults and juveniles is a mechanism that reduces the incidence of cannibalism in some inshore fishes (Blaber 1979) and may be a reason for the lack of cannibalism found in the present study. Clark (1985) also found no evidence of cannibalism in the diet of hoki from the Campbell

Hoki Diet – Discussion

Plateau, New Zealand, suggesting that either cannibalism doesn't occur or that there is good physical separation between the adults and juveniles which prevents it occurring. Hoki spawn on the west coast of the South Island, New Zealand, and the Cook Strait (Livingston 1990, Zeldis et al. 1998). Both of these areas are physically separated from the Chatham Rise, this study area, and the Campbell plateau, Clark's (1985) study area. By the time the juvenile fish have reached the Chatham Rise nursery ground they may have reached a sufficient size to avoid predation by adult hoki.

Kuo and Tanaka (1984b) found evidence of diel migration based on catch per unit effort (CPUE) where less fish were caught during the night (666-710kg/hr), 1800h – 0600h, than during the day (1293-3475kg/hr), 0600h – 1800h. McClatchie and Dunford (2003) recorded evidence of mesopelagic fish on the Chatham Rise vertically migrating, rising off the bottom at dusk and forming a distinct layer during the night in the epipelagic zone and then descending to join the bottom layer of fish at dawn. Bull (2000) found hoki rose off the bottom at evening and returned at dawn. This result would explain the differences in CPUE found by Kuo and Tanaka (1984b) and Langley et al. (2001), indicating a decrease in hoki on the bottom during the night. Samples for the present study were collected from trawling only during daylight hours (Hurst et al. 1992) in order to prevent a bias in catch rates caused by this difference between day and night. Tows that were done during the night were experimental and only for diet comparison, and not included in biomass estimates.

No clear evidence for diurnal feeding periodicity was found by Kuo and Tanaka (1984b), as was found in this study. They propose that this lack of clear evidence may be due to the sampling period with the most samples collected was between 0600h – 1400h. We assume here that feeding takes place on a daily basis, and that that the majority of fish leave the bottom to feed on the mid water layers, however we don't know the rate of gastric evacuation for hoki nor do we know whether all fish feed every day. We can however, conclude that all fish do not leave the bottom to feed every dusk, as catch rates of hoki at night were not zero. This may obscure the results of stomach fullness in this study.

McClatchie et al. (2005) suggests that there are hot spots of mesopelagic fish on the Mernoo and Veryan Banks at the western end of the Chatham Rise. These hot spots may also harbour a higher abundance of euphausiids and small mesopelagic fish due to the similarity in their diet and the presence of an abundant primary productivity food source (Dilling et al. 1998, Schmaltz 2008), which might contribute to the diet bias of small hoki, caught predominantly on the western end of the Chatham Rise, towards EUP and LAN. This source of food, coupled with comparatively shallow, warm water, which may enhance the growth rate of small fish, may also explain why small hoki, less than 50cm are predominantly only caught on the western end of the Chatham Rise.

Significant differences were found between the different sized of fish, different depths and different areas with the ANOSIM test. The associated R values (all less than 0.2) indicated these differences were small and a large amount of overlap was found in the diet. This result is displayed in the MDS ordinations, showing no clear separation between the individuals in each group. The groups that showed the largest difference in diet were depth, followed by size. This result may be due to prey distribution, with different prey being found in deeper water (eg. myctophid hot spots on Mernoo and Veryan banks (McClatchie and Dunford 2003)), however not enough is known about prey species distribution on the Chatham Rise to draw strong conclusions about this. Hoki distribution may also contribute somewhat to the differences found between different sizes and depths, with small fishes only being found in the shallowest depth range. However some large fish were found in the shallow water, also eating different prey to the small fish, giving weight to the hypotheses that differences are more due to prey distribution and patchiness. Gape size will also contribute to differences in diet, with small fish being physically incapable of consuming the prey variety of larger fish. The area differences found may alias the differences in size and depth, given that there was no clear trend in the area differences when holding the size and depth constant, and when compared to the overall area comparisons. It may rather be a reflection of fish distribution over the Chatham Rise, with predominantly small fish being found in the western end and larger fish being found to the east.

Optimal foraging theory predicts that larger fish will search for and consume single larger items of prey that provide more energy, rather than expend excessive energy chasing smaller prey items that are comparatively energy poor (Krebs 2001). This also suggests a reason why there is a change in diet as fish size increases. Bulman and Blaber (1986) calculated the energy components of different prey in hoki diet, and found the lantern fish *Lampanyctodes hectoris* to have the highest energy component of teleosts examined. *L. hectoris* is a large part of the composite group LAN used in the analysis, and may be why this group is consistently being consumed by fish of all size ranges. Bulman and Blaber (1986) also found the energy components for euphausiids and the prawn *Pasiphaea* sp. to be high. This may explain their dominance in the diet of hoki, Small fish may not be able to consume the larger bodied *Pasiphaea* sp. and therefore consume the smaller euphausiids in their absence. Then once the small fish have reached a large enough size, where they can physically consume the larger *Pasiphaea* sp., they may switch to or start consuming the more energy rich *Pasiphaea* sp., possibly explaining the increase in PRA in the larger fish. Euphausiids are the sole component of the group EUP, while *Pasiphaea* sp. make up a large proportion of the PRA species group used in the analysis. Alternatively, the depth distribution of *Pasiphaea* sp. is relatively unknown and they may in fact live in the deeper water, where the smaller fish are not, and therefore the increase in larger fishes diet may be due to a migration deeper rather than a change in size.

2.5.1 Summary

From the investigations I found hoki are not distributed homogeneously across the Chatham Rise. Different sized fish are found in different areas and at different depths. Diet differences were found between fish of different size and from different depths. Fish from different depths and of different sizes exploit different resources in the different areas, and probably have different competitors for prey at different sizes and places depending on the species composition of the fish assemblages they live in. This is important to note when looking at the ecosystem as a whole, rather than just at a species specific level. This study has only investigated the interactions of hoki on diet and does

Hoki Diet – Discussion

not consider the interactions that hoki may have with other fish species through diet on the Chatham Rise.

Further research is needed to understand the distribution and abundance patterns of the prey of hoki. From there we can begin to understand the impact these prey, and their abundances in time and space, may have on the diet variability of hoki. Investigation in to hoki diet over the course of a year is needed to test for seasonal variation. Comparing the prey composition in hoki diet with local prey abundances and fluctuations over time would give information on whether hoki is actively selecting certain for prey types or just eating the most abundant prey.

LITERATURE CITED

Anderson, O.F., Bagley, N.W., Hurst, R.J., Francis, M.P., Clark, M.R., and McMillan, P.J. 1998. Atlas of New Zealand fish and squid distributions from research bottom trawls. NIWA Technical Report **42**(Unpublished report held in NIWA library, Wellington).

Annala, J.H., Sullivan, K.J., Smith, N.W.M., Griffiths, M.H., Todd, P.R., Mace, P.M., and Connell, A.M. 2004. Report from the Fisheries Assessment Plenary, May 2004: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 690.

Blaber, S.J.M. 1979. The biology of filter feeding teleosts in Lake St Lucia, Zululand. *Journal of Fish Biology* **15**(1): 37-59.

Braccini, J.M., Gillanders, B.M., and Walker, T.I. 2005. Sources of variation in the feeding ecology of the piked spurdog (*Squalus megalops*): implications for inferring predator-prey interactions from overall dietary composition. *ICES Journal of Marine Science* **62**(6): 1076.

Bradford-Grieve, J., Boyd, P., Chang, F., Chiswell, S., Hadfield, M., Hall, J., James, M., Nodder, S., and Shushkina, E. 1999. Pelagic ecosystem structure and functioning in the subtropical front region east of New Zealand in austral winter and spring 1993. *Journal of Plankton Research* **21**(3): 405-428.

Bradford-Grieve, J.M., Chang, F.H., Gall, M., Pickmere, S., and Richards, F. 1997. Size-fractionated phytoplankton standing stocks and primary production during austral winter and spring 1993 in the Subtropical Convergence region near New Zealand. *New Zealand Journal of Marine and Freshwater Research* **31**: 201-224.

Bull, B. 2000. An acoustic study of the vertical distribution of hoki on the Chatham Rise. *New Zealand Fisheries Assessment Report* **5**.

Bull, B., Bagley, N.W., and Hurst, R.J. 2000. Proposed survey design for the Southern Plateau trawl survey of hoki, hake and ling in November-December 2000. Final Research

Hoki Diet – Literature Cited

Report to the Ministry of Fisheries for Project MDT199/01 Objective 1.(Unpublished report held by Ministry of Fisheries, Wellington): 31.

Bull, B., and Livingston, M.E. 2001. Links between climate variation and year class strength of New Zealand hoki(*Macruronus novaezelandiae*): an update. *New Zealand Journal of Marine and Freshwater Research* **35**(5): 871-880.

Bulman, C.M., and Blaber, S.J.M. 1986. Feeding ecology of *Macruronus novaezelandiae* (Hector) (Teleostei: Merlucciidae) in south-east Australia. *Australian Journal of Marine and Freshwater Research* **37**(5): 621-639.

Campo, D., Mostarda, E., Castriota, L., Scarabello, M.P., and Andaloro, F. 2006. Feeding habits of the Atlantic bonito, *Sarda sarda* (Bloch, 1793) in the southern Tyrrhenian sea. *Fisheries Research* **81**(2-3): 169-175.

Castriota, L., Scarabello, M.P., Finoia, M.G., Sinopoli, M., and Andaloro, F. 2005. Food and feeding habits of pearly razorfish, *Xyrichtys novacula* (Linnaeus, 1758), in the southern Tyrrhenian Sea: variation by sex and size. *Environmental Biology of Fishes* **72**(2): 123-133.

Clark, M. 1985. The food and feeding of seven fish species from the Campbell Plateau, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **19**: 339-363.

Clarke, K.R., and Gorley, R.N. 2006. *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth.

Clarke, K.R., and Warwick, R.M. 2001. *Changes in marine communities: an approach to statistical analysis and interpretation*, 2nd edition. PRIMER-E, Plymouth.

Cortés, E. 1997. A critical review of methods of studying fish feeding based on analysis of stomach contents: Application to elasmobranch fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **54**(3): 726-738.

Hoki Diet – Literature Cited

- Dilling, L., Wilson, J., Steinberg, D., and Alldredge, A. 1998. Feeding by the euphausiid *Euphausia pacifica* and the copepod *Calanus pacificus* on marine snow. *Marine Ecology Progress Series* **170**: 189-201.
- Dunn, M. 2005. Voyage Report (TAN0501) For Ministry of Fisheries project ZBD2004/02. Unpublished Report held at NIWA library, Wellington, NZ.
- Dunn, M. 2006. Voyage Report (TAN0601) For Ministry of Fisheries project ZBD2004/02. Unpublished Report held at NIWA library, Wellington, NZ.
- Dunn, M. 2007. Voyage Report (TAN0601) For Ministry of Fisheries project ZBD2004/02. Unpublished Report held at NIWA library, Wellington, NZ.
- Ferry, L.A., and Caillet, G.M. 1996. Sample size and data analysis: are we characterizing and comparing diet properly? *In* Feeding Ecology and Nutrition in Fish, Symposium Proceedings. *Edited by* D. MacKinlay and K. Shearer. American Fisheries Society, San Francisco. pp. 71-80.
- Francis, R., Hadfield, M.G., Bradford-Grieve, J.M., Renwick, J.A., and Sutton, P.J.H. 2006. Links between climate and recruitment of New Zealand hoki (*Macrurus novaezelandiae*) now unclear. *New Zealand Journal of Marine and Freshwater Research* **40**: 547-560.
- Francis, R.C., Hixon, M.A., Clarke, M.E., Murawski, S.A., and Ralston, S. 2007. Fisheries Management-Ten Commandments for Ecosystem-Based Fisheries Scientists. *Fisheries-American Fisheries Society* **32**(5): 217-233.
- Francis, R.I.C.C. 1984. An adaptive strategy for stratified random trawl surveys. *New Zealand Journal of Marine and Freshwater Research* **18**: 59-71.
- Francis, R.I.C.C. 2003. Analyses supporting the 2002 stock assessment of hoki. *New Zealand Fisheries Assessment Report* **5**.
- Haddon, M. 2001. *Modelling and Quantitative Methods in Fisheries*. Chapman & Hall.

Hoki Diet – Literature Cited

- Horn, P. 2007. Ageing a fish—how and why? *New Zealand Science Review* **63**: 58.
- Horn, P.L., and Sullivan, K.J. 1996. Validated aging methodology using otoliths, and growth parameters for hoki (*Macruronus novaezelandiae*) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research* **30**(2): 161-174.
- Hurst, R., Bagley, N.W., Chatterton, T., Hanchet, S., Schofield, K.A., and Vignaux, M. 1992. Standardisation of hoki/middle depth time series trawl surveys. MAF Fisheries Greta Point Internal Report **194**(Unpublished report held in NIWA library, Wellington): 89.
- Hurst, R.J., and Bagley, N.W. 1994. Trawl survey of middle depth and inshore bottom species off Southland, February-March 1993 (TAN9301). *New Zealand Fisheries Data Report* **52**: 58.
- Hyslop, E.J. 1980. Stomach contents analysis-a review of methods and their application. *Journal of Fish Biology* **17**(4): 411-429.
- Kerstan, M., and Sahrhage, D. 1980. Biological investigations on fish stocks in the waters off New Zealand. Bundesforschungsanstalt für Fischerei, Hamburg. *Mitteilungen aus dem Institut für Seefischerei*: 29.
- Koen Alonso, M., Alberto Crespo, E., Aníbal García, N., Noemí Pedraza, S., Ariel Mariotti, P., and Judith Mora, N. 2002. Fishery and Ontogenetic Driven Changes in the Diet of the Spiny Dogfish, *Squalus acanthias*, in Patagonian Waters, Argentina. *Environmental Biology of Fishes* **63**(2): 193-202.
- Krebs, C.J. 2001. *Ecology: The experimental analysis of distribution and abundance* - 5th Edition. Benjamin Cummings, an imprint of Addison Wesley Longman, Inc, San Francisco.
- Kröger, K. 2003. Recovery of subtidal benthic macroinvertebrate communities following natural and experimental disturbances. Unpublished PhD Thesis, Victoria University of Wellington.

Hoki Diet – Literature Cited

Kuo, C., and Tanaka, S. 1984a. Distribution and migration of hoki *Macruronus novaezelandiae* (Hector) in waters around New Zealand. Bulletin of the Japanese Society of Scientific Fisheries/Nissuishi **50**(3): 391-396.

Kuo, C., and Tanaka, S. 1984b. Feeding habit of hoki *Macruronus novaezelandiae* (Hector) in waters around New Zealand. Bulletin of the Japanese Society of Scientific Fisheries/Nissuishi **50**(5): 783-786.

Langley, A.D., Dunn, A., and Ballara, S.L. 2001. Determination of CPUE indices from catch effort data for hoki (*Macruronus novaezelandiae*) for the 1986-87 to 1998-99 fishing years. New Zealand Fisheries Assessment Report **49**: 70 p.

Livingston, M.E. 1990. Spawning hoki(*Macruronus novaezelandiae* Hector) concentrations in Cook Strait and off the east coast of the South Island, New Zealand, August-September 1987. New Zealand Journal of Marine and Freshwater Research **24**(4): 503-517.

Livingston, M.E. 2000. Links between climate variation and the year class strength of New Zealand hoki(*Macruronus novaezelandiae*) Hector. New Zealand Journal of Marine and Freshwater Research **34**(1): 55-69.

Livingston, M.E., Bull, B., Stevens, D.W., and Bagley, N.W. 2002. A review of hoki and middle depth trawl surveys of the Chatham Rise, January 1992-2001. NIWA Technical Report **113**.

Livingston, M.E., and Stevens, D.W. 2005. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2004 (TAN0401). New Zealand Fisheries Assessment Report **21**.

Livingston, M.E., Stevens, D.W., O'Driscoll, R.L., and Francis, R.I.C.C. 2004. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2003 (TAN0301). New Zealand Fisheries Assessment Report **16**.

MacDonald, J.S., Waiwood, K.G., and Green, R.H. 1982. Rates of digestion of different prey in Atlantic cod (*Gadus morhua*), ocean pout (*Macrozoarces americanus*), winter

Hoki Diet – Literature Cited

flounder (*Pseudopleuronectes americanus*), and American plaice (*Hippoglossoides platessoides*). Canadian Journal of Fisheries and Aquatic Science **39**(5): 651-659.

Marasco, R.J., Goodman, D., Grimes, C.B., Lawson, P.W., Punt, A.E., and Quinn, T.J., II. 2007. Ecosystem-based fisheries management: some practical suggestions. Canadian Journal of Fisheries and Aquatic Sciences **64**(6): 928(912).

McClatchie, S., and Dunford, A. 2003. Estimated biomass of vertically migrating mesopelagic fish off New Zealand. Deep Sea Research Part I: Oceanographic Research Papers **50**(10-11): 1263-1281.

McClatchie, S., Macaulay, G.J., and Coombs, R.F. 2004. Acoustic backscatter and copepod secondary production across the Subtropical Front to the east of New Zealand. Journal of Geophysical Research **109**.

McClatchie, S., Millar, R.B., Webster, F., Lester, P.J., Hurst, R., and Bagley, N. 1997. Demersal fish community diversity off New Zealand: Is it related to depth, latitude and regional surface phytoplankton? Deep Sea Research Part I: Oceanographic Research Papers **44**(4): 647-667.

McClatchie, S., Pinkerton, M., and Livingston, M.E. 2005. Relating the distribution of a semi-demersal fish, *Macruronus novaezelandiae*, to their pelagic food supply. Deep Sea Research Part I: Oceanographic Research Papers **52**(8): 1489-1501.

Ministry of Fisheries. 2008. Report from the Fisheries Assessment Plenary, May 2008: stock assessments and yield estimates. 347-371.

Montgomery, J.C., Foster, B.A., Milton, R.C., and Carr, E. 1993. Spatial and temporal variations in the diet of nototheniid fish in McMurdo Sound, Antarctica. Polar Biology **13**(6): 429-431.

Murdoch, R.C., Guo, R., and McCrone, A. 1990. Distribution of hoki (*Macruronus novaezelandiae*) eggs and larvae in relation to hydrography in eastern Cook Strait, September 1987. New Zealand Journal of Marine and Freshwater Research **24**(4): 529-539.

Hoki Diet – Literature Cited

- Murphy, R.J., Pinkerton, M.H., Richardson, K.M., Bradford-Grieve, J.M., and Boyd, P.W. 2001. Phytoplankton distributions around New Zealand derived from SeaWiFS remotely-sensed ocean colour data. *New Zealand Journal of Marine and Freshwater Research* **35**: 343-362.
- Nodder, S.D., Pilditch, C.A., Probert, P.K., and Hall, J.A. 2003. Variability in benthic biomass and activity beneath the Subtropical Front, Chatham Rise, SW Pacific Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* **50**(8): 959-985.
- Paine, R.T. 1988. Road Maps of Interactions or Grist for Theoretical Development? *Ecology* **69**(6): 1648-1654.
- Pauly, D., Trites, A.W., Capuli, E., and Christensen, V. 1998. Diet composition and trophic levels of marine mammals. *ICES Journal of Marine Science: Journal du Conseil* **55**(3): 467.
- Pinkas, L.M., Oliphant, M.S., and Iverson, I.L.K. 1971. Food habits of albacore, bluefin tuna and bonito in Californian waters. *California Fish and Game* **152**: 1-105.
- Probert, P.K., and McKnight, D.G. 1993. Biomass of bathyal macrobenthos in the region of the Subtropical Convergence, Chatham Rise, New Zealand. *Deep Sea Research Part I: Oceanographic Research Papers* **40**(5): 1003-1007.
- Rickard, G. 2008a. Current maps of the Chatham Rise at depths. NIWA. p. 2.
- Rickard, G. 2008b. Temperature maps of Chatham Rise at depths. NIWA. p. 2.
- Robertson, D.A., Roberts, P.E., and Wilson, J.B. 1978. Mesopelagic faunal transition across the Subtropical Convergence east of New Zealand. *New Zealand Journal of Marine and Freshwater Research* **12**: 295-312.
- Scharf, F.S., Juanes, F., and Rountree, R.A. 2000. Predator size-prey size relationships of marine fish predators: interspecific variation and effects of ontogeny and body size on trophic-niche breadth. *Marine Ecology Progress Series* **208**: 229-248.

Hoki Diet – Literature Cited

Schmaltz, J. 2008. Phytoplankton bloom off New Zealand. In <http://earthobservatory.nasa.gov/IOTD/view.php?id=8381>. NASA.

Stevens, D.W., and O'Driscoll, R.L. 2006. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2005 (TAN0501). New Zealand Fisheries Assessment Report **13**.

Stevens, D.W., and O'Driscoll, R.L. 2007. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2006 (TAN0601). New Zealand Fisheries Assessment Report **5**.

Sutton, P. 2001. Detailed structure of the Subtropical Front over Chatham Rise, east of New Zealand. Journal of Geophysical Research. C. Oceans **106**: 31045-31056.

Zeldis, J.R., Murdoch, R.C., Cordue, P.L., and Page, M.J. 1998. Distribution of hoki (*Macruronus novaezelandiae*) eggs, larvae, and adults off Westland, New Zealand, and the design of an egg production survey to estimate hoki biomass. Canadian Journal of Fisheries and Aquatic Sciences **55**(7): 1682-1694.

APPENDIX 2.1: Temperature maps.

Temperature gradients at various depths over the Chatham Rise – Plots by Graham Rickard (NIWA), 2008.

APPENDIX 2.2: Current maps.

Currents on the Chatham Rise at various depths - Plots created by Graham Rickard (NIWA), 2008.

APPENDIX 2.3: Stomach removal protocol.

Protocol for collection of stomach samples: (Dunn 2005, 2006, 2007).

For each species and tow the sample will be selected in two phases: the first phase is a random sample; the second phase is a sample of the smallest and largest individuals. The second phase samples are required to tell us about changes in diet as the fish grow.

Using hoki as an example, if 20 randomly selected biological samples are taken from the tow for the trawl survey (for otoliths etc), then 5 of these fish will be sampled for stomachs. When these 20 are finished, another 3 fish should be sampled for stomachs, selected from the smallest and biggest available. The total number of biological samples would therefore be 23, of which 8 have been sampled for stomachs (5 random, 3 non-random). Fish from each sample (random, non-random) must be identified as such in the database, by putting “S” or “SN” respectively in the comments field. This protocol was modified at sea and the whole target number of fish was taken from the 20 “biological” fish with the 5 random samples being taken first and the remaining 3 non-random samples being the biggest or the smallest of the remaining fish.

If the tow is very busy, and it is clear that all the samples cannot be taken, then the priority will be to take the random samples.

Fish with clearly regurgitated or everted stomachs do not count towards the targets. All other fish are included (i.e. this includes those which look like they may have little or nothing in their stomachs).

Where a stomach is taken as part of a random sample, put an “SR” in the comments field.

Where a stomach is taken as part of a non-random sample, put an “SN” in the comments field.

To take the stomach:

1. Push any stomach contents from the oesophagus back into the stomach.
2. Seal the front of the stomach by fixing a zip-tie around the oesophagus.
3. Cut the oesophagus as far forward as possible in front of the tie.
4. Make a cut at the posterior end of the stomach at the pyloric caeca (i.e. below the pyloric sphincter), or fix a second tie.
5. Remove stomach and put in a seal-top plastic bag.
6. Complete the stomach sample label, and place the label in the bag
7. Seal the bag.
8. Freeze

Every stomach must be individually bagged and labelled. Ideally, keep stomachs from each station together in a single larger bag.

Smaller fish do not need to have stomachs removed, but can be returned whole.

Hoki Diet – Appendix 3

Stomachs must be labelled with trip code, tow (station number), species and fish number. Without these details the stomach can not be linked back to the biological fish details that are collected at sea (e.g. length, weight, sex), as well as environmental or physical factors like depth and location and time of day.

APPENDIX 2.4: Categorised prey groupings.

Table of prey contained in the 11 categories used in the multivariate analysis.

The groups are based largely on morphological similarities. Small bodied lantern fish, the morphologically similar Lighthouse fish *Vinciguerrria* spp., Pearlside *Maurolicus australis* and small mesopelagic fish were grouped together as LAN. Larger bodied Lantern fish, the morphologically similar Viper fish *Chauliodus sloani*, Hatchet fish and Snipe eel were grouped together as LAN 1. Similarly sized mesopelagic teleosts were grouped as MES 1. Unknown teleost remains and benthic teleosts were grouped as FIS. All rattails, including Javelin fish, were grouped together as RAT. All prawns, mysids and crustacea that could not be clearly identified as either prawns, euphausiids or mysids were grouped as PRA. Euphausiids were grouped independently as EUP. All other crustaceans were grouped together as CRU. All squids were grouped together as SQX. All prey that was unidentified was grouped as UNF. And all other prey, such as shell fragments, salps, foraminifera and radiolarians were grouped together as OTH (for other).

Species Code	Scientific name	Common name / Family	Category
APH	<i>Amphipoda</i>	Amphipod	CRU
APP	<i>Phronima</i> sp.	Hyperiid amphipod	CRU
CYM	<i>Cylopus magellanicus</i>	Sm brown amphipod	CRU
TGA	<i>Themisto gaudichaudi</i>	Amphipod	CRU
VBI	<i>Vibilia</i> spp.	Amphipod	CRU
COP	<i>Copepoda</i>	Copepod	CRU
CRB	Crab	Unknown crab	CRU
CRU	<i>Crustacea</i>	Crustacea	CRU
MGA	<i>Munida gracilis</i>	Galatheidæ	CRU
EUP	<i>Euphausiidae</i>	Euphausid	EUP
CDO	<i>Capromimu abbreviatus</i>	Capro dory	FIS
DCO	<i>Notophycis marginata</i>	Dwarf cod	FIS
FIS	Fish - Unknown	Fish - Unidentifiable	FIS
OTO	Otolith	Otolith	FIS
SCL	Scale	Scale	FIS
LAN	<i>Myctophidae</i>	Lantern fish	LAN
ELT	<i>Electrona</i> spp.	Lantern fish	LAN
PRO	<i>Protomyctophum</i> spp.	Lantern fish	LAN
VIN	<i>Vinciguerrria</i> spp.	Lighthouse fishes	LAN
LHE	<i>Lampanyctodes hectoris</i>	Lantern fish	LAN
MES	mesopelagic fish	Mid water fish	LAN
MMU	<i>Maurolicus australis</i>	Pearlside	LAN
DDA	<i>Diaphus danae</i>	Lantern fish	LAN 1
DIA	<i>Diaphus</i> spp.	Lantern fish	LAN 1
GYM	<i>Gymnoscopelus</i> spp.	Lantern fish	LAN 1

Hoki Diet – Appendix 4

Species Code	Scientific name	Common name / Family	Category
GYP	<i>Gymnoscopelus piabilis</i>	Lantern fish	LAN 1
LPA	<i>Lampanyctus</i> sp.	Lantern fish	LAN 1
LPD	<i>Lampadena</i> sp.	Lantern fish	LAN 1
SYM	<i>Symbolophorus</i> spp.	Lantern fish	LAN 1
HAT	<i>Sternoptychidae</i>	Hatchet fish	LAN 1
NEX	<i>Nemichthyidae</i>	Snipe eel	LAN 1
NSC	<i>Nemichthys scolapaceus</i>	Snipe eel	LAN 1
CHA	<i>Chauliodus sloani</i>	Viper fish	MES 1
MST	<i>Melanostomiidae</i>	Scaleless black dragonfish	MES 1
PAL	<i>Paralepididae</i>	Barracudinas	MES 1
SPL	<i>Scopelasaurus</i> spp.	Waryfishes	MES 1
APG	<i>Apigonidae</i>	Cardinal fish	MES 1
DSS	<i>Bathylagus</i> spp.	Deepsea smelt	MES 1
GST	<i>Gonostomatidae</i>	Lightfishes	MES 1
NAN	<i>Nansenia</i> spp.	Deepsea smelt	MES 1
PER	<i>Perparsia kopua</i>	Tubeshoulder	MES 1
SSI	<i>Argentina elongata</i>	Silverside	MES 1
PHO	<i>Photichthys argenteus</i>	Lighthouse fishes	MES 1
FLL	Shell Fragments	Shell Fragments	OTH
HTU	<i>Hyalinoecia tubicola</i>	Quill worm	OTH
ZFR	Foraminifera and radiolarians	Foraminifera and radiolarians	OTH
SAL	<i>Salpidae</i>	Salp	OTH
EPN	Euphausiid/Prawn/Mysid	Euphausiid/Prawn/Mysid	PRA
PAS	<i>Pasiphaea</i> spp.	Prawn	PRA
NAT	Natant decapod	Natant decapod	PRA
PRA	Prawn	Prawn - Unidentified	PRA
ACA	<i>AcanthePHYra</i> spp.	Prawn	PRA
APE	<i>AcanthePHYra pelagica</i>	Prawn	PRA
FUN	<i>Funchalia</i> spp.	Prawn	PRA
GNA	<i>Gnathophausia I.</i>	Mysid	PRA
MYS	<i>mysidae</i>	Mysid	PRA
SYD	<i>Systellapis debilis</i>	Prawn	PRA
NMA	<i>Notopandalus magnoculus</i>	Prawn	PRA
ONO	<i>Oplophorus novaeseelandiae</i>	Prawn	PRA
SER	<i>Sergestes</i> spp.	Prawn	PRA

Hoki Diet – Appendix 4

Species Code	Scientific name	Common name / Family	Category
COL	<i>Caelorinchus oliverianus</i>	Oliver's rattail	RAT
JAV	<i>Lepidorhynchus denticulatus</i>	Javelin fish	RAT
RAT	<i>Macrouridae</i>	Rattail	RAT
CAE	<i>Caelorinchus</i> spp.	Rattail	RAT
COY	<i>Coryohenoides</i> spp.	Rattail	RAT
CSU	<i>Caelorinchus subserrulatus</i>	Rattail	RAT
ASQ	<i>Nototodarus sloanii</i> & <i>gouldi</i>	Arrow squid	SQX
NOS	<i>Nototodarus sloanii</i>	Arrow squid	SQX
CPH	<i>Cephalopoda</i>	Cephlopod	SQX
MIQ	<i>Moroteuthis ingens</i>	Warty squid	SQX
NOD	<i>Nototeuthis dimegacotyle</i>	Squid	SQX
OSQ	<i>Octopoteuthiidae</i>	Octopus squid	SQX
SEQ	<i>Sepiolidae</i>	Bobtail squid (micky mouse)	SQX
SQX	Squid	Squid - unidentified	SQX
VSQ	<i>Histiotheuthis</i> spp.	Violet squid	SQX
UNF	Unidentifiable	Unidentifiable	UNF
UNI	Unidentified	Unidentified	UNF

Hoki Diet – Appendix 5

APPENDIX 2.5: All prey %IRI table.

Overall dietary compositions. Prey sorted by taxonomic group. Percentage weight (%W), percentage number (%N), percentage frequency of occurrence (%O), Index of Relative importance (IRI), and percent Index of Relative Importance (%IRI), and 95% confidence intervals. Unid = unidentified.

Prey	Code	%O	Lower 95%	Upper 95%	%W	Lower 95%	Upper 95%	%C	Lower 95%	Upper 95%	IRI	Lower 95%	Upper 95%	%IRI	Lower 95%	Upper 95%
Polychaeta																
<i>Hyalinoecia tubicola</i>	HTU	0.08	0.00	0.31	0.02	0.00	0.06	0.02	0.00	0.06	0.00	0.00	0.04	0.00	0.00	0.00
Crustacea																
Amphipoda	APH	6.61	4.66	8.72	0.04	0.02	0.07	2.34	1.36	3.69	15.76	6.84	30.47	0.56	0.23	1.14
Hyperiid Amphipoda	APP	0.31	0.00	0.76	0.10	0.00	0.29	0.25	0.00	0.79	0.11	0.00	0.71	0.00	0.00	0.03
<i>Cylopus magellanicus</i>	CYM	0.62	0.08	1.49	0.01	0.00	0.01	0.13	0.01	0.34	0.09	0.00	0.52	0.00	0.00	0.02
<i>Themisto gaudichandi</i>	TGA	2.41	1.26	3.50	0.03	0.01	0.04	0.53	0.27	0.81	1.34	0.36	2.86	0.05	0.01	0.10
<i>Vibilia</i> spp	VBI	0.70	0.15	1.31	0.01	0.00	0.02	0.18	0.03	0.35	0.13	0.01	0.43	0.01	0.00	0.02
<i>AcanthePHYra pelagica</i>	APE	0.08	0.00	0.38	0.05	0.00	0.24	0.02	0.00	0.07	0.01	0.00	0.12	0.00	0.00	0.00
<i>AcanthePHYra</i> spp	ACA	0.16	0.00	0.48	0.04	0.00	0.16	0.06	0.00	0.21	0.02	0.00	0.17	0.00	0.00	0.01
Decapoda - crab	CRB	0.08	0.00	0.39	0.00	0.00	0.01	0.02	0.00	0.07	0.00	0.00	0.03	0.00	0.00	0.00
Decapoda - prawn	PRA	9.80	7.53	12.41	0.20	0.12	0.34	2.11	1.48	2.99	22.64	13.17	38.31	0.80	0.44	1.44
<i>Funchalia</i> spp	FUN	0.08	0.00	0.32	0.48	0.00	2.09	0.02	0.00	0.07	0.04	0.00	0.74	0.00	0.00	0.03
<i>Munida gracilis</i>	MGA	0.08	0.00	0.32	0.01	0.00	0.03	0.02	0.00	0.07	0.00	0.00	0.03	0.00	0.00	0.00
<i>Notopandalus magnoculus</i>	NMA	1.48	0.50	2.70	0.34	0.07	0.82	0.56	0.12	1.25	1.33	0.12	5.19	0.05	0.00	0.18
<i>Oplophorus novaezeelandiae</i>	ONO	1.09	0.45	1.93	0.54	0.19	1.06	0.27	0.10	0.53	0.87	0.13	2.80	0.03	0.01	0.10
<i>Pasiphaea</i> spp	PAS	19.30	15.95	22.66	5.20	3.29	8.00	9.88	6.64	14.36	290.74	177.20	459.98	10.30	6.10	16.78
<i>Sergestes</i> spp	SER	11.70	8.90	14.67	1.76	1.03	2.84	4.97	2.86	7.80	79.04	39.12	136.00	2.81	1.38	5.07
<i>Systellaspis debilis</i>	SYD	0.08	0.00	0.33	0.00	0.00	0.01	0.02	0.00	0.07	0.00	0.00	0.03	0.00	0.00	0.00
Euphausiacea	EUP	29.50	24.90	34.04	1.30	0.75	2.16	40.30	24.11	53.21	1224.60	671.71	1744.80	43.50	25.95	56.42
<i>Gnathophausia</i> spp	GNA	0.08	0.00	0.32	0.01	0.00	0.06	0.02	0.00	0.07	0.00	0.00	0.04	0.00	0.00	0.00
Mysidae	MYS	0.54	0.08	1.10	0.01	0.00	0.01	0.12	0.02	0.27	0.07	0.00	0.30	0.00	0.00	0.01
Unid. Crustacean	CRU	0.70	0.16	1.40	0.00	0.00	0.01	0.15	0.03	0.32	0.11	0.01	0.43	0.00	0.00	0.02
Copepoda	COP	0.93	0.24	1.79	0.00	0.00	0.00	0.21	0.06	0.43	0.19	0.02	0.73	0.01	0.00	0.03
Malacostraca	EPN	5.60	3.84	7.35	0.05	0.02	0.10	1.36	0.84	2.10	7.89	3.68	15.39	0.28	0.13	0.56
Teleostei																
Nemichthyidae	NEX	0.16	0.00	0.47	0.24	0.00	0.84	0.03	0.00	0.09	0.04	0.00	0.40	0.00	0.00	0.01
<i>Nemichthys scolapaceus</i>	NSC	0.08	0.00	0.38	0.13	0.00	0.57	0.02	0.00	0.07	0.01	0.00	0.22	0.00	0.00	0.01

Hoki Diet – Appendix 5

Prey	Code	%O			%W			%N			IRI			%IRI		
		%O	Lower 95%	Upper 95%	%W	Lower 95%	Upper 95%	%C	Lower 95%	Upper 95%	IRI	Lower 95%	Upper 95%	%IRI	Lower 95%	Upper 95%
Paralepididae	PAL	0.08	0.00	0.32	0.52	0.00	2.24	0.02	0.00	0.07	0.04	0.00	0.75	0.00	0.00	0.03
<i>Scopelosaurus</i> spp	SPL	0.31	0.00	0.82	5.32	0.00	14.33	0.06	0.00	0.16	1.67	0.00	11.03	0.06	0.00	0.40
<i>Austrophycis marginata</i>	DCO	0.08	0.00	0.39	0.00	0.00	0.01	0.02	0.00	0.08	0.00	0.00	0.03	0.00	0.00	0.00
<i>Caelorinchus oliverianus</i>	COL	1.17	0.48	2.05	4.70	0.23	11.61	0.34	0.13	0.68	5.88	0.33	22.08	0.21	0.01	0.80
<i>Coryphaenoides</i> spp	COY	0.16	0.00	0.55	0.35	0.00	1.46	0.03	0.00	0.11	0.06	0.00	0.75	0.00	0.00	0.03
<i>Coryphaenoides subterrulatus</i>	CSU	0.08	0.00	0.32	0.85	0.00	3.51	0.02	0.00	0.06	0.07	0.00	1.10	0.00	0.00	0.04
<i>Lepidorhynchus denticulatus</i>	JAV	1.94	1.00	3.20	21.70	7.19	37.92	0.38	0.18	0.69	42.86	9.33	106.33	1.52	0.32	3.88
<i>Caelorinchus</i> spp.	CAE	0.47	0.08	1.09	0.69	0.00	2.72	0.09	0.01	0.21	0.37	0.00	2.24	0.01	0.00	0.09
Macrouridae	RAT	1.40	0.55	2.38	2.79	0.22	6.40	0.30	0.11	0.57	4.32	0.32	15.02	0.15	0.01	0.55
<i>Diaphus danae</i>	DDA	0.78	0.23	1.56	0.68	0.12	1.65	0.15	0.04	0.33	0.64	0.05	2.86	0.02	0.00	0.10
<i>Diaphus</i> spp	DIA	0.78	0.23	1.50	0.27	0.04	0.63	0.15	0.04	0.32	0.32	0.02	1.31	0.01	0.00	0.05
<i>Electrona</i> spp	ELT	0.70	0.16	1.38	0.26	0.03	0.65	0.13	0.03	0.28	0.27	0.02	1.09	0.01	0.00	0.04
<i>Gymnoscopelus</i> spp	GYM	0.47	0.08	1.03	0.42	0.02	1.45	0.09	0.01	0.20	0.24	0.00	1.35	0.01	0.00	0.05
<i>Lampadena</i> spp	LPD	0.16	0.00	0.52	0.17	0.00	0.56	0.03	0.00	0.10	0.03	0.00	0.33	0.00	0.00	0.01
<i>Lampanyctodes hectoris</i>	LHE	12.20	9.43	14.92	4.10	2.45	6.45	4.66	3.23	6.68	106.90	61.13	176.71	3.79	2.21	6.26
<i>Lampanyctus</i> spp	LPA	0.31	0.00	0.97	0.99	0.00	3.97	0.18	0.00	0.68	0.36	0.00	3.84	0.01	0.00	0.14
Myctophidae	LAN	33.50	29.46	37.36	7.94	4.16	13.86	14.70	10.90	19.74	757.66	509.35	1068.90	26.90	18.21	36.83
<i>Protomyctophum</i> spp	PRO	0.08	0.00	0.39	0.00	0.00	0.00	0.03	0.00	0.15	0.00	0.00	0.06	0.00	0.00	0.00
<i>Symbolophorus</i> spp	SYM	0.31	0.00	0.79	0.32	0.00	1.07	0.07	0.00	0.21	0.12	0.00	0.86	0.00	0.00	0.03
Unid. Mesopelagic teleost	MES	6.38	4.76	8.71	1.01	0.39	1.99	3.45	1.52	6.61	28.42	11.93	67.16	1.01	0.40	2.43
Actinopterygii - otolith	OTO	1.94	0.85	3.35	0.01	0.00	0.01	0.56	0.21	1.07	1.10	0.18	3.35	0.04	0.01	0.13
Actinopterygii - scale	SCL	19.10	16.08	22.20	0.10	0.04	0.20	3.61	2.64	4.94	70.73	46.32	107.23	2.51	1.49	4.16
Unid. Teleost	FIS	8.01	5.67	10.32	6.47	2.13	14.53	1.92	1.12	2.92	67.13	24.71	150.05	2.38	0.83	5.35
<i>Nansenia</i> spp	NAN	0.23	0.00	0.66	0.78	0.00	3.11	0.04	0.00	0.13	0.19	0.00	1.71	0.01	0.00	0.06
<i>Persparia kopua</i>	PER	0.23	0.00	0.65	1.20	0.00	3.96	0.04	0.00	0.13	0.29	0.00	2.48	0.01	0.00	0.09
Apogonidae	APG	0.08	0.00	0.33	0.08	0.00	0.35	0.02	0.00	0.07	0.01	0.00	0.15	0.00	0.00	0.01
<i>Argentina elongata</i>	SSI	0.08	0.00	0.32	0.56	0.00	2.59	0.02	0.00	0.07	0.05	0.00	0.89	0.00	0.00	0.03
<i>Bathylagus</i> spp	DSS	0.16	0.00	0.53	0.80	0.00	3.47	0.03	0.00	0.10	0.13	0.00	1.39	0.01	0.00	0.05
Melanostomiidae	MST	0.08	0.00	0.32	0.14	0.00	0.59	0.02	0.00	0.06	0.01	0.00	0.20	0.00	0.00	0.01
<i>Chauliodus sloani</i>	CHA	0.39	0.00	0.89	1.26	0.00	4.35	0.09	0.00	0.23	0.53	0.00	3.41	0.02	0.00	0.12
<i>Maurollicus australis</i>	MMU	7.15	4.85	9.33	1.92	0.61	4.24	2.73	1.40	4.63	33.25	11.44	72.97	1.18	0.40	2.72
<i>Photichthys argenteus</i>	PHO	2.57	1.45	3.90	16.20	6.55	26.01	0.56	0.30	0.91	42.99	11.05	94.78	1.53	0.38	3.44
Sternoptychidae	HAT	0.31	0.00	0.78	0.03	0.00	0.09	0.06	0.00	0.15	0.03	0.00	0.17	0.00	0.00	0.01
<i>Vinciguerra</i> spp	VIN	0.23	0.00	0.67	0.03	0.00	0.10	0.04	0.00	0.13	0.02	0.00	0.13	0.00	0.00	0.01

Hoki Diet – Appendix 5

Prey	Code	%O	%O Lower 95%	Upper 95%	%W	%W Lower 95%	Upper 95%	%C	%N Lower 95%	Upper 95%	IRI	IRI Lower 95%	Upper 95%	%IRI	%IRI Lower 95%	Upper 95%
<i>Capromimus abbreviatus</i>	CDO	0.31	0.00	0.87	1.13	0.00	3.52	0.06	0.00	0.18	0.37	0.00	2.99	0.01	0.00	0.12
Mollusca																
Cephalopoda	CPH	1.32	0.47	2.34	0.04	0.00	0.16	0.28	0.09	0.53	0.43	0.05	1.46	0.02	0.00	0.05
Sepiolidae	SEQ	0.16	0.00	0.54	0.02	0.00	0.11	0.03	0.00	0.10	0.01	0.00	0.10	0.00	0.00	0.00
<i>Nototodarus</i> spp	ASQ	0.31	0.00	0.76	0.28	0.00	1.08	0.06	0.00	0.15	0.11	0.00	0.75	0.00	0.00	0.03
<i>Moroteuthis ingens</i>	MIQ	0.08	0.00	0.39	0.10	0.00	0.50	0.02	0.00	0.07	0.01	0.00	0.22	0.00	0.00	0.01
<i>Nototeuthis dimegacotyle</i>	NOD	0.08	0.00	0.37	0.01	0.00	0.03	0.02	0.00	0.07	0.00	0.00	0.04	0.00	0.00	0.00
<i>Nototodarus sloanii</i>	NOS	0.16	0.00	0.55	3.24	0.00	11.44	0.03	0.00	0.11	0.51	0.00	6.37	0.02	0.00	0.23
Octopoteuthidae	OSQ	0.31	0.00	0.84	0.01	0.00	0.02	0.06	0.00	0.16	0.02	0.00	0.15	0.00	0.00	0.01
Teuthoidea	SQX	1.79	0.91	2.89	0.65	0.03	2.68	0.38	0.18	0.69	1.85	0.27	7.49	0.07	0.01	0.27
<i>Histioteuthis</i> spp.	VSQ	0.54	0.08	1.19	0.60	0.00	2.44	0.10	0.02	0.24	0.38	0.00	2.34	0.01	0.00	0.09
Shell fragments Mollusca	FLL	0.16	0.00	0.55	0.00	0.00	0.01	0.03	0.00	0.11	0.01	0.00	0.07	0.00	0.00	0.00
Salpidae	SAL	2.18	1.16	3.30	0.24	0.07	0.53	0.72	0.25	1.63	2.10	0.41	6.38	0.08	0.01	0.24
Foraminifera/Radiozoa	ZFR	0.08	0.00	0.38	0.00	0.00	0.01	0.03	0.00	0.14	0.00	0.00	0.05	0.00	0.00	0.00
Unidentifiable	UNF	0.16	0.00	0.54	0.04	0.00	0.19	0.03	0.00	0.10	0.01	0.00	0.14	0.00	0.00	0.01
Unidentified	UNI	0.16	0.00	0.48	0.03	0.00	0.14	0.03	0.00	0.10	0.01	0.00	0.10	0.00	0.00	0.00

APPENDIX 2.6: Categorical data %IRI table.

Overall dietary compositions. Prey in groups used for multivariate analysis. Percentage weight (%W), percentage number (%N), percentage frequency of occurrence (%O), Index of Relative importance (IRI), and percent Index of Relative Importance (%IRI), and 95% confidence intervals.

Prey	% O			%W			%N			IRI			%IRI		
	%O	Lower 95%	Upper 95%	%W	Lower 95%	Upper 95%	%N	Lower 95%	Upper 95%	IRI	Lower 95%	Upper 95%	%IRI	Lower 95%	Upper 95%
LAN	48.06	43.95	52.65	15.24	10.02	23.44	25.72	18.80	33.87	1968.31	1419.48	2649.88	38.43	29.06	48.48
EUP	29.47	24.74	33.92	1.30	0.82	2.14	40.25	25.27	54.49	1224.59	702.43	1781.77	23.91	13.34	35.06
PRA	41.37	36.84	45.60	8.69	5.95	12.62	19.38	13.96	25.96	1160.87	794.56	1608.29	22.67	16.15	30.30
FIS	29.08	24.88	33.29	7.70	2.91	15.83	6.16	4.37	8.42	403.19	221.58	716.60	7.87	4.38	13.40
RAT	4.82	3.21	6.59	31.06	15.88	46.55	1.15	0.68	1.76	155.27	63.77	286.38	3.03	1.21	5.76
MES 1	4.20	2.67	5.95	27.17	14.34	40.63	0.91	0.53	1.43	117.92	41.45	233.92	2.30	0.79	4.73
CRU	11.98	9.33	14.72	0.19	0.07	0.39	3.82	2.49	5.32	47.96	24.72	78.22	0.94	0.48	1.51
SQX	4.74	3.22	6.48	4.95	0.43	12.84	0.97	0.61	1.49	28.10	5.51	76.41	0.55	0.10	1.52
LAN 1	3.03	1.79	4.40	3.37	1.30	6.60	0.78	0.39	1.46	12.60	3.39	30.94	0.25	0.07	0.60
OTH	2.41	1.37	3.73	0.26	0.09	0.52	0.80	0.32	1.67	2.55	0.64	7.20	0.05	0.01	0.14
UNF	0.31	0.00	0.79	0.08	0.00	0.24	0.06	0.00	0.16	0.04	0.00	0.30	0.00	0.00	0.01

APPENDIX 2.7: Prey species codes.

Prey species codes used in the dietary analysis of hoki listed alphabetically by species code.

Species Code	Scientific name	Common name / Family
ACA	<i>AcanthePHYra</i> spp.	Prawn
APE	<i>AcanthePHYra pelagica</i>	Prawn
APG	<i>Apigonidae</i>	Cardinal fish
APH	<i>Amphipoda</i>	Amphipod
APP	<i>Phronima</i> sp.	Hyperiid amphipod
ASQ	<i>Nototodarus sloanii & gouldi</i>	Arrow squid
CAE	<i>Caelorinchus</i> spp.	Rattail
CDO	<i>Capromimu abbreviatus</i>	Capro dory
CHA	<i>Chauliodus sloani</i>	Viper fish
COL	<i>Caelorinchus oliverianus</i>	Oliver's rattail
COP	<i>Copepoda</i>	Copepod
COY	<i>Coryohenoides</i> spp.	Rattail
CPH	<i>Cephlopoda</i>	Cephlopod
CRB	Crab	Unknown crab
CRU	<i>Crustacea</i>	Crustacea
CSU	<i>Caelorinchus subserrulatus</i>	Rattail
CYM	<i>Cyllopus magellanicus</i>	Sm brown amphipod
DCO	<i>Notophycis marginata</i>	Dwarf cod
DDA	<i>Diaphus danae</i>	Lantern fish
DIA	<i>Diaphus</i> spp.	Lantern fish
DSS	<i>Bathylagus</i> spp.	Deepsea smelt
ELT	<i>Electrona</i> spp.	Lantern fish
EPN	Euphausid/Prawn/Mysid	Euphausid/Prawn/Mysid
EUP	<i>Euphausidae</i>	Euphausid
FIS	Fish - Unknown	Fish - Unidentifiable
FLL	Shell Fragments	Shell Fragments
FUN	<i>Funchalia</i> spp.	Prawn
GNA	<i>Gnathophausia I.</i>	Mysid
GST	<i>Gonostomatidae</i>	Lightfishes
GYM	<i>Gymnoscopelus</i> spp.	Lantern fish
GYP	<i>Gymnoscopelus piabilis</i>	Lantern fish
HAT	<i>Sternoptychidae</i>	Hatchet fish
HTU	<i>Hyalinoecia tubicola</i>	Quill worm
JAV	<i>Lepidorhynchus denticulatus</i>	Javelin fish

Hoki Diet – Appendix 7

Species Code	Scientific name	Common name / Family
LAN	<i>Myctophidae</i>	Lantern fish
LHE	<i>Lampanyctodes hectoris</i>	Lantern fish
LPA	<i>Lampanyctus</i> sp.	Lantern fish
LPD	<i>Lampadena</i> sp.	Lantern fish
MES	mesopelagic fish	Mid water fish
MGA	<i>Munida gracilis</i>	Galatheididae
MIQ	<i>Moroteuthis ingens</i>	Warty squid
MMU	<i>Maurololicus australis</i>	Pearlside
MST	<i>Melanostomiidae</i>	Scaleless black dragonfish
MYS	<i>mysidae</i>	Mysid
NAN	<i>Nansenia</i> spp.	Deepsea smelt
NAT	Natant decapod	Natant decapod
NEX	<i>Nemichthyidae</i>	Snipe eel
NMA	<i>Notopandalus magnoculus</i>	Prawn
NOD	<i>Nototeuthis dimegacotyle</i>	Squid
NOS	<i>Nototodarus sloanii</i>	Arrow squid
NSC	<i>Nemichthys scolapaceus</i>	Snipe eel
ONO	<i>Oplophorus novaeseelandiae</i>	Prawn
OSQ	<i>Octopoteuthiidae</i>	Octopus squid
OTO	Otolith	Otolith
P01	<i>Cestoda</i>	Tape worm
P02	<i>Nematoda</i>	Round worm
P03	<i>Trematoda</i>	Fluke
P04	<i>Codepoda</i>	Parasitic copepod
PAL	<i>Paralepididae</i>	Barracudinas
PAS	<i>Pasiphaea</i> spp.	Prawn
PER	<i>Perparsia kopua</i>	Tubeshoulder
PHO	<i>Photichthys argenteus</i>	Lighthouse fishes
PRA	Prawn	Prawn - Unidentified
PRO	<i>Protomyctophum</i> spp.	Lantern fish
RAT	<i>Macrouridae</i>	Rattail
SAL	<i>Salpidae</i>	Salp
SCL	Scale	Scale
SEQ	<i>Sepiolidae</i>	Bobtail squid (micky mouse)
SER	<i>Sergestes</i> spp.	Prawn
SLN	Stomach lining	Stomach lining
SPL	<i>Scopelasaurus</i> spp.	Waryfishes
SQX	Squid	Squid - unidentified

Hoki Diet – Appendix 7

Species Code	Scientific name	Common name / Family
SSI	<i>Argentina elongata</i>	Silverside
SYD	<i>Systellapis debilis</i>	Prawn
SYM	<i>Symbolophorus</i> spp.	Lantern fish
TGA	<i>Themisto gaudichaudi</i>	Amphipod
UNF	Unidentifiable	Unidentifiable
UNI	Unidentified	Unidentified
VBI	<i>Vibilia</i> spp.	Amphipod
VIN	<i>Vinciguerria</i> spp.	Lighthouse fishes
VSQ	<i>Histiotheuthis</i> spp.	Violet squid
ZFR	Foraminifera and radiolarians	Foraminifera and radiolarians

3 The Mid-Domain Effect and Body-Size Depth Distribution in a Demersal Marine Fish Assemblage in New Zealand.

3.1 ABSTRACT

The cause for the patterns in species distribution has long been a question ecologists have tried to answer. One school of thought leans towards adaptive or environmental processes that shape the communities and the niches the species found there inhabit. The other side of the coin is neutral theory which predicts that species within a community are allocated by random chance. The mid-domain effect predicts species richness, and thus distribution, to be due to geometric constraints with the greatest species richness to be found at the centre of a geographically constrained domain. In marine systems there is another pattern often seen, that of bigger individuals being found in deeper water. Here I show evidence for the mid-domain effect on species richness on a demersal fish assemblage on the Chatham Rise, New Zealand. I found no interspecific pattern for body-size depth distribution with the entire species assemblage, nor when the assemblage was split by class into Osteichthyes (bony fish) and Chondrichthyes (cartilaginous fish), which supports our findings of the mid-domain effect. At a community level, species distribution over the studied depth range was largely explained by the geometric constraints of the mid-domain effect, while at a species level, distribution over depth was often a reflection of body size. Some species had large individuals deep while other species had small. Overall this supports the hypothesis that competition or adaptation works more strongly at a population or species level, than on the overall community who's species distribution can more often be attributed to random chance.

3.2 INTRODUCTION

Changes in species richness with latitude, topographical elevation and depth are patterns that have been frequently observed (Brown 2001, Macpherson and Duarte 1994, Rex and Etter 1998, Rohde 1992, Rohde et al. 1993, Stevens 1989). The question of which mechanisms are driving or causing these gradients remains unanswered. Colwell and Hurtt (1994) and Willig and Lyons (1998) suggested that species latitudinal ranges could be randomly shuffled within the geometric constraints of a biogeographical domain, with the requirement that species ranges did not extend beyond the outer limits of the domain, and consequently produce a pattern of species richness with the richness peak in the centre of the domain and reducing towards the domain boundaries. Colwell and Lees (2000) called this the mid-domain effect (MDE), and suggested that it might explain a significant portion of the observed latitudinal gradients. This species richness approach has also been applied to depth (Kendall and Haedrich 2006, Rohde et al. 1993, Smith and Brown 2002) and species richness gradients over topographical elevation (Brehm et al. 2007, McCain 2004). Strict mid-domain effect models attempt to solely explain species richness gradients without the considerations of external factors such as environmental or evolutionary events. These have been considered null models, as the observed pattern of species richness should be identical to that predicted by the model based on observed species ranges (Colwell et al. 2004, 2005). The mid-domain effect is contentious with some arguing that it does not fulfil the requirements of a null model (Hawkins and Diniz-Filho 2002, Hawkins et al. 2005, Zapata et al. 2005), as it is not truly independent of external factors. A number of studies have found support for the mid-domain effect (Dunn et al. 2007, Jetz and Rahbek 2001, Lees and Colwell 2007, Pineda and Caswell 1998, Rahbek et al. 2007, Rohde et al. 1993, Romdal et al. 2005), while others have reported little relationship between the observed and expected species richness patterns as predicted by the mid-domain effect (Diniz-Filho et al. 2002, Hawkins and Diniz-Filho 2002, Kendall and Haedrich 2006, Smith and Brown 2002).

Two contrasting paradigms exist in describing community ecology, that of neutrality and niche theory. Neutral theory was developed by Hubbell (1979, 2001, 2003, 2005). The

theory essentially considers the ecological properties of every individual in the population to be identical. Individuals compete for the same, fully exploited pool of resources, but the identity of the winner of any single contest is left to chance (Whitfield 2002). Hubbell (2001) has shown that neutral theory is capable of summarizing a surprising array of empirical patterns of species abundances. Neutral theory is considered by some ecologists as a radical shift from established niche theories (Chave 2004). Niche theory looks at how the differences between species allows them to co-exist (Whitfield 2002). There are two types of niche: the fundamental niche, the range of environmental conditions within which a species can live indefinitely in the absence of negative interactions (i.e. competition, predation, parasitism); and the realised niche, the environmental conditions / resources that the species actually exists in / uses in the presence of negative interactions (Krebs 2001). The nice concept implicitly assumes that neighbouring species have negative impacts (or niche shrinking affect) on one another and is firmly bound to the notion of competitive exclusion (that no two species can occupy the same niche) (Krebs 2001). Niche theory considers communities to be structured / organised based on species assemblages of those species which can best utilise the available resources (Nybakken 2001). However niche and neutral theories do not need to be conflicting, but can rather be viewed together as complementary (Leibold and McPeck 2006). Theories of coexistence by niche differentiation are mostly concerned with deterministic processes, and a small number of species that interact through fixed rules, as prescribed in the Lotka-Volterra equations. By contrast, the neutral theory is primarily concerned with species-rich communities (tropical forests, coral reefs) with many rare species, where the role of stochasticity at an individual scale becomes unavoidable (Chave 2004). Neutral theory has also been used as the null hypothesis when testing null models (Dornelas et al. 2006). In this vein the mid-domain effect has been suggested as a null model by some, due to its exclusion of environmental factors (Connolly 2005, Kendall and Haedrich 2006). Colwell et al. (2004) suggest that rather than being considered a null that can be rejected or accepted, the mid-domain effect should be assessed on an equal statistical footing as other possible explanations of varying species richness, with the context of multiple causality.

MDE & Body-Size Depth Distribution – Introduction

The ocean has different conditions at different depths, and different species inhabit these depths. Shallower water is warmer, has less hydrostatic pressure, and is potentially more nutrient rich or contains an abundant primary food source (i.e. phytoplankton, zooplankton) while deeper water is colder, has greater hydrostatic pressure and has no primary productivity, besides that which ‘rains’ down from the photic zone (Nybakken and Bertness 2005). One easily testable difference between species from different locations is body size. It has been hypothesised that differences in body size between species from shallower and deeper water may be an adaptation for survival in the different environments (Macpherson and Duarte 1991).

A large number of studies have observed that large individuals of marine demersal species are found in deeper water (Macpherson 1981, Macpherson and Duarte 1991, Perry et al. 2005, Sarda and Cartes 1993, Swain 1997). This pattern has so frequently been observed that it has been considered a law (Heinecke’s law) (Cushing and Walsh 1976). There has been much speculation on the cause of this phenomenon, as well as disagreement on the validity of its “law” status (Olabarria and Thurston 2003, Rex and Etter 1998, Uiblein et al. 1996). Those in favour of this phenomenon have hypothesised that the causes are inter and intraspecific competition (Helfman 1978) or predation (Cushing 1975, Cushing and Walsh 1976). One method of predator avoidance is spatial segregation (Keast 1978). It has also been postulated that temperature is a dominant causal factor for this “large-deep and small-shallow” phenomenon. Warmer shallow waters are more favourable for growth rates, therefore they may be taken advantage of by smaller or younger fish in the early stages of life when growth rates are high. Cooler, deeper waters have also been found to reduce the metabolic energy costs of fish (Sogard and Olla 1996). Large fish have a higher energy need than smaller fish and therefore it is advantageous for them to minimise their energy expenditure by any means possible. Larger fish have also been found to have a larger thermal tolerance than smaller fish and therefore are physically able to survive in colder, deeper waters where smaller individuals are not (Brill et al. 1999).

MDE & Body-Size Depth Distribution – Introduction

There are two broad classes of fish, Class Chondrichthyes and Superclass Osteichthyes which have some differences between them, morphologically, reproductively and physiologically. The Osteichthyes or better known as bony fish, have an ossified (or bony) skeleton, external fertilisation, are usually egg laying and usually have a larval stage of development (Bone and Moore 2008). The Chondrichthyes have a cartilaginous skeleton, internal fertilisation, give birth to live young and don't have a larval stage of development (Carrier et al. 2004a).

Competition occurs between species as well as within a single species. Spatial separation is a mechanism that decreases competition between individuals. If two individuals are present in the same place then the direct competition between them is lessened. Thus separation between the large and small individuals of a species may enhance the survival rates of the small fish by lessening competition pressure. Additionally, when fish are small they may be considered prey for the larger individuals of the species and spatial separation may also be a way of predator avoidance.

This study tests for evidence of the mid-domain effect. It also investigates whether overall, larger bodied fish were found in deeper water, whether there was a difference between Osteichthyes (bony fish, e.g. barracouta) and Chondrichthyes (cartilaginous fish, e.g. sharks) in body-size depth trends, and on an individual species basis, whether these trends were present and in what way.

3.3 MATERIALS AND METHODS

3.3.1 Study Area

The Chatham Rise is a prominent bathymetric ridge that projects about 500 nautical miles (n. miles) (926 km) east from Banks Peninsula, on the east coast of the South Island, to the Chatham Islands, New Zealand (Figure 3.1). It has a depth varying from about 250 m down to about 3000 m. Pacific deep water is found at the depths below about 2000 m,

and has a temperature of 1–2°C with no appreciable seasonal variation. Above this, the Chatham Rise is an area where the subtropical convergence forms, as warmer and more saline subtropical water from the north meets subantarctic water from the south (Sutton 2001). The surface water temperatures on the Chatham Rise recorded during trawl surveys have consequently been about 2–3°C warmer on the northern side, and varies seasonally by about 3–4°C. The bottom water temperatures at depths greater than 700 m, recorded during trawl surveys have been much less variable, being warmer on the northern side, but by <1°C, and varying seasonally by <1°C. At 300m depth the average temperature on the Northern Rise is 10–11°C and at 800m it is 7–8°C, while in comparison the average temperature on the Southern Rise at 300m is 8–9°C and 4–6°C at 800m. The subtropical convergence runs from about 45° S on the west coast of the South Island, around the bottom of the South Island, up the east coast, then extends (to a variable extent) through the Mernoo Gap at the western end of the Chatham Rise, before turning southwards and running eastwards along the Chatham Rise. Currents on the Chatham Rise move from west to east, they come down the eastern coast of the North Island from the north and up the eastern coast of the South Island from the south, hit the Rise and move east, oscillating north and south as it does so on either side of the Rise.

Primary productivity is relatively high on the Chatham Rise compared with other offshore areas (Bradford-Grieve et al. 1999, Bradford-Grieve et al. 1997, Murphy et al. 2001). Zooplankton productivity has also been observed to be high (McClatchie et al. 2004) and mesopelagic biomass has been reported as high (McClatchie and Dunford 2003). This high productivity is widely believed to be the underlying reason why the Chatham Rise supports major fisheries such as hoki (Annala et al. 2004). When compared to the nearby subantarctic, the Chatham Rise has higher species diversity and richness (McClatchie et al. 1997).

Figure 3.1: New Zealand with 1000 m depth contour, in relation to the Chatham Rise, dark rectangle (Top). The Chatham Rise from Banks Peninsula on the left to the Chatham Islands on the right (Bottom).

3.3.2 Sampling

Sampling took place on New Zealand Ministry of Fisheries funded annual trawl surveys for hoki and middle depth species on the Chatham Rise, New Zealand that have been run since 1992 by the National Institute of Water and Atmospheric Research Ltd (NIWA). Survey details and results can be found in Stevens and O'Driscoll (2006, 2007). Briefly, sampling was conducted for roughly four weeks from late December to late January each year. Samples for this study were collected from surveys fishing in January 2005, 2006, and 2007. Fish were captured in bottom trawls, the net was an 8-seam hoki bottom trawl with 100m sweeps, 50m bridles, 12m backstops, 58.8m ground rope, 45m headline and 60mm cod end mesh (see Hurst and Bagley (1994) for net plan and rigging details). The

sample design was 2-phase stratified random, after Francis (1984). The same 26 strata were used consistently. Phase-1 trawl stations were allocated from simulations based on catch rates from the previous three annual trawl surveys, after Bull et al. (2000). At each station sampled the trawl was towed for approximately 3 nautical miles (n. miles) at a speed over ground of 3.5 knots (kts) (Hurst et al. 1992). The catch was sorted by species and either the whole catch or a representative sub sample of the fishes are weighed, to the nearest gram (on electronic, calibrated, motion compensated balances) and measured to the nearest millimetre. Fish length, weight and sex were recorded with the latitudinal and longitudinal location of each trawl station. The fish species used in this analysis are the 30 most abundant species caught over the three sample years. The criteria of most abundant were considered to be those species with an estimated biomass of > 800 tonnes.

3.3.3 Mid-Domain Effect

To test for a mid-domain effect the data were analysed using Colwell's (2006) RangeModel. The mid-domain effect predicts a parabolic curve in species richness over the domain with the peak of species richness being found in the centre of the domain, due to the random shuffling of observed species ranges. This approach that species with large ranges have their range mid point towards the centre of the domain, while species with smaller ranges can have their range mid points towards the domain edges (Colwell and Lees 2000). Following Lees et al. (1999) and Romdal et al. (2005) the predictive power of the RangeModel was tested by the R^2 value of a linear regression on the observed and expected species richness. The likelihood of sampling a specific species increases with the abundance of that species and the amount of sampling effort. Given that sampling effort was comparatively evenly spread over the study area, species abundances will affect the likelihood of collecting individuals of a particular species from the minimum and maximum of their depth range. There is a greater likelihood of sampling the actual minimum and maximum of abundant species ranges, while under estimating the range limits of comparatively rarer species. To take into account sampling bias of the more abundant species, one or two standard deviations around the average depth were used as the depth limits of a species; the average depth was considered the range mid point. This

reduced the range size of all the species, giving a more standardised approach across all species sampled.

3.3.4 Body-size depth relationship

Depth variation is a mechanism that may separate predators, remove competition, or limit niche overlap. Average body size was plotted against average depth for the complete species assemblage, a linear regression was run to test for significance of any relationship found. A significant relationship would be either a positive or negative trend of body size increasing with increasing depth with a p-value of less than 0.05. Given their differences, samples were also split by Class Osteichthyes and Chondrichthyes, both cumulatively and by species to see whether the same relationship of body size and depth was shown, using linear regressions of size variance to test for significance of any relationship. Intraspecific length / depth relationships were plotted for the 30 species studied, and linear regressions run to again test for the significance of any relationship. Nine species were not plotted individually as there were less than ten sample records or there had been inconsistent sampling over the entire depth range. Regression analyses were performed using SPSS v16 software (SPSS 2007).

3.4 RESULTS

3.4.1 Mid-domain effect

The 30 species conformed to the predictions of the mid-domain effect. Species with large ranges have their range mid-point towards the centre of the domain, as well as showing that species richness increased towards the centre of the domain (Figure 3.2). The observed species richness fit the RangeModel predictions well (Figure 3.3 (A)), falling predominately within the 95% confidence intervals, but with some deviations. Deviations occurred at the shallowest and deepest depths (Figure 3.3 (A)). The R^2 value from the plot of the observed species richness against the expected species richness (Figure 3.3 (B))

MDE & Body-Size Depth Distribution – Results

Figure 3.2: Output of the RangeModel showing the depth ranges of the 30 study species displayed as horizontal lines centred around their depth midpoints.

Figure 3.3: (A) Observed species richness (circles) over the depth domain, plotted with the mean expected species richness (solid line), as predicted by Colwell's Range Model, and the upper and lower 95% confidence intervals (broken lines). (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel. Line of best fit fitted through the data. $R^2 = 0.7732$ ($p < 0.001$).

was $R^2 = 0.7732$ ($p < 0.001$) indicating a good fit to the model's predictions. Using one and two standard deviations around the average depth of each species for the species

range, which removed the sample bias given to more abundant species, the predictions of the mid-domain effect were still met (Figure 3.4 and Figure 3.5). Both produced a parabolic curve of species richness peaking towards the middle of the domain, with the observed species richness falling predominately within the 95% confidence intervals of the expected species richness, but with some deviation. Deviations occurred at the shallowest and middle depth for the single standard deviation range size (Figure 3.4 A) and to either side of the species richness peak in the middle depth for the two standard deviation range size (Figure 3.5 A). The R^2 value of the observed versus the expected for the single standard deviation was $R^2 = 0.5949$ ($p < 0.001$) and for the two standard deviation depth range $R^2 = 0.6667$ ($p < 0.001$). Both degrees of restriction on the species depth ranges resulted in more than 50% of the species richness being explained by 'geometric constraints, or in other words the mid-domain effect.

3.4.2 Body-size depth relationship

Body size (length) and depth relationships between the 30 species were found to be non significant ($p = 0.400$) (Figure 3.6 A). Both the Osteichthyes (Figure 3.6 B) and the Chondrichthyes (Figure 3.6 C) exhibited a non significant relationship between body size and depth. On an individual species basis 15 out of the 30 species plotted showed a significant positive relationship between body size and depth, six showed a significant negative relationship and nine showed no relationship (Figure 3.7). Osteichthyes predominately had species with a positive relationship while Chondrichthyes predominately had no relationship (Table 3.1).

Figure 3.4: (A) Observed species richness (circles) over the depth domain, plotted with mean expected species richness (solid line) as predicted by the RangeModel, and the upper and lower 95% confidence intervals (broken lines) for species with ranges set at one standard deviation of the observed depth ranges. (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel for species with depth ranges set at one standard deviation of observed depth ranges. $R^2 = 0.6667$ ($p < 0.001$).

Figure 3.5: (A) Observed species richness (circles) over the depth domain, plotted with mean expected species richness (solid line) as predicted by the RangeModel, and the upper and lower 95% confidence intervals (broken lines) for species with ranges set at two standard deviations of the observed depth ranges. (B) Observed species richness plotted against expected species richness, as predicted by the RangeModel for species with depth ranges set at two standard deviations of observed depth ranges. $R^2 = 0.5949$ ($p < 0.001$).

MDE & Body-Size Depth Distribution – Results

Figure 3.6: Interspecific relationship of average body size and average depth caught at for (A) all 30 species studied, (B) Osteichthyes only, and (C) Chondrichthyes only.

Table 3.1: Number of species, total and split by Class, and the type of body-size depth relationship shown. Significance at 0.05 level.

Relationship	Osteichthyes	Chondrichthyes	Total
Positive	12	3	15
Negative	4	2	6
No relationship	5	4	9
Sum	21	9	30

MDE & Body-Size Depth Distribution – Results

Figure 3.7: Intraspecific relationship of body size and depth caught at for each of the 23 species studied. For ease of interpretation the regression lines are shown over the entire length range, but were only fitted over the length and depth of the actual species data. No regression line was plotted for non-significant results. Species codes are given in APPENDIX 3.1.

3.5 DISCUSSION

I found moderately strong empirical support for the hypothesis that the species richness gradient of abundant demersal fish, between 200-900m on the Chatham Rise, New Zealand, is determined by ‘geometric constraints, indicating a mid-domain effect. Other studies on marine fish (Kendall and Haedrich 2006, Smith and Brown 2002) have found little or no evidence of the mid-domain effect on species distribution gradients. Kendall and Haedrich (2006) observed patterns that showed a decline in species richness with depth and did not match the richness patterns produced by the null model (the mid-domain effect). Applied to bathymetric ranges, Rapoport’s rule predicts that species richness decreases and range sizes increase with depth and latitude. Kendall and Haedrich (2006) found the rule explained the decrease in fish species richness with depth and between latitudes, but did not appear to explain the increase in range sizes with depth. Smith and Brown(2002) studied the species distribution of pelagic fish in the North East Pacific Ocean and found a very nested species assemblage and no evidence of the mid-domain effect. They suggested the inverse relationship between pelagic fish species diversity and depth supported the hypotheses that implicate temperature, productivity or both as causal mechanisms.

Considering that this study uses a subset of the depth range that Smith and Brown (2002) studied, at the shallow end of the scale, based on their findings I would expect to find a strongly left skewed distribution of species richness, however, this was not the case, as the species distribution I observed is right skewed. When sampling intensity was taken into account species richness still exhibited a peak towards the middle of the domain. It was not skewed to the right, like the plot for the unadjusted depth ranges, but rather more centred. This, while exhibiting more similarity to the findings of Smith and Brown (2002) than the unadjusted depths, still has greater than 50% of the species richness explained by the mid-domain effect. Given that the distribution of species richness of the unadjusted depths is right skewed, one might wonder if the 1000m domain boundary is too shallow and is closer to the domain mid-point and more sampling to a deeper depth would have

provided a more thorough investigation of the species present and possibly a more centred curve of species richness.

Methodological constraints on the definition of a hard boundary according to Colwell and Hurr (1994), as one of the conditions of a mid-domain effect model, is ‘a point beyond which species ranged can not extend’. Although mid-domain effect models have now been loosened to more ‘permeable’ or ‘soft’ boundaries (Connolly 2005), mid-domain effect curves are none the less sensitive to their placement, so cautions must be exercised in defining domain limits (Carranza et al. 2008). The depth limits observed here would be considered ‘soft’ domain boundaries as they are limited to 1000m due to the sampling restrictions of the survey data was collected on, but records for some species indicate that their bathymetric distributions may reach >1000m as well as < 200m. This discrepancy may be seen as artificially biasing the bathymetric range so that the mid-point lies in either a shallower or deeper position than it naturally occurs. McClain et al. (2007) lists three approaches commonly used for dealing with species whose ranges are only partially contained within the domain being analysed. These are: (1) exclude them completely from the analysis (Colwell and Lees 2000, Jetz and Rahbek 2001), (2) cut their ranges at the domain edge and treat the part of the range occurring within the domain as the entire range (used by Diniz-Filho et al. 2002, McCain 2003), and (3) cut the species range at the domain edge and force these range fragments to remain attached to the domain edge during analysis (Colwell et al. 2004). We followed the second approach.

Rare species ranges can be under estimated as occurrences at the shallow and deep edges of their ranges may not be captured. A species range is considered to be between the deepest and shallowest occurrences of that species within the sampling regime. The sampling methods used also limits the species captured due to gear selectivity. There is a minimum size of fish able to be caught due to trawl mesh size, which in this study is ~10cm. Sampling also only occurs within 7m of the sea floor (Stevens and O'Driscoll 2006, 2007, Stevens et al. 2008) due to the head line height of the net used, therefore restricting the study to comparatively large, abundant and demersal species. Different capture methods used by different studies will have different species selectivity and

catchability. For example, a mid water trawl with fine cod end mesh will capture a different species assemblage to that from a bottom trawl with large cod end mesh at the same depth (Stevens and O'Driscoll 2006, 2007, Stevens et al. 2008). Caution is needed when comparing species richness at particular depths between studies, to ensure that they are truly comparable. What is considered a rare species may also be a result of the sampling method or gear used, as some gear does not sample some, for example small, species very well. As a precaution only abundant species were used in this study to ensure even sampling of all species across all depths.

This study found no trend between body size and depth at the community level when looking at all species, as well as when the data were split by class into Osteichthyes and Chondrichthyes. This finding supports a neutral theory of species distribution. On a species level, however, both positive, negative and no trends were found between increasing body size with increasing depth. At a species level environmental or evolutionary factors may be affecting the species depth distribution. In the study by Smith and Brown (2002) they found depth differences between the Chondrichthyes and Osteichthyes in their size depth distributions, as well as significant trends in their interspecific distributions. Whether this is due to them having a larger sample size or the effect not being present on the Chatham Rise is still open for debate. The entire array of species from the Chatham Rise was not used and this may affect the significance of any trend found. The Chatham Rise, and New Zealand, does not have the massive manta rays or whale sharks that were present in Smith and Brown's (2002) study, which may facilitate the result they found. They suggest the negative relationship within the Chondrichthyes to be in part related to food sources, with the larger species being predominantly planktivorous and therefore limited to the comparatively shallow waters of the productive photic zone. It was also suggested that the result of internal fertilisation and increased maternal investment in Chondrichthyes, through either live birth or hatching, gives young an early advantage in life (Carrier et al. 2004b) and therefore they are better able to exist in the nutrient poor deeper water than their Osteichthyes counterparts. However finding only two of the Chondrichthyes species studied had negative length depth relationships food and reproduction may only be part of the story. Collins et

al (2005) also found differences in the depth distributions of Chondrichthyes and Osteichthyes, they conclude that something, though unclear, in the wide anatomical, behavioural and physiological differences between the two classes is the cause for the differences in depth distributions found. This suggests that on the basis of their differences both the Chondrichthyes and Osteichthyes are exploiting different niches.

Temperature changes with depth, and may explain some of the pattern seen in fish distribution. Small fish, within a species, are often found in shallower water than larger individuals of the same species (Collins et al. 2005, Macpherson and Duarte 1991). Water temperature affects fish metabolism, with warmer water being conducive to greater growth rates and cooler water slowing metabolic rates (Brill et al. 1999, Sogard and Olla 1996). Reduced metabolic costs implies that a greater fraction of the resources (food) consumed by an individual can be allocated to reproduction, therefore potentially increasing an individuals reproductive output. During ontogeny there are also physiological changes in fish, allowing them to tolerate colder temperatures and exploit the deeper regions that were previously unavailable to them (Brill et al. 1999, Swain 1997). This argument does not hold, however, for the species that display a negative size depth relationship. Instead it may be that the energy requirements for the larger fish is too great to be supported by the limited food resources, therefore requiring them to inhabit the shallower more nutrient rich waters. Smaller individuals, with their lower energetic needs may be better suited to inhabiting the deeper waters (Collins et al. 2005).

Overall it is difficult to determine one factor that is influencing the size distribution of fish with depth. Different species have different mechanism driving their distributions, and what may drive one species, may have little effect on another species. Testing the effects of hydrostatic pressure, light, temperature and food availability at different depths on changes in body size is difficult, as all variables are intertwined and difficult to tease apart. We found evidence of a mid-domain effect in the species distribution of the 30 most abundant demersal species on the Chatham Rise, between 200 – 1000m. This was supported by the lack of trends found between body size and depth in the total species

MDE & Body-Size Depth Distribution – Discussion

assemblage and when split by class into Osteichthyes and Chondrichthyes. Competition and adaption are also implied to work more strongly at a population level rather than at a community level as body-size depth trends were found in the majority of the species studied, although the nature of the relationship, be it positive or negative for body size increases with depth, differed between the individual species.

3.6 LITERATURE CITED

Annala, J.H., Sullivan, K.J., Smith, N.W.M., Griffiths, M.H., Todd, P.R., Mace, P.M., and Connell, A.M. 2004. Report from the Fisheries Assessment Plenary, May 2004: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 690.

Bone, Q., and Moore, R.H. 2008. Biology of fishes. Taylor & Francis Group, New York.

Bradford-Grieve, J., Boyd, P., Chang, F., Chiswell, S., Hadfield, M., Hall, J., James, M., Nodder, S., and Shushkina, E. 1999. Pelagic ecosystem structure and functioning in the subtropical front region east of New Zealand in austral winter and spring 1993. *Journal of Plankton Research* **21**(3): 405-428.

Bradford-Grieve, J.M., Chang, F.H., Gall, M., Pickmere, S., and Richards, F. 1997. Size-fractionated phytoplankton standing stocks and primary production during austral winter and spring 1993 in the Subtropical Convergence region near New Zealand. *New Zealand Journal of Marine and Freshwater Research* **31**: 201-224.

Brehm, G., Colwell, R.K., and Kluge, J. 2007. The role of environment and mid-domain effect on moth species richness along a tropical elevational gradient. *Global Ecology and Biogeography* **16**(2): 205-219.

Brill, R.W., Block, B.A., Boggs, C.H., Bigelow, K.A., Freund, E.V., and Marcinek, D.J. 1999. Horizontal movements and depth distribution of large adult yellowfin tuna (*Thunnus albacares*) near the Hawaiian Islands, recorded using ultrasonic telemetry: implications for the physiological ecology of pelagic fishes. *Marine Biology* **133**(3): 395-408.

Brown, J.H. 2001. Mammals on mountainsides: elevational patterns of diversity. *Global Ecology & Biogeography* **10**(1): 101-109.

MDE & Body-Size Depth Distribution – Literature Cited

- Bull, B., Bagley, N.W., and Hurst, R.J. 2000. Proposed survey design for the Southern Plateau trawl survey of hoki, hake and ling in November-December 2000. Final Research Report to the Ministry of Fisheries for Project MDT199/01 Objective 1.(Unpublished report held by Ministry of Fisheries, Wellington): 31.
- Carranza, A., Colwell, R.K., and Rangel, T. 2008. Distribution of megabenthic gastropods along environmental gradients: the mid-domain effect and beyond. *Marine Ecology Progress Series* **367**: 193-202.
- Carrier, J.C., Musick, J.A., and Heithaus, M.R. 2004a. Biology of sharks and their relatives. CRC Press, Boca Raton.
- Carrier, J.C., Pratt, H.L., Jr., and Castro, J.I. 2004b. Reproductive Biology of Elasmobranchs. *In* Biology of Sharks and their Relatives. *Edited by* J.C. Carrier, J.A. Musick and M.R. Heithaus. CRC Press, Boca Raton. p. 596.
- Chave, J. 2004. Neutral theory and community ecology. *Ecology Letters* **7**(3): 241-253.
- Collins, M.A., Bailey, D.M., Ruxton, G.D., and Priede, I.G. 2005. Trends in body size across an environmental gradient: A differential response in scavenging and non-scavenging demersal deep-sea fish. *Proceedings of the Royal Society B: Biological Sciences* **272**(1576): 2051-2057.
- Colwell, R.K. 2006. RangeModel A Monte Carlo simulation tool for assessing geometric constraints on species richness. Version 5. User's Guide and application published at: <http://viceroy.eeb.uconn.edu/rangemodel>.
- Colwell, R.K., and Hurtt, G.C. 1994. Nonbiological Gradients in Species Richness and a Spurious Rapoport Effect. *The American Naturalist* **144**(4): 570-595.
- Colwell, R.K., and Lees, D.C. 2000. The mid-domain effect: geometric constraints on the geography of species richness. *Trends in Ecology and Evolution* **15**(2): 70-76.
- Colwell, R.K., Rahbek, C., and Gotelli, N.J. 2004. The mid-domain effect and species richness patterns: what have we learned so far. *The American Naturalist* **163**(3).

MDE & Body-Size Depth Distribution – Literature Cited

- Colwell, R.K., Rahbek, C., and Gotelli, N.J. 2005. The mid-domain effect: there's a baby in the bathwater. *The American Naturalist* **166**(5): 149-154.
- Connolly, Sean R. 2005. Process-Based Models of Species Distributions and the Mid-Domain Effect. *The American Naturalist* **166**(1): 1-11.
- Cushing, D.H. 1975. *Marine ecology and fisheries*. Cambridge University Press.
- Cushing, D.H., and Walsh, J.J. 1976. *The ecology of the seas*. Blackwell.
- Diniz-Filho, J.A.F., de Sant'Ana, C.E.R., de Souza, M.C., and Rangel, T.F. 2002. Null models and spatial patterns of species richness in South American birds of prey. *Ecology Letters* **5**(1): 47-55.
- Dornelas, M., Connolly, S.R., and Hughes, T.P. 2006. Coral reef diversity refutes the neutral theory of biodiversity. *Nature* **440**(7080): 80-82.
- Dunn, R.R., McCain, C.M., and Sanders, N.J. 2007. When does diversity fit null model predictions? Scale and range size mediate the mid-domain effect. *Global Ecology & Biogeography* **16**(3): 305-312.
- Francis, R.I.C.C. 1984. An adaptive strategy for stratified random trawl surveys. *New Zealand Journal of Marine and Freshwater Research* **18**: 59-71.
- Hawkins, B.A., and Diniz-Filho, J.A.F. 2002. The mid-domain effect cannot explain the diversity gradient of Nearctic birds. *Global Ecology & Biogeography* **11**(5): 419-426.
- Hawkins, B.A., Diniz-Filho, J.A.F., and Weis, A.E. 2005. The mid-domain effect and diversity gradients: is there anything to learn? *The American Naturalist* **166**(5): 140-143.
- Helfman, G.S. 1978. Patterns of community structure in fishes: summary and overview. *Environmental Biology of Fishes* **3**(1): 129-148.
- Hubbell, S.P. 1979. Tree Dispersion, Abundance, and Diversity in a Tropical Dry Forest. *Science* **203**(4387): 1299-1309.

MDE & Body-Size Depth Distribution – Literature Cited

- Hubbell, S.P. 2001. The Unified Neutral Theory of Biodiversity and Biogeography. Princeton University Press.
- Hubbell, S.P. 2003. Modes of speciation and the lifespans of species under neutrality: a response to the comment of Robert E. Ricklefs. *Oikos* **100**(1): 193-199.
- Hubbell, S.P. 2005. Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional Ecology* **19**(1): 166-172.
- Hurst, R., Bagley, N.W., Chatterton, T., Hanchet, S., Schofield, K.A., and Vignaux, M. 1992. Standardisation of hoki/middle depth time series trawl surveys. MAF Fisheries Greta Point Internal Report **194**(Unpublished report held in NIWA library, Wellington): 89.
- Hurst, R.J., and Bagley, N.W. 1994. Trawl survey of middle depth and inshore bottom species off Southland, February-March 1993 (TAN9301). New Zealand Fisheries Data Report **52**: 58.
- Jetz, W., and Rahbek, C. 2001. Geometric constraints explain much of the species richness pattern in African birds. *Proceedings of the National Academy of Sciences* **98**(10): 5661.
- Keast, A. 1978. Trophic and spatial interrelationships in the fish species of an Ontario temperate lake. *Environmental Biology of Fishes* **3**(1): 7-31.
- Kendall, V.J., and Haedrich, R.L. 2006. Species richness in Atlantic deep-sea fishes assessed in terms of the mid-domain effect and Rapoport's rule. *Deep Sea Research Part I: Oceanographic Research Papers* **53**(3): 506-515.
- Krebs, C.J. 2001. *Ecology: the Experimental Analysis of Distribution and Abundance*. Benjamin Cummings, San Francisco.
- Lees, D.C., and Colwell, R.K. 2007. A strong Madagascan rainforest MDE and no equatorward increase in species richness: re-analysis of The missing Madagascan mid-

MDE & Body-Size Depth Distribution – Literature Cited

domain effect, by Kerr JT, Perring M. Currie DJ (Ecology Letters 9: 149159, 2006). Ecology Letters **10**(9): E4-E8.

Lees, D.C., Kremen, C., and Andriamampianina, L. 1999. A null model for species richness gradients: bounded range overlap of butterflies and other rainforest endemics in Madagascar. Biological Journal of the Linnean Society **67**(4): 529-584.

Leibold, M.A., and McPeck, M.A. 2006. Coexistence of the niche and neutral perspectives in community ecology. Ecology **87**(6): 1399-1410.

Macpherson, E. 1981. Resource partitioning in a Mediterranean demersal fish community. Marine Ecology Progress Series **4**: 183-193.

Macpherson, E., and Duarte, C.M. 1991. Bathymetric trends in demersal fish size: Is there a general relationship? Marine Ecology Progress Series **71**(2): 103-112.

Macpherson, E., and Duarte, C.M. 1994. Patterns in species richness, size, and latitudinal range of east Atlantic fishes. Ecography **17**(3): 242-248.

McCain, C.M. 2003. North American desert rodents: A test of the mid-domain effect in species richness. Journal of Mammalogy **84**(3): 967.

McCain, C.M. 2004. The mid-domain effect applied to elevational gradients: species richness of small mammals in Costa Rica. Journal of Biogeography (J. Biogeogr.) **31**: 19-31.

McClain, C.R., White, E.P., and Hurlbert, A.H. 2007. Challenges in the application of geometric constraint models. Global Ecology & Biogeography **16**(3): 257-264.

McClatchie, S., and Dunford, A. 2003. Estimated biomass of vertically migrating mesopelagic fish off New Zealand. Deep Sea Research Part I: Oceanographic Research Papers **50**(10-11): 1263-1281.

MDE & Body-Size Depth Distribution – Literature Cited

- McClatchie, S., Macaulay, G.J., and Coombs, R.F. 2004. Acoustic backscatter and copepod secondary production across the Subtropical Front to the east of New Zealand. *Journal of Geophysical Research* **109**.
- McClatchie, S., Millar, R.B., Webster, F., Lester, P.J., Hurst, R., and Bagley, N. 1997. Demersal fish community diversity off New Zealand: Is it related to depth, latitude and regional surface phytoplankton? *Deep Sea Research Part I: Oceanographic Research Papers* **44**(4): 647-667.
- Murphy, R.J., Pinkerton, M.H., Richardson, K.M., Bradford-Grieve, J.M., and Boyd, P.W. 2001. Phytoplankton distributions around New Zealand derived from SeaWiFS remotely-sensed ocean colour data. *New Zealand Journal of Marine and Freshwater Research* **35**: 343-362.
- Nybakken, J.W. 2001. *Marine biology : an ecological approach*. Benjamin Cummings, San Francisco.
- Nybakken, J.W., and Bertness, M.D. 2005. *Marine biology: An ecological approach*. Benjamin Cummings, San Francisco.
- Olabarria, C., and Thurston, M.H. 2003. Latitudinal and bathymetric trends in body size of the deep-sea gastropod *Troschelia berniciensis* (King). *Marine Biology* **143**(4): 723-730.
- Perry, A.L., Low, P.J., Ellis, J.R., and Reynolds, J.D. 2005. Climate change and distribution shifts in marine fishes. *American Association for the Advancement of Science*. pp. 1912-1915.
- Pineda, J., and Caswell, H. 1998. Bathymetric species-diversity patterns and boundary constraints on vertical range distributions. *Deep Sea Research Part II: Topical Studies in Oceanography* **45**(1-3): 83-101.
- Rahbek, C., Gotelli, N.J., Colwell, R.K., Entsminger, G.L., Rangel, T., and Graves, G.R. 2007. Predicting continental-scale patterns of bird species richness with spatially explicit models. *Proceedings of the Royal Society B: Biological Sciences* **274**(1607): 165-174.

MDE & Body-Size Depth Distribution – Literature Cited

- Rex, M.A., and Etter, R.J. 1998. Bathymetric patterns of body size: implications for deep-sea biodiversity. *Deep Sea Research Part II: Topical Studies in Oceanography* **45**(1): 103-127.
- Rohde, K. 1992. Latitudinal Gradients in Species Diversity: The Search for the Primary Cause. *Oikos* **65**(3): 514-527.
- Rohde, K., Heap, M., and Heap, D. 1993. Rapoport's Rule Does Not Apply to Marine Teleosts and Cannot Explain Latitudinal Gradients in Species Richness. *The American Naturalist* **142**(1): 1.
- Romdal, T.S., Colwell, R.K., and Rahbek, C. 2005. The influence of band sum area, domain extent, and range sizes on the latitudinal mid-domain effect. *Ecology* **86**(1): 235-244.
- Sarda, F., and Cartes, J.E. 1993. Relationship between size and depth in decapod crustacean populations on the deep slope in the Western Mediterranean. *Deep Sea Research Part I: Oceanographic Research Papers* **40**(11): 2389-2400.
- Smith, K.F., and Brown, J.H. 2002. Patterns of diversity, depth range and body size among pelagic fishes along a gradient of depth. *Global Ecology & Biogeography* **11**(4): 313-322.
- Sogard, S.M., and Olla, B.L. 1996. Food deprivation affects vertical distribution and activity of a marine fish in a thermal gradient: potential energy-conserving mechanisms. *Marine ecology progress series. Oldendorf* **133**(1): 43-44.
- SPSS. 2007. SPSS Statistics Version 16.0.1 for windows. SPSS Inc.
- Stevens, D.W., and O'Driscoll, R.L. 2006. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2005 (TAN0501). *New Zealand Fisheries Assessment Report* **13**.

MDE & Body-Size Depth Distribution – Literature Cited

- Stevens, D.W., and O'Driscoll, R.L. 2007. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2006 (TAN0601). New Zealand Fisheries Assessment Report **5**.
- Stevens, D.W., O'Driscoll, R.L., and Gauthier, S. 2008. Trawl survey of hoki and middle depth species on the Chatham Rise, January 2007 (TAN0701). New Zealand Fisheries Assessment Report **52**.
- Stevens, G.C. 1989. The Latitudinal Gradient in Geographical Range: How so Many Species Coexist in the Tropics. *The American Naturalist* **133**(2): 240-256.
- Sutton, P. 2001. Detailed structure of the Subtropical Front over Chatham Rise, east of New Zealand. *Journal of Geophysical Research. C. Oceans* **106**: 31045-31056.
- Swain, D.P. 1997. Sex-specific temperature distribution of American plaice (*Hippoglossoides platessoides*) and its relation to age and abundance. *Canadian Journal of Fisheries and Aquatic Sciences* **54**(5): 1077-1087.
- Uiblein, F., Bordes, F., and Castillo, R. 1996. Diversity, abundance and depth distribution of demersal deep-water fishes off Lanzarote and Fuerteventura, Canary Islands. *Journal of Fish Biology* **49**: 75-90.
- Whitfield, J. 2002. Neutrality versus the niche. *Nature* **417**(6888): 480-481.
- Willig, M.R., and Lyons, S.K. 1998. An Analytical Model of Latitudinal Gradients of Species Richness with an Empirical Test for Marsupials and Bats in the New World. *Oikos* **81**(1): 93-98.
- Zapata, F.A., Gaston, K.J., and Chown, S.L. 2005. The mid-domain effect revisited. *The American Naturalist* **166**: 144-148.

APPENDIX 3.1: Species codes.

List of species codes used for each of the individual species studied, common name and scientific name.

Code	Common name	Scientific name
BAR	Barracouta	<i>Thyristes atun</i>
BBE	Banded Bellowsfish	<i>Centriscops humerosus</i>
BOE	Black Oreo	<i>Allocyttus niger</i>
BYS	Alfonsino	<i>Beryx splendens</i>
CAS	Oblique Banded Rattail	<i>Caelorinchus aspercephalus</i>
CBO	Bollons Rattail	<i>Caelorinchus bollonsi</i>
COL	Oliver's Rattail	<i>Caelorinchus oliverianus</i>
CSQ	Centrophorus squamosus	<i>Centrophorus squamosus</i>
CYP	Cetnroscymnus crepidater	<i>Cetnroscymnus crepidater</i>
ETB	Baxter's Lantern Dogfish	<i>Etmopterus baxteri</i>
GSH	Dark Ghost Shark	<i>Hydrolagus novaezelandiae</i>
GSP	Pale Ghost Shark	<i>Hydrolagus bemisi</i>
HAK	Hake	<i>Merluccius australis</i>
HOK	Hoki	<i>Macruronus novaezelandiae</i>
JAV	Javelin fish	<i>Lepidorhynchus denticulatus</i>
LCH	Long-nosed Chimera	<i>Harriotta raleighana</i>
LDO	Lookdown Dory	<i>Cyttus traverse</i>
LIN	Ling	<i>Genypterus blacodes</i>
OPE	Orange Perch	<i>Lepidoperca aurantia</i>
RCO	Red Cod	<i>Pseudophycis bachus</i>
SOR	Spiky Oreo	<i>Pseudocyttus maculatus</i>
SND	Shovelnose Spiny Dogfish	<i>Deania calcea</i>
SPD	Spiny Dogfish	<i>Squalus acanthias</i>
SPE	Sea Perch	<i>Helicolenus spp.</i>
SRB	Southern Ray's Bream	<i>Brama spp.</i>
SSO	Smooth Oreo	<i>Neocyttus rhomboidalis</i>
SSK	Smooth Skate	<i>Dipturus innominatus</i>
STA	Giant Stargazer	<i>Kathetostoma giganteum</i>
SWA	Silver Warehou	<i>Seriolella punctata</i>
WWA	White Warehou	<i>Seriolella caerulea</i>

4 GENERAL DISCUSSION

The Chatham Rise is a highly productive environment with high primary productivity facilitated by its location at the meeting point of two large water masses called the Subtropical convergence (STC) (Bradford-Grieve et al. 1996). This environment of high primary productivity is believed to be the underlying cause for the regions to support a number of commercial fisheries (Annala et al. 2004). These fisheries are all managed on a single species basis, as are most New Zealand's fisheries. Globally there is a trend moving away from single species fisheries management toward a more holistic ecosystem based management approach (Francis et al. 2007, Marasco et al. 2007). Little is known about the species interactions of the commercially exploited species and the trophic structure of the community on the Chatham Rise (Annala et al. 2004, Ministry of Fisheries 2007, 2008). Diet studies are a good method of investigating trophic interactions in marine communities (Link 1999).

The first goal of this study was to investigate the effects of several sources of variation in the feeding ecology of hoki (*Macruronus novaezelandiae*). The specific objectives were to investigate the distribution of hoki on the Chatham Rise by size, depth, and sex; characterise the diet of hoki on the Chatham Rise, New Zealand, between 200-800m; and test for the effects of fish size, depth, location and sex on diet variability of hoki.

The second goal of this study was to test for evidence of the mid-domain effect on 30 of the abundant demersal fish species associated with hoki on the Chatham Rise, as well as look at body-size depth relationships between the 30 species studied, within class Osteichthyes (bony fish, e.g. barracouta) and Chondrichthyes (cartilaginous fish, e.g. sharks) and on an individual species basis.

4.1 Hoki diet

The diet of hoki (*Macruronus novaezelandiae*) was studied through the identification of prey and analysis of stomach contents. The samples were collected during January of 2005, 2006 and 2007 creating a snapshot of the diet, rather than a picture that included the full potential seasonal variability. Hoki was found to have a diet of predominately pelagic origins. Their diet contained a range of teleosts, largely from the family Myctophidae, natant decapods and euphausiids, as well as some cephalopods (largely squid). These findings are similar to historical studies by Kerstan and Sahrhage (1980), Kuo and Tanaka (1984), Clark (1985), and Bulman and Blaber (1986), however some differences were also found. Clark (1985) found amphipods to feature highly in the diet of hoki in the Subantarctic. This presence of amphipods was not found in the current study. Speculation on the reason for this difference may lie in the distribution of amphipods which may have been in high abundance in the Subantarctic during the sampling period of Clark (1985), while they may not have been abundant on the Chatham Rise. Lack of knowledge on amphipod distribution prevents this hypothesis from being tested. Alternatively there may be a shift in diet as fish migrate from the Chatham Rise to the Subantarctic. No evidence of cannibalism was found in this study, unlike Kuo and Tanaka (1984), and Bulman and Blaber (1986). This lack of cannibalism may be due to the physical separation between large and small hoki, with small fish found in shallow water while large fish were found in deeper water. Alternatively the small fish may have reached a size where they were too big to be predated on by the large hoki present on the Chatham Rise. Due to sampling restrictions, trawl mesh size, hoki of less than about 30cm total length were not caught as part of the sampling for this study. From these surveys, it isn't known whether hoki smaller than about 30cm were present on the Chatham Rise.

Differences in depth were found to explain the most variability in hoki diet. Fish found in deeper water were found to eat different prey from fish found in shallower water. This difference in diet due to depth was still found when fish size was held constant. Fish size also explained a large amount of diet variability, although only as a gradual change through ontogeny rather than a dramatic diet shift as has been found in other fish species

General Discussion

(Garicía-Berthou and Moreno-Amich 2000). Only a small amount of variation in diet was found between the four areas, North West, North East, South West and South East, and no consistent pattern when depth and size were held constant. The differences between the areas may be aliasing both size and depth distributions. Hoki are not evenly distributed over the Chatham Rise, and over the depth range samples. Small fish were found predominately on the Western end of the rise and shallower than 500m.

Hoki displayed a significant relationship between body size and depth, with body size increasing with increasing depth. This change in depth distribution with size was also reflected in a change in dietary composition. Larger fish are physically able to take larger prey, due to gape size increasing, and therefore are able to exploit a different range of species, and therefore niche, than smaller fish. Given that very little is known about prey distribution it is difficult to determine whether the cause for diet differences is because of the differences in depth, fish size, or prey patchiness, or a combination of all three.

4.2 Body-size depth relationship

At a general community level no body-size depth relationship was found between the 30 species studied. When split by class Chondrichthyes and Osteichthyes, no trend was also found. On an individual species basis positive, negative and no trends were found in body-size depth relationships. What drives these trends is still unclear although there are some potential explanations. Temperature decreases with depth and in some species small fish are physiologically unable to live at low temperatures (Brill et al. 1999). Warmer water is more conducive to greater growth rates, while cooler water slows metabolic rates (Brill et al. 1999, Sogard and Olla 1996). Taking advantage of these different effects at different life stages may be beneficial. When fish are small their main life objective is to reach the size of sexual maturity, therefore putting all energy into growth. Once sexual maturity is reached their main objective is to direct all energy into breeding, therefore living in an environment that reduces metabolic cost would be an advantage. This argument doesn't hold true for species that have a negative body-size depth relationship. Instead it may be that the energy requirements of the larger fish are too great to be

General Discussion

supported by the limited food resources, therefore requiring them to inhabit the shallower more nutrient rich waters. Smaller individuals, with their lower energetic needs may be better suited to inhabiting the deeper waters (Collins et al. 2005).

The differences in body-size depth relationships between the Chondrichthyes and Osteichthyes at a species level have been hypothesised to be due largely to differences in reproductive strategies. Chondrichthyes have internal fertilisation and give birth to live young, with no larval stage (Carrier et al. 2004a). Most Osteichthyes lay or release eggs and their young develop through a larval stage which is vulnerable to predation (Bond 1996). It is the result of internal fertilisation and increased maternal investment in Chondrichthyes, through either live birth or hatching, that gives young an early advantage in life (Carrier et al. 2004b) and therefore they are better able to exist in the nutrient poor deeper water than their Osteichthyes counter parts.

Hoki exhibited differences in diet that reflected the differences in size distribution found, with small fish found in shallower water and larger fish found in deeper water. From this result it could be inferred that the pattern of diet differences between different sized fish might also be evident in other species that showed body-size depth trends. To validate this inference investigation into the diet of these species would be required.

Predator species have different distributions with depth, both within the specific species, as found from the body-size depth investigation of this study, and between the species, with some species preferring shallower or deeper water to others, seen from the different limits to of the species ranges. It would be logical to question prey distribution and hypothesise that similar patterns of distribution may be displayed. Not a lot is known about the distribution of prey species. Robertson et al. (1978) studied the distribution of myctophids over the subtropical convergence, over the Chatham Rise, and found subtropical species in higher abundance on the northern side while subantarctic species were in higher abundance on the southern side. Robertson et al. (1978) also found differences in the species distribution of prawns over the subtropical convergence, with some species predominately found in the warmer waters to the north and other species

General Discussion

found in the colder waters to the south. Robertson et al.'s (1978) study looked at the spatial distribution of species down to 400m depth, but didn't look at the potential differences in species composition with depth. It is known that there is a large amount of vertical migration by mesopelagic fish species (McClatchie and Dunford 2003), and this vertical migration may confound any body-size depth relationship, as all sizes may vertically migrate between similar depths. As with predator species there may be a change in the composition of species found with increasing depth. Anecdotal evidence from midwater research trawls for mesopelagic reference materials for dietary identification found that one species of Myctophid (*Lampanyctodes hectoris*) and one Sternoptychidae (*Maurolicus muelleri*) were very abundant in shallow tows (100-200m) and that their abundance decreased as tow depth increased. With the tow depth increase so to did the abundance of other myctophid species. This change in species composition is also reflected to a degree in hoki diet, with larger bodied myctophids and other mesopelagic fish species featuring only in the diet of large hoki, and hoki from deep water.

The lack of body-size depth relationship found between the 30 species studied, and within the two classes, Chondrichthyes and Osteichthyes, supports a more neutral species distribution, rather than one explained by niche theory. On an individual species basis niche theory appears to better describe the distribution of individuals, as a strong relationship between body-size and depth distribution exists for some species.

4.3 Mid-domain effect

The mid-domain effect is the shuffling of species latitudinal ranges within the geometric constraints of a biogeographical domain, providing species ranges do not extend beyond the outer limits of the domain, which results in a pattern of species richness with the richness peak in the centre of the domain and reducing towards the domain boundaries (Colwell and Hurtt 1994, Willig and Lyons 1998). Moderately strong support was found for the hypothesis that the species richness gradient of abundant demersal fish in this study is determined by 'geometric constraints' giving evidence of the mid-domain effect.

General Discussion

This is in contrast to the findings of Smith and Brown (2002) in the Pacific ocean and Kendall and Haedrich (2006) in the North Atlantic. Kendall and Haedrich (2006) compare their findings against the mid-domain effect and state it's usefulness at eliminating the argument for a random distribution of species across the studied depth gradient.

McClatchie et al. (1997) looked at demersal fish community diversity off New Zealand and tried to relate it to depth, latitude and regional surface primary production. They did not test specifically for the mid-domain effect over depth, but found an increase in species richness with increasing depth (within the studies depth limit of 1000m). They also found a decrease in species richness with latitude between the Chatham Rise and the subantarctic. They hypothesised that the difference in species richness with between latitudes could be due to differences in primary productivity, being higher on the Chatham Rise, due to the presence of the subtropical convergence, than the subantactic. McClatchie et al. (1997) continued to note that caution is needed when drawing conclusions on the correlation between species richness and primary productivity, as benthic topographical relief (steepness of bottom slope) is also correlated with species richness. Rapid changes in relief bring about rapid changes in habitat type, probable overlap in fish communities, and associated hard bottom invertebrate communities that support corals, hydroids and sponges (McClatchie et al. 1997). These habitats are unlikely to occur in more sedimented low relief area (McClatchie et al. 1997). The subantarctic is a comparatively flat plateau with little steep relief, where as the Chatham Rise rises fairly rapidly from abyssal depths. McClatchie et al. (1997) gave no hypothesis to potentially explain the species richness gradient observed over depth.

This study used a sub-set of the depth range used by both Smith and Brown (2002) and Kendall and Haedrich (2006). Based on their finding it was expected that species richness would be skewed left (towards the shallower depths) however this was not found, as a peak in species richness was found towards the middle of the domain even when sampling bias was taken into account. One of the conditions of a mid-domain effect model is that a hard domain boundary is 'a point beyond which species ranges can not

extend' (Colwell and Hurtt 1994). Mid-domain models have been loosened to more 'permeable' or soft boundaries (Connolly 2005), allowing studies such as this one to include species known to have ranges that extend beyond the domain boundaries. This study follows the examples of Diniz-Filho et al. (2002) and McCain (2003) by truncating the species ranges, that extended beyond the domain boundaries, at the domain boundaries and including the reduced ranges in the analysis. Truncating the ranges may artificially bias the bathymetric ranges so that the range mid-point lies in either a shallower or deeper position than naturally occurs. Mid-domain curves are none the less sensitive to domain boundary placement, so caution must be exercised in defining domain limit (Carranza et al. 2008).

4.4 Conclusions

The Chatham Rise is a diverse ecosystem covering a broad range of depths. The study on hoki diet found a change in diet related to the different depths at which fish were caught. Differences in diet were also found between large and small fish. This difference between large and small fish was still found when depth was held constant, as was the difference between depths when size was constant. The findings in this study on diet characteristics are largely similar to historical studies in the same location. Studies from different locations also found evidence of a pelagic feeding strategy, however some difference in prey types was found between studies. The difference in prey types found may be due to differences in prey distribution, however very little is known about prey distribution patterns around New Zealand. A large difference between this study and some of the historical studies is the lack of cannibalism found on the Chatham Rise. Physical separation between small and large fish is hypothesised as a potential explanation.

Hubbell (2001, 2003, 2005) put forward the hypothesis of neutral theory, with random chance explaining species distribution. Using the mid-domain effect model evidence was found that the abundant, demersal fish assemblage on the Chatham Rise was randomly

General Discussion

distributed over depth. This was in contrast to two other studies which found nested species distribution and no evidence of the mid-domain effect.

When investigated at a species level the effects of adaption were evident in body-size depth relationships. Some species exhibited a positive trend, others a negative trend and some species had no relationship at all. This lends weight to the hypothesis that competition and predation work at a population level rather than at a broad community scale.

4.5 Future directions

This study highlighted the lack of knowledge on prey species distribution over the Chatham Rise. Further investigation into this would be beneficial for distinguishing whether differences in diet within a species is due to patchy prey distribution or another variable such as changes in body size or depth. The investigation of hoki diet is one step towards the data needed to create a food web on the Chatham Rise, and eventually a trophic model that can be used in fisheries management. The diet of 24 other commercial and common by-catch species on the Chatham Rise is being investigated by NIWA as part of a Ministry of Fisheries funded project to investigate the ecosystem-scale trophic relationships of middle-depth, demersal fish through diet composition and guild structure (Dunn et al. in prep).

A lot is known about the hydrodynamics and currents of the Chatham Rise, the amount of primary productivity, the commercial fishing effort, and commercial species distribution. Little, however, is known about the direct link between the primary productivity and commercial fish species. The diet part of this study looks at what prey is important to one commercial fish species (hoki). Unfortunately little is also known about the distribution of the prey species with depth and over the entire Chatham Rise area, and the link between the prey species and primary productivity. Further investigation into the link between prey species and primary productivity would potentially identify whether there is a link between primary productivity and hoki abundance in particular location.

General Discussion

Further research is needed to understand the distribution and abundance patterns of the prey of hoki. From there we can begin to understand the impact these prey, and their abundances in time and space, may have on the diet variability of hoki. Investigation in to hoki diet over time is needed to test for seasonal variation. Comparing the prey composition in hoki diet with local prey abundances and fluctuations over time would give information on whether hoki is actively selecting certain prey types or just eating the most abundant prey.

The location of hoki spawning aggregations have been identified and egg and larvae (less than 18mm total length) distribution have been noted in the vicinity of the spawning aggregation (Murdoch et al. 1990). The feeding of these larvae (less than 18mm total length) has been investigated by Murdoch (1990), who found them to eat copepods of various species. Little is known about the distribution and feeding of larvae and juvenile hoki, between 18mm total length and about 30cm total length, before they begin to be caught on the Chatham Rise.

Further investigation into mid-domain effect on species distribution with depth over the Chatham Rise would be interesting. Using a greater depth range as domain boundaries would make a more comparable study those in the North Atlantic by Kendall and Haedrich (2006) and the pelagic Pacific Ocean by Smith and Brown (2002). Increased sampling effort or the easing of conditions used for the classification of 'abundant' would allow for more species to be included in the analysis and it would be interesting to see whether the mid-domain effect still explained a large amount of the species richness gradient seen with depth.

4.6 LITERATURE CITED

Annala, J.H., Sullivan, K.J., Smith, N.W.M., Griffiths, M.H., Todd, P.R., Mace, P.M., and Connell, A.M. 2004. Report from the Fisheries Assessment Plenary, May 2004: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 690.

Bond, C.E. 1996. Biology of Fishes. Brooks/Cole, Melbourne.

Bradford-Grieve, J.M., Murdoch, R.C., James, M.R., Oliver, M., and Hall, J. 1996. Vertical distribution of zooplankton > 39 µm in relation to the physical environment off the west coast of South Island, New Zealand. New Zealand Journal of Marine and Freshwater Research **30**: 285-300.

Brill, R.W., Block, B.A., Boggs, C.H., Bigelow, K.A., Freund, E.V., and Marcinek, D.J. 1999. Horizontal movements and depth distribution of large adult yellowfin tuna (*Thunnus albacares*) near the Hawaiian Islands, recorded using ultrasonic telemetry: implications for the physiological ecology of pelagic fishes. Marine Biology **133**(3): 395-408.

Bulman, C.M., and Blaber, S.J.M. 1986. Feeding ecology of *Macruronus novaezelandiae* (Hector) (Teleostei: Merlucciidae) in south-east Australia. Australian Journal of Marine and Freshwater Research **37**(5): 621-639.

Carranza, A., Colwell, R.K., and Rangel, T. 2008. Distribution of megabenthic gastropods along environmental gradients: the mid-domain effect and beyond. Marine Ecology Progress Series **367**: 193-202.

Carrier, J.C., Musick, J.A., and Heithaus, M.R. 2004a. Biology of sharks and their relatives. CRC Press, Boca Raton.

General Discussion – Literature Cited

- Carrier, J.C., Pratt, H.L., Jr., and Castro, J.I. 2004b. Reproductive Biology of Elasmobranchs. *In* Biology of Sharks and their Relatives. *Edited by* J.C. Carrier, J.A. Musick and M.R. Heithaus. CRC Press, Boca Raton. p. 596.
- Clark, M. 1985. The food and feeding of seven fish species from the Campbell Plateau, New Zealand. *New Zealand Journal of Marine and Freshwater Research* **19**: 339-363.
- Collins, M.A., Bailey, D.M., Ruxton, G.D., and Priede, I.G. 2005. Trends in body size across an environmental gradient: A differential response in scavenging and non-scavenging demersal deep-sea fish. *Proceedings of the Royal Society B: Biological Sciences* **272**(1576): 2051-2057.
- Colwell, R.K., and Hurtt, G.C. 1994. Nonbiological Gradients in Species Richness and a Spurious Rapoport Effect. *The American Naturalist* **144**(4): 570-595.
- Connolly, Sean R. 2005. Process-Based Models of Species Distributions and the Mid-Domain Effect. *The American Naturalist* **166**(1): 1-11.
- Diniz-Filho, J.A.F., de Sant'Ana, C.E.R., de Souza, M.C., and Rangel, T.F. 2002. Null models and spatial patterns of species richness in South American birds of prey. *Ecology Letters* **5**(1): 47-55.
- Dunn, M., Connell, A.M., Forman, J., Griggs, L., Horn, P., Notman, P., Pinkerton, M., Stevens, D.W., and Wood, B. in prep. Ecosystem-scale trophic relationships: diet composition and guild structure of middle-depth fish on the Chatham Rise. Final Research Report to the Ministry of Fisheries for Project ZBD2004-02 Objectives 1-5.
- Francis, R.C., Hixon, M.A., Clarke, M.E., Murawski, S.A., and Ralston, S. 2007. Fisheries Management-Ten Commandments for Ecosystem-Based Fisheries Scientists. *Fisheries-American Fisheries Society* **32**(5): 217-233.
- Garicía-Berthou, E., and Moreno-Amich, R. 2000. Food of introduced pumpkinseed sunfish: ontogenetic diet shift and seasonal variation. *Journal of Fish Biology* **57**(1): 29-40.

General Discussion – Literature Cited

Hubbell, S.P. 2001. *The Unified Neutral Theory of Biodiversity and Biogeography*. Princeton University Press.

Hubbell, S.P. 2003. Modes of speciation and the lifespans of species under neutrality: a response to the comment of Robert E. Ricklefs. *Oikos* **100**(1): 193-199.

Hubbell, S.P. 2005. Neutral theory in community ecology and the hypothesis of functional equivalence. *Functional Ecology* **19**(1): 166-172.

Kendall, V.J., and Haedrich, R.L. 2006. Species richness in Atlantic deep-sea fishes assessed in terms of the mid-domain effect and Rapoport's rule. *Deep Sea Research Part I: Oceanographic Research Papers* **53**(3): 506-515.

Kerstan, M., and Sahrhage, D. 1980. Biological investigations on fish stocks in the waters off New Zealand. Bundesforschungsanstalt für Fischerei, Hamburg. *Mitteilungen aus dem Institut für Seefischerei*: 29.

Kuo, C., and Tanaka, S. 1984. Feeding habit of hoki *Macruronus novaezelandiae* (Hector) in waters around New Zealand. *Bulletin of the Japanese Society of Scientific Fisheries/Nissuishi* **50**(5): 783-786.

Link, J. 1999. Reconstructing food webs and managing fisheries. *In* *Proceedings of the 16th Lowell Wakefield Fisheries Symposium*. pp. 99-01.

Marasco, R.J., Goodman, D., Grimes, C.B., Lawson, P.W., Punt, A.E., and Quinn, T.J., II. 2007. Ecosystem-based fisheries management: some practical suggestions. *Canadian Journal of Fisheries and Aquatic Sciences* **64**(6): 928(912).

McCain, C.M. 2003. North American desert rodents: A test of the mid-domain effect in species richness. *Journal of Mammalogy* **84**(3): 967.

McClatchie, S., and Dunford, A. 2003. Estimated biomass of vertically migrating mesopelagic fish off New Zealand. *Deep Sea Research Part I: Oceanographic Research Papers* **50**(10-11): 1263-1281.

General Discussion – Literature Cited

McClatchie, S., Millar, R.B., Webster, F., Lester, P.J., Hurst, R., and Bagley, N. 1997. Demersal fish community diversity off New Zealand: Is it related to depth, latitude and regional surface phytoplankton? Deep Sea Research Part I: Oceanographic Research Papers **44**(4): 647-667.

Ministry of Fisheries. 2007. Report from the Fisheries Assessment Plenary, May 2007: stock assessment and yield estimates. (Unpublished report held in NIWA library, Wellington.): 344-370.

Ministry of Fisheries. 2008. Report from the Fisheries Assessment Plenary, May 2008: stock assessments and yield estimates. 347-371.

Murdoch, R.C. 1990. Diet of hoki larvae (*Marcuronus novaezelandiae*) off Westland, New Zealand. New Zealand Journal of Marine and Freshwater Research **24**: 518-527.

Murdoch, R.C., Guo, R., and McCrone, A. 1990. Distribution of hoki (*Macruronus novaezelandiae*) eggs and larvae in relation to hydrography in eastern Cook Strait, September 1987. New Zealand Journal of Marine and Freshwater Research **24**(4): 529-539.

Robertson, D.A., Roberts, P.E., and Wilson, J.B. 1978. Mesopelagic faunal transition across the Subtropical Convergence east of New Zealand. New Zealand Journal of Marine and Freshwater Research **12**: 295-312.

Smith, K.F., and Brown, J.H. 2002. Patterns of diversity, depth range and body size among pelagic fishes along a gradient of depth. Global Ecology & Biogeography **11**(4): 313-322.

Sogard, S.M., and Olla, B.L. 1996. Food deprivation affects vertical distribution and activity of a marine fish in a thermal gradient: potential energy-conserving mechanisms. Marine ecology progress series. Oldendorf **133**(1): 43-44.

Willig, M.R., and Lyons, S.K. 1998. An Analytical Model of Latitudinal Gradients of Species Richness with an Empirical Test for Marsupials and Bats in the New World. Oikos **81**(1): 93-98.