Molecular studies of New Zealand Fucales: Phylogeography, phylogeny and taxonomy in *Carpophyllum* and *Cystophora* (Phaeophyceae)

by

Joseph Buchanan

A thesis submitted to the Victoria University of
Wellington in fulfilment of the requirements for the
degree of Doctor of Philosophy in Marine Biology

Victoria

UNIVERSITY OF WELLINGTON

*Te Whare Wānanga
o te Ūpoko o te Ika a Māui*

School of Biological Sciences
Victoria University of Wellington
Te Whare Wānanga o te Ūpoko o te Ika a Māui

2011

*On the rocky shore of the island in the offing, the lovely seaweed – I
will miss it when it goes under the flowing tide.*

– Yamabe no Akahito
composed in the fifth of the tenth month, 724.

Abstract

Genetic variation in *Carpophyllum* Greville and *Cystophora* J. Agardh (Fucales, Phaeophyceae) was investigated at a variety of scales. An extensive survey of mitochondrial spacer variation in *Carpophyllum maschalocarpum* from 32 populations around New Zealand shows strong population differentiation at relatively small scales (50–100 kilometres), but also pathways of long distance dispersal that connect populations over much greater distances. In addition, historical climate change appears to have restricted *C. maschalocarpum* to the northern North Island during the last glacial maximum, with subsequent southward range expansion revealed by low genetic diversity in southern populations. These results are consistent with limited dispersal at the gamete and zygote stage, expected in fucalean algae, but with occasional long distance dispersal by detached floating thalli. The genetic signature suggests these two modes of dispersal are decoupled.

Internal Transcribed Spacers sequences show little differentiation between *C. maschalocarpum* and *C. angustifolium*, and hybridisation was found in several populations where these species are broadly sympatric. In the Bay of Plenty the two species had different ITS ribotypes, but most *C. angustifolium* specimens had a mitochondrial spacer haplotype that clustered with *C. maschalocarpum* haplotypes. This indicates mitochondrial introgression from *C. maschalocarpum* into *C. angustifolium*. In Northland species were difficult to separate by morphology or molecular markers, and some populations appear to be comprised entirely of hybrids.

Genetic distances between different species of *Cystophora* are very variable, and in some cases intra-species distances are similar to inter-species distances. This is problematic for DNA barcoding methods that rely on thresholds between inter-species and intra-species genetic distances. In some (but not all) cases, the absence of molecular differentiation can be attributed to oversplitting of *Cystophora* species by

morphological methods, and I synonymise *C. congesta* with *C. retroflexa*, and *C. distenta* with *C. scalaris*. These studies exemplify the difficulties of delimiting species in brown algae: Morphology is often misleading or uninformative; genetic differentiation of species is very variable and often low; and species' histories show complex patterns of isolation and secondary contact. I argue for an explicitly historical concept of species, with species' history included in species descriptions.

Acknowledgements

Many people contributed to the completion of this thesis in a variety of ways. My supervisor, Joe Zuccarello, provided tireless support, advice and enthusiasm, both for the preparation of the thesis and in all matters phycological. I could not have found a better person to oversee this project. Pete Ritchie undertook the role of secondary supervisor and was always available when advice was needed.

Maria McMillan provided tireless support for me, my work and my family during this time. Abbie and Lily provided welcome distraction.

Wendy Nelson, Tracy Farr, Kate Neill, Peter Martin, Erasmo Macaya, Fiona Hodge, Kyatt Dixon and Maria helped with collecting specimens. Chris Hepburn, Rochelle Dewdney, Francisco “Pancho” Neira and an anonymous Australian surfer were helpful in directing me to some excellent collecting sites.

Transport and accommodation during fieldwork were provided by Claire Haglund, Simon Hart and Andrea McMillan, Woodford Rouse and family, Maraehako Backpackers, Stuart and Nancy McMillan, Marty Smith and Phil (Napier) and Department of Conservation Chatham Islands staff.

Martin Thiel provided advice on Chapter 2. Alan Millar from the Sydney Royal Botanic Gardens Herbarium and Jenn Dalen from Te Papa Tongarewa (WELT) assisted with access to herbaria. Members of the Wellington phycological group and the VUW Plant Club provided feedback and advice at different stages of the project.

Financial assistance was provided by the Victoria University of Wellington Vice-Chancellor’s Strategic Research Scholarship and by the Education New Zealand Post Graduate Study Abroad Scheme. The Department of Conservation assisted with fieldwork costs.

Many Victoria University staff assisted in many different ways, in particular I would like to thank Cam, Jo, Chris, Mary and Patricia for logistical support.

Essential to completion of this project was the assistance with childcare. In particular, I would like to thank the entire VUW crèche staff, Jane Buchanan, Lenka Rocheford and Pip, and Pip Adam.

Contents

Abstract.....	iv
Acknowledgements.....	vii
List of Figures.....	x
Chapter 1: Introduction and taxonomic background.....	1
1.1 Introduction.....	2
1.2 <i>Carpophyllum</i> Greville.....	3
1.3 <i>Cystophora</i> J. Agardh.....	6
1.4 Research questions.....	8
1.4.1 Hybridisation.....	8
1.4.2 Genetic variation and population structure.....	8
1.4.3 Phylogeny.....	9
1.4.4 Taxonomic questions.....	10
1.5 Structure of this thesis.....	10
1.6 References.....	11
Chapter 2: Decoupling of short and long distance dispersal pathways in the endemic New Zealand seaweed <i>Carpophyllum maschalocarpum</i> (Phaeophyceae, Fucales).....	15
2.1 Abstract.....	16
2.2 Introduction.....	17
2.3 Materials and methods.....	21
2.4 Results.....	27
2.5 Discussion.....	34
2.6 Acknowledgements.....	39
2.7 References.....	39
Chapter 3: Patterns of hybridisation in <i>Carpophyllum</i> (Phaeophyceae, Fucales).....	47
3.1 Abstract.....	48
3.2 Introduction.....	49
3.3 Materials and methods.....	51
3.4 Results.....	60
3.5 Discussion.....	67
3.6 Acknowledgements.....	73
3.7 References.....	73
Chapter 4: Barcoding brown algae: Does <i>cox1</i> identify species of <i>Cystophora</i> (Fucales, Phaeophyceae)?.....	77
4.1 Abstract.....	78
4.2 Introduction.....	78
4.3 Materials and methods.....	81
4.4 Results.....	95
4.5 Discussion.....	99
4.6 Acknowledgements.....	105
4.7 References.....	105

Chapter 5: Taxonomic oversplitting in the New Zealand members of the brown algal genus *Cystophora* (Phaeophyceae, Sargassaceae):

Synonymisation of two species.....	111
5.1 Abstract.....	112
5.2 Introduction.....	112
5.3 Materials and methods.....	118
5.4 Results.....	118
5.5 Discussion.....	121
5.5.1 <i>Cystophora congesta</i> and <i>C. retroflexa</i>	123
5.5.2 Amended species description: <i>Cystophora retroflexa</i>	126
5.5.3 <i>Cystophora distenta</i> and <i>C. scalaris</i>	127
5.5.4 Amended species description: <i>Cystophora scalaris</i>	128
5.6 Conclusion.....	131
5.7 Acknowledgements.....	131
5.8 References.....	131

Chapter 6: Species delimitation in brown algae.....133

6.1 Abstract.....	134
6.2 Introduction.....	134
6.3 Species delimitation.....	135
6.4 Morphological approaches.....	136
6.5 Culture experiments.....	137
6.6 Crossing experiments and reproductive isolation.....	137
6.7 Hybridisation as natural crossing experiments.....	139
6.8 Molecular data.....	142
6.9 Species delimitation in <i>Carpophyllum</i>	147
6.10 Species delimitation in <i>Cystophora</i>	148
6.11 Dispersal and history.....	148
6.12 Genetic distance – molecular data without history.....	150
6.13 Model-based methods.....	155
6.14 Synthesising species concepts.....	156
6.15 References.....	158

Chapter 7: Appendices.....173

7.1 Appendix 1. Alignment of mitochondrial spacer sequences from <i>Carpophyllum maschalocarpum</i>	175
7.2 Appendix 2. Alignment of ITS2 sequences from <i>Carpophyllum</i>	178
7.3 Appendix 3. Alignment of cloned ITS sequences from <i>Carpophyllum</i>	183
7.4 Appendix 4. Alignment of <i>cox1</i> sequences from <i>Cystophora</i>	187
7.5 Appendix 5. Alignment of ITS sequences from <i>Cystophora</i>	199
7.6 Appendix 6. Alignment of mitochondrial spacer sequences from <i>Cystophora retroflexa</i> and related species.....	212
7.7 Appendix 7: Additional work.....	214
7.8 Appendix 8: Poster: Phylogeography of <i>Carpophyllum maschalocarpum</i>	215
7.9 Appendix 9: Hodge <i>et al.</i> (2010) Hybridisation in <i>Carpophyllum</i>	216

List of Figures

Chapter 2

Fig. 2.1. Map of New Zealand showing major currents and bioregions.....	20
Fig. 2.2a. Distribution of mitochondrial haplotypes of <i>Carpophyllum</i> <i>maschalocarpum</i>	29
Fig. 2.2b. Statistical parsimony network inferred from haplotypes of <i>C.</i> <i>maschalocarpum</i>	29
Fig. 2.3. SAMOVA groupings of populations of <i>C. maschalocarpum</i>	31
Fig. 2.4. Mismatch distributions of <i>C. maschalocarpum</i> haplotypes.....	33

Chapter 3

Fig. 3.1. ITS2 statistical parsimony network for <i>C. maschalocarpum</i> and <i>C.</i> <i>angustifolium</i>	62
Fig. 3.2. Geographical patterns of ITS2 ribotypes in <i>Carpophyllum</i>	68

Chapter 4

Fig. 4.1. Map showing sampling locations of <i>Cystophora</i> spp.....	82
Fig. 4.2. Distance tree inferred from <i>cox1</i> sequences from <i>Cystophora</i> spp.....	96
Fig. 4.3. Maximum likelihood tree inferred from <i>cox1</i> and ITS sequences from <i>Cystophora</i> spp.....	97

Chapter 5

Fig. 5.1. Graphs showing morphological variation in receptacles of <i>Cystophora</i> <i>scalaris</i>	119
Fig. 5.2. Statistical parsimony network inferred from mitochondrial spacer sequences of <i>Cystophora retroflexa</i> and <i>C. subfarcinata</i>	120
Fig. 5.3. Morphological variation in <i>Cystophora retorta</i>	122
Fig. 5.4. Morphological variation in <i>Cystophora retroflexa</i>	124
Fig. 5.5. Morphological variation in receptacles of <i>Cystophora scalaris</i>	125
Fig. 5.6. Organisation of ramuli and receptacles of <i>Cystophora retroflexa</i> and <i>C. scalaris</i>	130

Chapter 1

Introduction and background

1.1 Introduction

THE BROWN ALGAE, or Phaeophyceae, are a monophyletic class within the Stramenopiles (Baldauf 2008). They are distantly related to the other eukaryote lineages that have evolved complex multicellular thalli (Metazoans, Fungi, Archaeplastida) and have unique characteristics (Charrier *et al.* 2007). In this project I investigate evolutionary processes in two genera of brown algae, *Carpophyllum* Greville and *Cystophora* J. Agardh. By comparing evolutionary processes in different lineages of life we can progress towards a general understanding of the variation and commonalities in evolutionary processes.

In a review of the state of knowledge of New Zealand marine macroalgae, Nelson (1994) wrote: “Speciation and endemism, and our lack of understanding about rates and processes involved, remain as outstanding issues.” Furoid algae are ideal organisms to investigate evolutionary processes as: (1) They are diverse, Australasia is the probable origin and centre of diversity of the Fucales (Clayton 1984); (2) New Zealand has endemic species and genera and others that are shared with Australia (Parsons 1985); and (3) Fuclean algae have a relatively simple life history, with a single life stage (Chapman 1995). This last character simplifies interpretation of genetic variation, as ecological and evolutionary processes are not acting differently on the dominant macrothallus phase and on the cryptic and phenotypically distinct microthallus phase.

Carpophyllum is endemic to New Zealand, *Cystophora* is found in New Zealand and southern Australia. Species of both genera are among the most common shallow marine macroalgae of New Zealand (Shears & Babcock 2007). Stands of *Carpophyllum* dominate the shallow sub-tidal around a large part of rocky New Zealand coastline. *Cystophora* species are common in the shallow subtidal, rock pools and lower intertidal. Species of both genera provide structure and shelter for other marine organisms and are a significant contributor to primary productivity (Schiel 1990).

Some difficulties arise in species identifications in both genera (Adams 1994). Some forms appear morphologically intermediate between species (Lindauer *et al.* 1961) and ecological studies often identify specimens to generic

level only. This lack of taxonomic information can lead to problems in identifying introduced species, recording distributional changes in species resulting from climate change and in investigating ecological and phylogenetic relationships of these algae.

This project aims to use molecular techniques to investigate genetic variation in these algae at different scales, to provide insights into processes effecting speciation. Specifically, I investigate phylogeny, taxonomy, dispersal and hybridisation. Here I review the taxonomic background and ecology of these genera and outline some questions that will be addressed.

1.2 *Carpophyllum* Greville 1830

Carpophyllum has four species¹, all endemic to New Zealand (Adams 1994). A fifth species of *Carpophyllum* is under consideration based on two specimens collected in Antarctica (M. Clayton & W. Nelson pers. comm.). *Carpophyllum flexuosum* (Esper) Greville is the most widespread species, extending throughout the North and South Islands, Stewart Island, and the Chatham Islands.

Carpophyllum maschalocarpum Turner (Greville) is the most abundant species (Shears & Babcock 2007) and is found around the North Island, Chatham Islands and South Island as far south as Fiordland and Banks Peninsula. *Carpophyllum plumosum* (A. Richard) J. Agardh and *C. angustifolium* J. Agardh have more restricted distributions. *Carpophyllum plumosum* is found only on the eastern coasts of the North Island, from Northland to the Wairarapa Coast, and on the Chatham Islands. *Carpophyllum angustifolium* is found only on the east coast of the North Island, from Northland to just north of East Cape. Adams & Nelson (1985) also recorded *C. angustifolium* and *C. plumosum* from the Three Kings Islands,² off the northern tip of New Zealand's North Island. *Carpophyllum* is generally absent from the sub-Antarctic islands, although *C. flexuosum* has been recorded from the Auckland Islands (Papenfuss 1964), but Adams (1994) suggests

¹ Algaebase (Guiry & Guiry 2011) includes *Carpophyllum macrophyllum* as a current species. This was created by Montagne (1845) from material from the Auckland Islands. According to Lindauer *et al.* (1961) *C. macrophyllum* is a synonym for *C. flexuosum*.

² I examined specimens of *C. angustifolium* collected from the Three Kings Islands held in WELT and consider them to be *C. angustifolium* × *C. maschalocarpum* hybrids.

reports from the sub-Antarctic islands are from drift material. Reports of *Carpophyllum* species from the Kermadec Islands appear to be based only on drift material (Adams & Nelson 1984).

Carpophyllum is placed in the Sargassaceae and is related to some species of *Sargassum* (Draisma *et al.* 2010). The phylogenetic relationships within *Sargassum* have yet to be completely elucidated (Mattio *et al.* 2008, Draisma *et al.* 2010), and *Carpophyllum*'s relationship to the various sub-groupings of *Sargassum* is not entirely clear. BLAST searches and preliminary trees using my sequence data suggest *Carpophyllum* is monophyletic and most closely related to some species of the *Sargassum* subg. *Phyllotricha*, with *S. verruculosum* the closest relative found (albeit with limited species sampling). Draisma *et al.* (2010), in the only published phylogeny that included *Carpophyllum*, placed *Carpophyllum maschalocarpum* as sister species to *S. decurrens* (in the clade *Sargassum*-2 in their system) in a phylogeny inferred from 23S rDNA and *psaA* chloroplast DNA (they did not include *S. verruculosum* in their dataset). My ITS data suggest a close relationship between *S. verruculosum* and *S. decurrens*. Draisma *et al.* (2010) also show a close relationship between *Carpophyllum* and *Nizamuddinina zanardinii* from Oman (suggested by Nelson 1994).

Carpophyllum was established by Greville (1830, xxxii), with two species, *C. maschalocarpum* and *C. flexuosum*. His description is brief: "Frons plana vel compresso-plana. Folia ramiformia, distichia. Receptacula minuta, tuberculata, cylindracea, in racemes marginalibus." The description emphasises the flattened thallus (found in all species except *C. angustifolium*) and distichous laterals. Marginal receptacles are only found in *C. flexuosum*, in other species receptacles are axillary. Lindauer *et al.* (1961, p. 306) listed six characters that, in combination, separate *Carpophyllum* from *Cystophora* and *Sargassum*: "1. the flattened stem; 2. elongate flattened holdfast with short haptera at intervals on edges; 3. rounded or elongate vesicles; 4. leafy appendages; 5. reproductive organs on special branchlets; 6. absence of special basal leaves (these are present in *C. flexuosum*).” I consider that only the flattened holdfast reliably separates *Carpophyllum* from *Sargassum*. *Carpophyllum* appears to have affinities with the distichous members of *Sargassum* subg. *Phyllotricha* (based on unpublished

molecular data³). The holdfast of all *Carpophyllum* species is flattened, creeping over the substrate, with haptera on the edges. In all species except *C. angustifolium*, the holdfast gives rise to a flattened main axis, with no intermediate terete stipe. In *C. angustifolium* the holdfast is flattened and gives rise to a terete stipe. Holdfasts in *Sargassum* subg. *Phyllotricha* species are conical or discoid and give rise to an initially terete stipe. In some species this stipe then becomes flattened and distichous (Womersley 1954, Rainbo Dixon, personal communication, January 16, 2011).

Carpophyllum species were collected by early visitors to New Zealand (Laing 1926) and returned to Europe for description. *Carpophyllum flexuosum* was described by Esper (1802) as *Fucus flexuosus*. *Carpophyllum maschalocarpum* was described by Turner (1811) as *Fucus maschalocarpus*. *Carpophyllum plumosum* was described by A. Richard (1832) as *Sargassum plumosum*, from material collected on Dumont D'Urville's second voyage to New Zealand. *Carpophyllum angustifolium* was described by J. Agardh (1877), from material supplied by Sven Berggren. Synonyms and other taxonomic details are recorded in Lindauer *et al.* (1961).

Some descriptive work on *Carpophyllum*, mainly related to the reproductive anatomy was undertaken by Delf (1939), Dawson (1940), and Naylor (1954). I can find no systematic treatment of *Carpophyllum* since the compilation of early work in Lindauer *et al.* (1961), although Dromgoole's (1973) largely ecophysiological study included some taxonomic notes. Other studies have investigated the chemistry of *Carpophyllum maschalocarpum* (Glombitza & Li 1991a, 1991b, Li & Glombitza 1991) and *C. angustifolium* (Glombitza & Schmidt 1999).

A major survey of macroalgal abundance (Shears & Babcock 2007) found *Carpophyllum maschalocarpum* was the second most abundant macroalgal species around New Zealand (after *Ecklonia radiata*). Other *Carpophyllum* species are locally abundant. Ecological studies have investigated relationships between *Carpophyllum* species and invertebrates, especially effects of urchin grazing, in northern New Zealand (Choat & Schiel 1982, Andrew 1988), the

³ These and other data suggests *Phyllotricha* should be elevated to a distinct genus (Rainbo Dixon, Pers. Comm. 3 May 2011). A paper addressing this is in preparation.

Chatham Islands (Schiel *et al.* 1995) and Fiordland (Villouta *et al.* 2001). Taylor (1998a and b) investigated epifauna on *Carpophyllum*. A study by Cole *et al.* (2001) investigated recruitment of *C. flexuosum* around Northland. Various studies have recorded *Carpophyllum* distribution and abundances in local areas (refs. in Hurd *et al.* 2004).

Carpophyllum species are all dioecious, Fertile thalli have axillary clusters of receptacles. *Carpophyllum flexuosum* also has marginal receptacles on some lateral lamina (Lindauer *et al.* 1961).

1.3 *Cystophora* J. Agardh 1848

Womersley (1964) monographed *Cystophora*, describing 23 species, separated mainly by characters of the vegetative morphology and the arrangement of conceptacles on the receptacle. Adams (1994) lists six species of *Cystophora* from New Zealand: *Cystophora torulosa* (R. Brown) J. Agardh and *C. retroflexa* (Labillardière) J. Agardh are widespread. *Cystophora distenta* J. Agardh, *C. platylobium* (Mertens) J. Agardh, *C. scalaris* J. Agardh, and *C. congesta* Womersley & Nizamuddin ex Womersley are found from Cook Strait southwards. Two species (*C. scalaris* and *C. distenta*) are reported from the Auckland Islands (Adams 1994). Other species extend south as far as Stewart Island and east to the Chatham Islands (Nelson 1994).

Lindauer *et al.* (1961) also list *C. dumosa* (Greville) J. Agardh, a species that Womersley (1964) rejected as a *nomen confusum* under the then current ICBN rules. Womersley (1964) found most of the material used to establish *Cystophora dumosa* consisted of fragments of *C. monilifera*. The earliest valid description of *C. dumosa* is by Sonder (1846). According to Womersley, Sonder's material is a mix of fragments of *C. monilifera* and probably *C. brownii*. Doubtful reports of Australian species from New Zealand include *Cystophora* (= *Caulocystis*) *cephalornithos*, *C. monilifera*, *C. paniculata* and *C. retorta*. These are either misidentifications or Australian specimens with localities mislabelled (Lindauer *et al.* 1961, Womersley 1964, 1987).

Womersley (1964) created a new species, *Cystophora congesta* Womersley & Nizamuddin that includes specimens recorded as *C. dumosa*,

including those from New Zealand. *Cystophora congesta* is reported from the New Zealand mainland from Wellington southwards and from Stewart Island and the Chatham Islands. According to Womersley (1964) it is closely related to *C. retroflexa* and “occasional intergrades occur.” Hybridisation has been reported in other species of *Cystophora*. Lindauer *et al.* (1961) states: “Within the New Zealand species [of *Cystophora*] a degree of hybridization occurs in certain areas where some of the species overlap. *Cystophora platylobium* seems quite distinct, but *C. torulosa*, *C. scalaris*, *C. retroflexa* and *C. dumosa* [= *distenta*] seem to intergrade. A full analysis of the degree of this variation needs to be made.”

Several studies have recorded ecological interactions involving *Cystophora*. *Cystophora* species are settled preferentially by some reef fish (McDermott & Shima 2005) and shelter higher numbers of invertebrates than other, less ramified, large brown algae (Taylor & Cole 1994). Other workers have investigated the chemistry of *Cystophora* (Glombitza *et al.* 1997). Amico (1995) included several *Cystophora* species in a chemotaxonomic study of the Cystoseiraceae.

Several phylogenetic studies have included specimens of *Cystophora*. Rousseau & De Reviers' (1999) phylogeny of the order Fucales, based on rDNA LSU and SSU, included three species of *Cystophora* and other Australian fuclean species. Cho *et al.* (2006) included a single (unidentified) specimen of *Cystophora* from Brighton Beach, New Zealand, in a phylogeny of the Fucales inferred from the *psaA* gene. A multi-locus phylogeny of the “crown” group of brown algae (Silberfield *et al.* 2010) included the Australian species *Cystophora retorta* and *C. grevillei*. Draisma *et al.* (2010) included two species of *Cystophora* in a multi-gene phylogeny. These studies have not elucidated relationships within *Cystophora*, but have identified *Landsburgia* as the sister genus to *Cystophora*, and placed *Cystophora* among other Cystoseiraceae, as well as showing the Cystoseiraceae are polyphyletic.

Most *Cystophora* species have bisexual conceptacles, although sometimes individual conceptacles have mainly oogonia or antheridia. Womersley (1964) notes that *C. monilifera* and *C. racemosa* can have unisexual conceptacles, but either single receptacles bear both types of conceptacle, or adjoining receptacles

bear different types. Only *C. siliquosa* is fully dioecious, with separate male and female plants.

1.4 Research questions

1.4.1 Hybridisation

Hybrids have been reported in *Carpophyllum* (Hodge *et al.* 2010) and *Cystophora* (Womersley 1964). Hybridisation has been extensively investigated in the northern hemisphere fucalean genus *Fucus*. Morphological variation in *Fucus* has been ascribed to both environmental (Chapman 1995) and genetic factors (Anderson & Scott 1998), and natural and artificial hybrids have been studied extensively (McLachlan *et al.* 1971, Coyer *et al.* 2002, Coyer *et al.* 2010).

Two questions immediately arise: (1) Are intermediate forms hybrids?; and (2) are hybrid zones widespread or geographically limited? A number of additional questions are likely to arise: If hybrids are widespread how are the species maintained? If they are limited do they represent zones of secondary contact following speciation due to climate change, geological processes or long distance dispersal (e.g., from Australia)? Further, if hybrids have persisted, is there evidence of hybridisation leading to speciation (reticulate evolution)?

1.4.2 Genetic variation and population structure

Large, negatively buoyant and non-motile eggs are assumed to limit dispersal ability at the gamete and zygote stage in fucalean algae, such as *Carpophyllum* and *Cystophora*, but drifting detached thalli might facilitate longer dispersal distances (Coleman & Brawley 2005, Coleman & Kelaher 2009, Fraser *et al.* 2009). Dispersal might also be extended by the retention of sporelings on the receptacles followed by release and rapid attachment (Schiel & Foster 2006).

Dispersal ability should be closely related to genetic variation within and between populations. Broad-scale studies of population genetic structure in macroalgae often find high population structure, and this is related to (1) limited dispersal ability (Dayton 1973, Coleman & Brawley 2005); and (2) historical

processes, especially historical climate change (Hoarau *et al.* 2007, Maggs *et al.* 2008, Coyer *et al.* 2010). These processes have mostly been studied in the northern hemisphere, especially Europe, and there is little understanding of the effect of these processes in New Zealand, although some phylogeographic work has been done on fish (Hickey *et al.* 2009) and invertebrates (Goldstien *et al.* 2006, 2009).

All *Carpophyllum* species are endemic to New Zealand, which simplifies sampling across the range of the species, and allows population level studies to capture the history of the species. I will use mitochondrial markers to attempt to discern historical patterns of range expansion and contemporary patterns of gene flow by dispersal. Trans-Tasman distributions of some species of *Cystophora* are presumed to arise from long-distance dispersal (Parsons 1985), but the degree of isolation of New Zealand and Australian populations has not been studied. This raises the question of whether there have been multiple dispersal events between New Zealand and Australia or whether dispersal events are rare and ancient, with subsequent genetic differentiation.

1.4.3 Phylogeny

The relationships between species of *Cystophora* and *Carpophyllum* are not clear. In *Carpophyllum* characters such as the terete main axis of *C. angustifolium* or the large basal lamina of *C. flexuosum* might link *Carpophyllum* to a shared common ancestor with *Sargassum*. Alternatively these might be derived characters relating to the sheltered habitat of *C. flexuosum* and the exposed habitat of *C. angustifolium*.

Womersley (1964) produced a tentative phylogeny of *Cystophora* based on the arrangement of branching of the main axis and laterals, and Amico (1995) used a chemotaxonomical approach to group *C. torulosa*, *C. congesta*, *C. scalaris* and *C. monilifera* together, separate from a second group containing *C. expansa* and *C. platylobium* together, with *C. monilifera* (which Amico (1995) considered “more developed”) distinct from both of these. It is not clear whether characters such as branching from the edge (rather than the face) of main axes are shared or

derived characters. In both genera a molecular phylogeny might shed light on the development of these characters.

1.4.4 Taxonomic questions

Characters used to separate New Zealand species of *Cystophora* vary between authors. Lindauer *et al.* (1961) recognise *C. dumosa* as a species with characters intermediate between *C. scalaris*, *C. retroflexa* and *C. torulosa*. Womersley (1964) created a new species, *C. congesta*, which included some specimens assigned to *C. dumosa*, but Womersley's characters are hardly intermediate between the above three species. Adams (1994) also recognised *C. congesta*, but her illustration (Adams 1994, p. 141) does not agree with Womersley's description, and her choice of characters provided in a key for New Zealand species appears closer to Lindauer *et al.*'s (1961) than Womersley's (1964). I have found this key difficult to use, with ambiguous outcomes for several specimens. Molecular markers might be useful for testing species limits in *Cystophora* and assessing the value of different morphological characters.

1.5 Structure of this thesis

In Chapter 2 I investigate the phylogeography of *Carpophyllum maschalocarpum* using extensive sampling of a mitochondrial spacer marker from 32 populations. In Chapter 3 I look at patterns of hybridisation between *Carpophyllum maschalocarpum* and *C. angustifolium* from several populations from the north-eastern North Island. In Chapter 4 I infer a phylogeny for *Cystophora* and compare this to species delimited using a DNA barcoding methodology. Chapter 5 examines the taxonomy of New Zealand species of *Cystophora* in some detail, resulting in the synonymisation of two pairs of species. Finally in Chapter 6 I present a summary of my findings, together with some additional data, in the context of approaches taken by other workers to species delimitation in brown algae.

Because this thesis was written as a series of independent papers, there is some unavoidable repetition of introductory material and general information in each chapter.

1.6 References

- ADAMS, N. M. (1994). *Seaweeds of New Zealand: An illustrated guide*. Christchurch, New Zealand: Canterbury University Press.
- ADAMS, N. M. & NELSON, W. A. (1984). Marine algae of the Kermadec Islands. A list of species. *National Museum of New Zealand Miscellaneous Series*, 10, 1–29.
- ADAMS, N. M. & NELSON, W. A. (1985). The marine algae of the Three Kings Islands. A list of species. *National Museum of New Zealand Miscellaneous Series*, 13, 1–29.
- AGARDH, J. G. (1848). *Species genera et ordines algarum, seu descriptiones succinctae specierum, generum et ordinum, quibus algarum regnum constituitur. Volumen Primum. Algas fucoideas complectens*. Lundae [Lund], Sweden: C.W.K. Gleerup.
- AGARDH, J. G. (1877). De Algis Novae Zelandiae marinis. *Lunds Universitets Års-Skrift, Afdelningen for Matematik och Naturvetenskap*, 14, 1–32.
- AMICO, V. (1995). Marine brown algae of family of Cystoseiraceae: Chemistry and chemotaxonomy. *Phytochemistry*, 39, 1257–1279.
- ANDERSON, C. I. H. & SCOTT, G. W. (1998). The occurrence of distinct morphotypes within a population of *Fucus spiralis*. *Journal of the Marine Biological Association of the U.K.*, 78, 1003–1006.
- ANDREW, N. L. (1988). Ecological aspects of the common sea-urchin, *Evechinus chloroticus*, in northern New Zealand – A review. *New Zealand Journal of Marine and Freshwater Research*, 22, 415–426.
- BALDAUF, S. L. (2008) An overview of the phylogeny and diversity of eukaryotes. *Journal of Systematics and Evolution*, 46, 263–273.
- CHAPMAN, A. R. O. (1995). Functional ecology of fucoid algae: Twenty-Three years of progress, *Phycologia*, 34, 1–32.
- CHARRIER, B., COELHO, S. M., LE BAIL, A., TONON, T., MICHEL, G., POTIN, P., KLOAREG, B., BOYEN, C., PETERS A. F. & COCK, J. M. (2007). Development and physiology of the brown alga *Ectocarpus siliculosus*: two centuries of research, *New Phytologist*, 177, 319–322.
- CHO, G. Y., ROUSSEAU, F., REVIERS, B. DE & BOO, S. M. (2006). Phylogenetic relationships within the Fucales (Phaeophyceae) assessed by the photosystem I coding *psaA* sequences. *Phycologia*, 45, 512–519.
- CHOAT, J. H. & SCHIEL, D. R. (1982). Patterns of distribution and abundance of of large brown algae and invertebrate herbivores in subtidal regional of northern New Zealand. *Journal of Experimental Marine Biology and Ecology*, 60, 129–162.
- CLAYTON, M. N. (1984). Evolution of the Phaeophyta with particular reference to the Fucales. *Progress in Phycological Research*, 3, 11–46.
- COLE, R. G., BABCOCK, R. C., TRAVERS, V. & CREESE, R. G. (2001). Distributional expansion of *Carpophyllum flexuosum* onto wave exposed reefs in northeastern New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35, 17–32.

- COLEMAN, M. A. & BRAWLEY, S. H. (2005). Are life history characteristics good predictors of genetic diversity and structure? A case study of the intertidal alga *Fucus spiralis* (Heterokontophyta; Phaeophyceae). *Journal of Phycology*, **41**, 753–762.
- COLEMAN, M. A. & KELAHER, B. P. (2009). Connectivity among fragmented populations of a habitat-forming alga, *Phyllospora comosa* (Phaeophyceae, Fucales) on an urbanised coast. *Marine Ecology Progress Series*, **381**, 63–70.
- COYER, J. A., HOARAU, G., COSTA, J. F., HOGERDIJK, B., BILLARD, E., VALERO, M., PEARSON, G. A. & OLSEN, J. L. (2010). Evolution and species diversification within the intertidal brown macroalgae *Fucus spiralis*/*F. vesiculosus* species complex in the North Atlantic. *Molecular Phylogenetics and Evolution*, doi:10.1016/j.ympev.2010.11.015.
- COYER, J. A., PETERS, A. F., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2002). Inheritance patterns of ITS1, chloroplasts and mitochondria in artificial hybrids of the seaweeds *Fucus serratus* and *F. evanescens* (Phaeophyceae). *European Journal of Phycology* **37**: 173–178.
- DAWSON, A. E. E. (1940). Studies in the Fucales of New Zealand – II. Observations on the female frond of *Carpophyllum flexuosum* (Esp.) Grev. = *Carpophyllum phyllanthus* (Turn.) Hook. & Harv. *New Phytologist*, **39**, 285–302.
- DAYTON, P. K. (1973). Dispersion, dispersal, and persistence of the annual intertidal alga *Postelsia palmaeformis* Ruprecht. *Ecology*, **54**, 433–438.
- DELFT, E. M. (1939). Studies in the fucales of New Zealand I. The genus *Carpophyllum* Grev. *Journal of Botany, London*, **77**, 129–138.
- DRAISMA, S. G. A., BALLESTEROS, E., ROUSSEAU, F. & THIBAUT, T. (2010). DNA sequence data demonstrate the polyphyly of the genus *Cystoseira* and other Sargassaceae genera (Phaeophyceae). *Journal of Phycology*, **46**, 1329–1345.
- DROMGOOLE, F. I. (1973). *A contribution to the biology of the genus Carpophyllum Grev.* Unpublished PhD. thesis, University of Auckland, Auckland, New Zealand.
- ESPER, E. J. C. (1802). *Icones fucorum cum characteribus systematicis, synonymis (sic) auctorum et descriptionibus novarum specierum. Abbildungen der Tange mit beygefügt systematischen Kennzeichen, Anführungen der Schriftsteller, und Beschreibungen der neuen Gattungen.* Vol. Zweyter Theil. Nürnberg: Raspe.
- FRASER, C. L., NIKULA, R., SPENCER, H. G. & WATERS, J. M. (2009). Kelp genes reveal effects of subantarctic sea ice during the Last Glacial Maximum. *Proceedings of the National Academy of Sciences of the U.S.A.*, **106**, 3249–3253.
- GLOMBITZA, K. W., KEUSGEN, M. & HAUPERICH, S. (1997). Fucophlorethols from the brown alga *Sargassum spinuligerum* and *Cystophora torulosa*. *Phytochemistry*, **46**, 1417–1422.
- GLOMBITZA, K. W. & LI, S. (1991a). Fucophlorethols from the brown alga *Carpophyllum maschalocarpum*. *Phytochemistry*, **30**, 3423–3427.
- GLOMBITZA, K. W. & LI, S. (1991b). Hydroxyphlorethols from the brown alga *Carpophyllum maschalocarpum*. *Phytochemistry*, **30**, 2741–2745.
- GLOMBITZA, K. W. & SCHMIDT, A. (1999). Trihydroxyphlorethols from the brown alga *Carpophyllum angustifolium*. *Phytochemistry*, **51**, 1095–1100.
- GOLDSTIEN, S. J., GEMMELL, N. J. & SCHIEL, D. R. (2009). Colonisation and connectivity by intertidal limpets among New Zealand, Chatham and Sub-Antarctic Islands. I. Genetic connections. *Marine Ecology Progress Series*, **388**, 111–119.
- GOLDSTIEN, S. J., GEMMELL, N. J., & SCHIEL, D. R. (2006). Comparative phylogeography of coastal limpets across a marine disjunction in New Zealand. *Molecular Ecology*, **15**, 3259–3268.
- GREVILLE, R. K. (1830). *Algae britannicae, or descriptions of the marine and other inarticulated plants of the British islands, belonging to the order Algae; with plates illustrative of the genera.* Edinburgh & London, United Kingdom: McLachlan & Stewart; Baldwin & Cradock.

- GUIRY, M. D. & GUIRY, G. M. (2011). *AlgaeBase*. Retrieved, January 21, 2011, from <http://www.algaebase.org>.
- HICKEY, A. J. R., LAVERY, S. D., HANNAN, D. A., BAKER, C. S. & CLEMENTS, K. D. (2009). New Zealand triplefin fishes (family Tripterygiidae): contrasting population structure and mtDNA diversity within a marine species flock. *Molecular Ecology*, 18, 680–696.
- HODGE, F., BUCHANAN, J. & ZUCCARELLO, G. C. (2010). Hybridization between the endemic brown algae *Carpophyllum maschalocarpum* and *C. angustifolium* (Fucales): Genetic and morphological evidence. *Phycological Research*, 58, 239–247.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- HURD, C. L., NELSON, W. A., FALSHAW, R. & NEIL, K. F. (2004). History, current status and future of marine macroalgal research in New Zealand: Taxonomy, ecology, physiology and human uses. *Phycological Research*, 52, 80–106.
- LAING, R. M. (1926). A reference list of New Zealand marine algae. *Transactions of the New Zealand Institute*, 57, 126–185.
- LI, S. & GLOMBITZA, K. W. (1991). Carmalols and phlorethofuhalols from the brown alga *Carpophyllum maschalocarpum*. *Phytochemistry*, 30, 3417–3421.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae. *Nova Hedwigia*, 3, 129–350.
- MAGGS, C. A., CASTILHO, R., FOLTZ, D., HENZLER, C., JOLLY, M. T., KELLY, J., OLSEN, J., PEREZ, K. E., STAM, W., VÄINÖLÄ, R., VIARD, F., & WARES, J. (2008). Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology*, 89, S108–S122.
- MATTIO, L., PAYRI, C. E. & STIGER-POUVREAU, V. (2008). Taxonomic revision of Sargassum (Fucales, Phaeophyceae) from French Polynesia based on morphological and molecular analyses. *Journal of Phycology*, 44, 1541–1555.
- MCDERMOTT, C. J. & SHIMA, J. S. (2006). Ontogenetic shifts in microhabitat preference of the temperate reef fish *Forsterygion lapillum*: implications for population limitation. *Marine Ecology Progress Series*, 320, 259–266.
- MCLACHLAN, J., CHEN, L. C-M. & EDELSTEIN, T. (1971). The culture of four species of *Fucus* under laboratory conditions. *Canadian Journal of Botany*, 49, 1463–1469.
- MONTAGNE, C. (1845). Plantes cellulaires. In: Voyage au Pôle Sud et dans l’Océanie sur les corvettes l’Astrolabe et la Zélée, pendant les années 1837–1838–1839–1840, sous le commandement de M.J. Dumont-d’Urville. Botanique. 1, i-xiv, 1–349.
- NAYLOR, M. (1954). The female conceptacle and young sporeling of the four New Zealand species of *Carpophyllum*. *Transactions and Proceedings of the Royal Society of New Zealand*, 82, 1–6.
- NELSON, W. A. (1994). Distribution of macroalgae in New Zealand – an archipelago in space and time. *Botanica Marina*, 37, 221–223.
- PAPENFUSS, G. F. (1964). Catalogue and bibliography of Antarctic and Sub-Antarctic benthic marine algae. In Lee, M. O. (Ed.), *Bibliography of the Antarctic Seas*, 1, 1–76. Washington DC, American Geophysical Union.
- PARSONS, M. J. (1985). New Zealand seaweed flora and its relationships. *New Zealand Journal of Marine and Freshwater Research*, 19, 131–138.
- RICHARD, A. (1832). Essai d'une flore de la Nouvelle-Zélande. In Lesson, A. & Richard, A. (Eds.) *Voyage de découvertes de l’Astrolabe exécuté pendant les années 1826–1827–1828–1829, sous le commandement de M.J. Dumont d’Urville*. Paris: J. Tastu, Éditeur.
- ROUSSEAU, F. & DE REVIERS, B. (1999). Circumscription of the order Ectocarpales (Phaeophyceae): bibliographical synthesis and molecular evidence. *Cryptogamie Algologie*, 20, 5–18.

- SCHIEL, D. R. (1990). Macroalgal assemblages in New Zealand: structure, interactions and demography. *Hydrobiologia*, 192, 59–76.
- SCHIEL, D. R. & FOSTER, M. S. (2006). The population biology of large brown seaweeds: Ecological consequences of multiphase life histories in dynamic coastal environments. *Annual Review of Ecology, Evolution and Systematics*, 37, 343–372.
- SCHIEL, D. R., ANDREW, N. L. & FOSTER, M. S. (1995). The structure of subtidal algae and invertebrate assemblages at the Chatham Islands, New Zealand. *Marine Biology*, 123, 355–367.
- SHEARS, N. T. & BABCOCK, R. C. (2007). *Quantitative description of New Zealand's shallow sub-tidal reef communities* (Science for Conservation No. 280). Wellington, New Zealand: New Zealand Department of Conservation.
- SILBERFELD, T., LEIGH, J. W., VERBRUGGEN, H., CRUAUD, C., DE REVIERS, B. & ROUSSEAU, F. (2010). A multi-locus time-calibrated phylogeny of the brown algae (Heterokonta, Ochrophyta, Phaeophyceae): Investigating the evolutionary nature of the “brown algal crown radiation.” *Molecular Phylogenetics and Evolution*, 56, 659–674.
- SONDER, O. G. (1846). Algae. In Lehmann, C. (Ed.) *Plantae Preissianae sive enumeratio plantarum quas in Australasia occidentali et meridionali-occidentali annis 1838–1841 collegit Ludovicus Preiss. Vol. 2*. Hamburgi [Hamburg], Germany: Sumptibus Meissneri.
- TAYLOR, R. B. (1998a). Seasonal variation in assemblages of mobile epifauna inhabiting three sub-tidal brown seaweeds in northeastern New Zealand. *Hydrobiologia*, 361, 25–35.
- TAYLOR, R. B. (1998b). Short-term dynamics of a seaweed epifaunal assemblage. *Journal of Experimental Marine Biology and Ecology*, 227, 67–82.
- TAYLOR, R. B. & COLE, R. G. (1994). Mobile epifauna on subtidal brown seaweeds in northeastern New Zealand. *Marine Ecology Progress Series*, 115, 271–282.
- TURNER, D. (1809–1811). *Fuci sive plantarum fucorum generi a botanicis ascriptarum icones descriptiones et historia. Fuci, or coloured figures and descriptions of the plants referred by botanists to the genus Fucus. Vol. 3*. Londini [London]: typis J. M’Creery, impensis J. et A. Arch.
- VILLOUTA, E., CHADDERTON, W. L., PUGSLEY, C. W. & HAY, C. H. (2001). Effects of the sea urchin (*Evechinus chloroticus*) grazing in Dusky Sound, Fiordland, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35, 1007–1024.
- WOMERSELY, H. B. S. (1964). The morphology and taxonomy of *Cystophora* and related genera (Phaeophyta). *Australian Journal of Botany*, 12, 53–110.
- WOMERSLEY, H. B. S. (1987). *The marine benthic flora of southern Australia. Part II*. Adelaide, Australia: South Australian Government Printing Division.
- WOMERSLEY, H. B. S. (1954). Australian species of *Sargassum*, subgenus *Phyllotrichia*. *Australian Journal of Botany*, 2, 337–354.

Chapter 2

Decoupling of short and long distance dispersal pathways in the endemic New Zealand seaweed *Carpophyllum maschalocarpum* (Phaeophyceae, Fucales).

2.1 Abstract

The processes that produce and maintain genetic structure in organisms operate at different time scales and on different life history stages. In marine macroalgae, gene flow occurs through gamete/zygote dispersal and rafting by adult thalli. Population genetic patterns arise from this contemporary gene flow interacting with historical processes. Spatial patterns of mitochondrial DNA variation were analysed to investigate contemporary and historical dispersal patterns in the New Zealand endemic fucalean brown alga *Carpophyllum maschalocarpum* (Turner) Greville. Populations bounded by habitat discontinuities were often strongly differentiated from adjoining populations over scales of tens of kilometres and intra-population diversity was generally low, except for one region of north-east New Zealand (the Bay of Plenty). There was evidence of strong connectivity between the northern and eastern regions of New Zealand's North Island and between the North and South Islands of New Zealand and the Chatham Islands (separated by 650 km of open ocean). Moderate haplotypic diversity was found in Chatham Islands populations while other southern populations showed low diversity consistent with Last Glacial Maximum retreat and subsequent recolonisation. I suggest that ocean current patterns and prevailing westerly winds facilitate long distance dispersal by floating adult thalli, decoupling genetic differentiation of Chatham Island populations from dispersal potential at the gamete/zygote stage. This study highlights the importance of encompassing the entire range of a species when inferring dispersal patterns from genetic differentiation, as realised dispersal distances can be contingent on local or regional oceanographic and historical processes.

Abbreviations:

AMOVA, Analysis of Molecular Variation; LGM, Last Glacial Maximum; CTAB, Cetyltrimethylammonium Bromide; SSCP, Single Stranded Conformational

Polymorphism; SAMOVA, Spatial Analysis of Molecular Variance; SST, Sea Surface Temperature; TBE, Tris-Borate-EDTA.

2.2 Introduction

MULTIPLE dispersal mechanisms are found in many organisms. In many plant species most seed dispersal is limited to local populations, but alternative dispersal mechanisms connect distant populations (Higgins *et al.* 2003). Short dispersal distances are also found in many sessile or demersal marine species, but occasional long distance dispersal events connect populations across wide habitat disjunctions (Pakker *et al.* 1996, Reed *et al.* 1988, Fraser *et al.* 2009). Dispersal distance in marine organisms is often correlated with the duration of a pelagic larval stage (Bohonak 1999), but many marine organisms have the potential to disperse as adults, either because of inherent buoyancy or by rafting on floating substrata (e.g., drift wood, algae or pumice) (Thiel & Gutow 2005a).

Long distance dispersal mechanisms differ from short distance mechanisms in two ways: Firstly, long distance dispersal mechanisms are often mediated by large scale processes (climate patterns, ocean currents, seasonal migrations of animal vectors), rather than local conditions (turbulence, wave regimes, inshore currents). Secondly, long distance dispersal events are rare, stochastic and will seldom be observed directly, but will become evident in a species' range or population genetic structure over long time scales. Consequently, factors that mediate long distance dispersal are decoupled from factors that mediate most observed dispersal (Kinlan *et al.* 2005, Thiel & Haye 2006).

Dispersal by floating thalli might be an important dispersal mechanism for macroalgae (Deysher & Norton 1982, van den Hoek 1987, Thiel & Gutow 2005b, Fraser *et al.* 2009). Floating macroalgal thalli are abundant and can release viable propagules for a considerable time (Kingsford 1992, Smith 2002, Macaya *et al.* 2005, Hernandez-Carmona *et al.* 2006, McKenzie & Bellgrove 2008), allowing dispersal between populations separated by habitat disjunctions such as soft sediment coasts or large stretches of open water (Dethier *et al.* 2003). Genetic evidence for long distance dispersal in macroalgae across open ocean is accumulating but still rare (Tatarenkov *et al.* 2007, Fraser *et al.* 2009).

Fucalean algae (Phaeophyceae) have a monophasic life history with dispersal from maternal thalli by non-motile, negatively buoyant eggs or zygotes (Kendrick & Walker 1991, 1995, Norton 1992, Reed *et al.* 1992, Chapman 1995). Populations are predicted to show strong genetic differentiation because of limited dispersal potential at this stage. Population genetic studies of northern hemisphere Fucaleae have shown population differentiation (Coyer *et al.* 2003, Coleman & Brawley 2005a, Hoarau *et al.* 2007, Tatarenkov *et al.* 2007), but often at larger scales than predicted from direct measurements of propagule dispersal. Long distance dispersal by detached, floating thalli is often invoked to explain population connectivity at large scales (van den Hoek 1987, Coleman & Brawley 2005b, Thiel & Gutow 2005b), especially where local population differentiation is high (Muhlin *et al.* 2008, Coleman & Kelaher 2009),

More generally, the use of molecular markers has uncovered higher than expected genetic population structure in marine organisms. Currents, upwelling, habitat disjunctions, and larval behaviour limit dispersal even for organisms with long pelagic duration (Sotka *et al.* 2004). Historical events are also evident in genetic structure. Marko (2004), for example, found different biogeographical histories in two species of the intertidal gastropod *Nucella* in the northeastern Pacific, despite similar dispersal modes. Retreat from high latitudes following glacial cooling, followed by recolonisation from either northern or southern refugia is a common pattern (Hoarau *et al.* 2007, Maggs *et al.* 2008) although marine species appear to show more variable responses to global cooling than terrestrial species (Wares & Cunningham 2001, Marko *et al.* 2010).

Carpophyllum maschalocarpum (Turner) Greville is a common fucalean alga that is endemic to New Zealand. All *Carpophyllum* species are dioecious. Female thalli produce large (c. 200 μm diameter) non-motile eggs that are extruded to the surface of receptacles but retained by a mucilaginous stalk during fertilisation (Naylor 1954, Clayton 1992). Following fertilisation, zygotes remain attached to the maternal thallus for up to 28 days before release (Delf 1939, Dromgoole 1973). As eggs and zygotes are negatively buoyant, the species is expected to have low dispersal ability at these life stages, but long distance dispersal is possible as adult thalli usually possess pneumatocysts and are

frequently found floating on the sea surface (Kingsford 1992). Floating female thalli bearing zygotes or germlings could establish new populations in the absence of fertile males.

Carpophyllum maschalocarpum forms extensive stands in the shallow subtidal of New Zealand's rocky coasts, where it is the second most abundant algal species by biomass and an important structuring organism (Shears & Babcock 2007). The species is present around both main islands of New Zealand and the Chatham Islands, but is absent from the south-east of the South Island and the sub-Antarctic islands (Morton & Miller 1968, Shears & Babcock 2007, Nelson *et al.* 1991, Schiel *et al.* 1995).

Carpophyllum maschalocarpum's extensive range, potential multiple dispersal modes, ecological importance and ubiquity in rocky habitats make it an ideal organism for investigating phylogeographic processes. The species' range is interrupted by significant habitat breaks (Fig. 2.1), including extensive soft sediment coast around both main islands of New Zealand and stretches of deep water between the main islands (Cook Strait), and between the mainland and the Chatham Islands, 650 km east of the mainland (Morton & Miller 1968, Nelson 1994, Shears *et al.* 2008). The factors mediating the two mechanisms of dispersal should be a major determinant of genetic structure in *Carpophyllum maschalocarpum*. If dispersal by floating thalli is rare, high population differentiation is expected, especially between populations separated by discontinuities in habitat. If floating dispersal is common, connectivity is expected to be mediated by current and wind patterns.

The aim of this study is to determine if the population genetic variation of *C. maschalocarpum* is congruent with dispersal modes. Specifically I compare connectivity between adjacent (10s of kilometres) and distant (100s of kilometres) populations, test for congruence between phylogeographic structure and biogeographic provinces proposed by Shears *et al.* (2008) (Fig. 2.1), and test for genetic isolation by distance and population expansion.

Fig. 2.1 New Zealand map showing major currents (Heath 1985) and locations referred to in the text. Boundaries of bioregions proposed by Shears *et al.* 2008 are indicated by dashed lines and named in italics. EAUC: East Auckland Current; ECC: East Cape Current; SC: Southland Current. Dashed lines show present day 15°C SST isotherm and estimated 15°C isotherm for the Last Glacial Maximum (Barrows & Juggins 2005).

2.3 Materials and Methods

Sampling and DNA extraction

Specimens identified in the field as *Carpophyllum maschalocarpum* were collected from 32 sites around New Zealand (Table 1.1), representing the entire range of *C. maschalocarpum* with the exception of Fiordland, where the species is relatively rare (Shears & Babcock 2007). Thalli were collected haphazardly, at least 1 m apart. Tips of thalli for DNA extraction were rapidly dried and stored in silica gel. In addition, one or more typical thalli from each species/site were prepared as voucher specimens. Five *Carpophyllum angustifolium* J. Agardh specimens and a number of putative *C. angustifolium* × *C. maschalocarpum* hybrid thalli were also included as outgroup specimens and to assist in genetic identification of hybrid specimens. DNA was extracted from approximately 2–5 mg of dry tissue using a modified CTAB buffer procedure (Zuccarello & Lokhorst 2005), with the addition of 1% polyvinylpyrrolidone to the extraction buffer.

PCR conditions and primers

The 23S-tRNA Lys mitochondrial spacer (225 bp) was amplified from a total of 651 samples, using primers (forward: fuc2625F 5'-GCTGTGAGGTTTTTAGCTGACC-3' and reverse: fuc3141R 5'-TCCACCACTCTAACCAACTGAG-3') designed from alignments of the *Fucus*, *Laminaria* and *Desmarestia* mitochondrial genomes (Oudot-Le Seq *et al.* 2002, 2006).

PCR amplifications used a touchdown routine with an initial denaturation step of 94°C for 4 min, followed by 5 cycles of 1 min at 94°C, 1 min at 55°C (–1°C/cycle) and 1 min at 72°C, followed by 30 cycles of 1 min at 94°C, 1 min at 50°C and 1 min at 72°C, with a final extension of 72°C for 10 min. The PCR mix contained 1 µL genomic DNA, 0.5 U Taq DNA polymerase (New England Biolabs, MA, USA) 1X ThermoPol reaction buffer (NEB), 7.5 pmoles of each primer, 200 nmoles dNTP, 5% DMSO and 0.01% BSA. Amplified products were checked for length and yield on 1% agarose gels stained with ethidium bromide.

Populations were screened for variable haplotypes by SSCP (Zuccarello *et al.* 1999, Sunnucks *et al.* 2000). 3–4 μ L PCR product was mixed with 9 μ L 98% formamide, 10 mM NaOH, 0.025% bromophenol blue, 0.025% xylene cyanol. Samples were denatured at 95–100°C for 5 min, then snap cooled on ice before loading. Gels (225 mm long and 0.75 mm thick, D-Code System, BioRad, Hercules, CA. USA) contained 9% 37.5:1 acrylamide/bis-acrylamide (Sigma Aldrich, St Louis, MO. USA) in 0.5X TBE buffer with the addition of 10% glycerol. Electrophoresis was carried out for 18–20 hrs at 7W in 0.5X TBE buffer at 4°C. After electrophoresis, gels were silver stained following protocols in Bassam *et al.* (1991) and banding patterns scored by eye. One or more of each haplotype indicated by SSCP was sequenced from each population, as well as any samples giving ambiguous SSCP profiles. Amplification for sequencing used the same PCR conditions and reverse primer as above, but used a novel forward primer (fuc2512F: 5' -CCGGGTAGCTACATCGAGAA-3') that anneals approximately 100 bp further toward the 5' end of the fragment, to ensure accurate reads of the 5' end of the spacer.

Generally SSCP banding and sequencing results were concordant. In very few cases haplotypes were not easily distinguished by SSCP and all samples were sequenced. PCR products were cleaned with ExoSAP-IT (USB, Cleveland, OH, USA) enzymes and sequenced commercially (Macrogen Inc., Seoul, South Korea). Sequences from 210 samples were included in the alignment.

Data analysis

Sequences were aligned using the ClustalW algorithm in MEGA 3.1 (Kumar *et al.* 2004) and checked by eye. Alignment was straightforward, requiring insertion of a single gap to accommodate outgroups. Haplotype networks were generated in TCS 1.21 (Clement *et al.* 2000). Arlequin 3.11 (Excoffier *et al.* 2005) was used to generate summary population genetic statistics and perform neutrality tests (Tajima's D, Fu & Li's F* and D*, and Fu's F_s). Neutrality indices were tested against 2000 coalescence simulations assuming demographic stability and neutrality.

Pairwise Φ_{st} values were calculated in Arlequin 3.11 for Bay of Plenty populations (in which shared haplotypes were reasonably frequent) and tested for

significance by 1023 permutations. Mantel tests, also implemented in Arlequin 3.11, were used to determine isolation by distance, first for all populations then for mainland populations (excluding Chatham Island populations). A matrix of shortest over-water geographic distances between populations was calculated and tested for correlation with pairwise Φ_{st} , with significance tested against 2000 permutations. A second test was made using only the north-south component of these distances.

AMOVAs, implemented in Arlequin 3.11, were used to compare population partitions found in *C. maschalocarpum* with proposed marine biogeographic regions (Shears *et al.* 2008). These were compared to genetic structure indicated by SAMOVA (Spatial Analysis of Molecular Variance, Dupanloup *et al.* 2002) implemented in SAMOVA 1.0. This uses a simulated annealing process to group adjoining populations into K-groups so as to maximise genetic distance between groups (maximum F_{ct}), while minimising distance within groups (minimum F_{sc}). K was iterated in successive runs until a maximum F_{ct} was calculated, and significance of fixation indexes tested by 1000 permutations.

Mismatch distribution analysis (Rogers & Harpending 1992), implemented in Arlequin 3.11, was used to test for signatures of spatial expansion in frequency distributions of pairwise sequence differences. Analyses were applied to the total data and separately to populations partitioned into two groups as indicated by SAMOVA and tested against 5000 coalescent simulations assuming spatial expansion. Results were interpreted in the context of simulations using stepping stone and infinite island models. Here unimodal frequency distributions of pairwise mutational distances are regarded as evidence for past population expansions, but this signature is weakened or removed where populations are subdivided with low migration rates (Ray *et al.* 2003, Excoffier 2004).

Table 2.1: Geographic locations, numbers of specimens sampled, mtDNA spacer haplotype designations and population diversity indices for *Carpophyllum maschalocarpum*. Hyphens in gene diversity and nucleotide diversity columns indicate populations where substantial introgression by *C. angustifolium* was detected and values were not calculated.

Map ID (Fig 2.2)	Site	N	Latitude	Longitude	Herbarium number of exemplars	Haplotypes present (n)	Gene diversity $h \pm$ S.D.	Nucleotide diversity $\pi \pm$ S.D.
1	Hooper Point, Northland	47	34° 24' 56" S	172° 51' 13" E	-	21 (10), 34 (34), 65 (1), 66 (1), 67 (1)	-	-
2	North Cape, Northland	10	34° 24' 57" S	173° 03' 09" E	WELT A031664	30 (8), 31 (1), 32 (1)	-	-
3	Matai Bay, Northland	18	34° 49' 37" S	173° 24' 54" E	-	68 (18)	0	0
4	Ahipara, Northland	25	35° 10' 10" S	173° 06' 30" E	-	40 (1), 52 (24)	0.0800 $\pm$ 0.0722	0.0004 $\pm$ 0.0008
5	Wekarua, Northland	19	34° 56' 42" S	173° 39' 12" E	WELT A031665	30 (7), 33 (1), 47 (7), 50 (4)	-	-
6	Cape Wiwiki, Northland	21	35° 09' 22" S	174° 07' 20" E	-	21 (21)	0	0
7	Bland Bay, Bay of Islands	20	35° 20' 31" S	174° 21' 58" E	WELT ASG290	13 (20)	0	0
8	Poor Knights Islands	10	35° 27' 40" S	174° 44' 20" E	WELT A031311, A031317, A031319	21 (10)	0	0
9	Waterfall Reef, Leigh, Northland	11	36° 16' 08" S	174° 47' 53" E	WELT A031318, A031320	01 (2), 13 (8), 36 (1)	0.4727 $\pm$ 0.1617	0.0067 $\pm$ 0.0049
10	Matheson Bay, Northland	34	36° 18' 05" S	174° 47' 58" E	-	01 (2), 13 (27), 55 (1), 56 (1), 57 (2), 66 (1)	0.3708 $\pm$ 0.1042	0.0063 $\pm$ 0.0044
11	Sailors Grave, Bay of Plenty	20	36° 57' 37" S	175° 50' 43" E	WELT A031312– A031316, A031331	20 (1), 21 (1), 25 (2), 27 (1), 28 (13), 29 (1), 70 (1)	0.5842 $\pm$ 0.1270	0.0084 $\pm$ 0.0056

Table 2.1 (Continued)

Map ID (Fig 2.2)	Site	N	Latitude	Longitude	Herbarium number of exemplars	Haplotypes present (n)	Gene diversity $h \pm$ S.D.	Nucleotide diversity $\pi \pm$ S.D.
12	Otanga, Bay of Plenty	25	37° 32' 59" S	178° 09' 41" E	-	14 (3), 16 (3), 21 (2), 22 (1), 26 (3), 28 (9), 39 (3), 43 (1)	0.8367 $\pm$ 0.0539	0.0152 $\pm$ 0.0090
13	Maraehako Bay, Bay of Plenty	25	37° 40' 20" S	177° 47' 48" E	-	14 (15), 15 (3), 16 (2), 17 (1), 18 (1), 19 (2), 23 (2), 38 (2), 71 (1), 72 (1)	0.7425 $\pm$ 0.0813	0.0065 $\pm$ 0.0045
14	Maketu, Bay of Plenty	24	37° 44' 41" S	176° 28' 13" E	WELT A031333	58 (21), 59 (1), 60 (2)	0.2355 $\pm$ 0.1093	0.0017 $\pm$ 0.0019
15	Opape, Bay of Plenty	21	37° 58' 20" S	177° 25' 15" E	-	01 (1), 14 (19), 69 (1),	0.1857 $\pm$ 0.1102	0.0021 $\pm$ 0.0021
16	Horoera, East Cape	24	37° 38' 17" S	178° 28' 29" E	WELT A031310	21 (19), 52 (1), 61 (1), 62 (1), 64 (2)	0.3768 $\pm$ 0.1224	0.0050 $\pm$ 0.0038
17	Whale Bay, Waikato	20	37° 49' 22" S	174° 48' 02" E	-	01 (17), 06 (1), 35 (1), 37 (1)	0.2842 $\pm$ 0.1284	0.0013 $\pm$ 0.0016
18	Kiritehere, Waikato	20	38° 19' 31" S	174° 42' 07" E	WELT A024176	01 (17), 02 (1), 04 (1), 05 (1)	0.2842 $\pm$ 0.1284	0.0022 $\pm$ 0.0022
19	Kaiti Beach, Gisborne	23	38° 40' 56" S	178° 01' 46" E	WELT A031332	21 (21), 63 (1), 67 (1)	0.1700 $\pm$ 0.1025	0.0054 $\pm$ 0.0040
20	Waipatiki, Hawke Bay	23	39° 18' 19" S	176° 58' 29" E	WELT A031324, A031325	08 (21), 11 (2)	0.1660 $\pm$ 0.0976	0.0015 $\pm$ 0.0017
21	Ahuriri St, Port Napier	15	39° 28' 42" S	176° 54' 13" E	WELT A031323	08 (15)	0	0
22	Pihama, South Taranaki	21	39° 31' 17" S	173° 54' 55" E	WELT A031321, A031322, A031330	01 (18), 07 (3)	0.2571 $\pm$ 0.1104	0.0011 $\pm$ 0.0015
23	Pukerua Bay, Wellington	20	41° 01' 29" S	174° 54' 06" E	WELT A031309	01 (20)	0	0

Table 2.1 (Continued)

Map ID (Fig 2.2)	Site	N	Latitude	Longitude	Herbarium number of exemplars	Haplotypes present (n)	Gene diversity $h \pm$ S.D.	Nucleotide diversity $\pi \pm$ S.D.
24	Riversdale, Wairarapa	15	41° 06' 25" S	176° 04' 11" E	WELT A031308	01 (12), 12 (3)	0.3429 $\pm$ 0.1278	0.0046 $\pm$ 0.0036
25	Wharanui, Kaikoura	23	41° 55' 20" S	174° 06' 00" E	*	01 (23)	0	0
26	Waipapa Point, Kaikoura	17	42° 12' 41" S	173° 52' 29" E	*	01 (17)	0	0
27	Ripapa Island, Banks Peninsula	24	43° 37' 07" S	172° 45' 13" E	*	01 (24)	0	0
28	Cape Three Points, Banks Peninsula	16	43° 49' 50" S	172° 54' 30" E	*	01 (16)	0	0
29	Wharekauri, Chatham Islands	10	43° 42' 18" S	176° 34' 55" W	WELT ASG257	24 (10)	0	0
30	Point Dorset, Chatham Islands	24	43° 49' 34" S	176° 42' 21" W	WELT A031328	01 (16), 03 (6), 24 (2)	0.5072 $\pm$ 0.0929	0.0060 $\pm$ 0.0043
31	Waitangi, Chatham Islands	22	43° 56' 42" S	176° 33' 42" W	WELT A031329	01 (6), 08 (1), 09 (3), 10 (5), 24 (4), 39 (3)	0.8398 $\pm$ 0.0368	0.0206 $\pm$ 0.0117
32	Manukau Reef, Chatham Islands	5	37° 44' 41" S	176° 28' 13" W	WELT A024213	08 (5)	0	0

* Vouchers for these sites were lost as a result of a chemical spillage accident.

2.4 Results

Haplotypic diversity and population structure

SSCP and subsequent sequencing identified 67 haplotypes (Genbank numbers HM070070 to HM070166; Appendix 1). While SSCP might underestimate haplotypic diversity, sequencing of a large proportion of samples should reduce the likelihood of significant underestimation of diversity. A statistical parsimony network (Fig. 2.2b) grouped haplotypes into three clusters, each separated by five or more steps. The largest cluster (labelled (1) in Fig. 2.2b) joined 52 haplotypes. These are considered to arise from the *C. maschalocarpum* mitochondrial lineage. A second cluster (labelled (2) in Fig. 2.2b) joined four closely related haplotypes from North Cape (pop. 2 in Fig. 2.2a) and Wekarua (pop. 5). This cluster was five steps removed from the nearest *C. maschalocarpum* haplotype and six steps removed from the nearest haplotype of the sister species *C. angustifolium*. It is unclear whether these haplotypes arise from the *C. angustifolium* or *C. maschalocarpum* maternal lineage. A third cluster (labelled (3) in Fig. 2.2b), joined eleven haplotypes from *C. angustifolium* and putative hybrid specimens, and was six steps removed from the nearest *C. maschalocarpum* haplotype. Haplotypes from *Carpophyllum angustifolium* and any specimens considered to have hybrid origin (including all of pops. 1, 2 and 5), are not included in population genetic analyses. Three haplotypes (15, 23 and 36) were found only in specimens identified morphologically as either *C. angustifolium* or hybrids, but which differed from haplotypes found in specimens morphologically identified as *C. maschalocarpum* by only one mutational step. These are considered to have arisen in the *C. maschalocarpum* mitochondrial lineage and are included in subsequent analyses.

Geographically restricted haplotypes were common (Fig. 2.2a). Forty-one *C. maschalocarpum* haplotypes were private (sampled from a single population only). At two sites, Matai Bay (pop. 3 in Fig 2.2a) and Maketu (pop. 14) all haplotypes sampled were private. At Ahipara (pop. 4) both haplotypes sampled were private aside from a single specimen of haplotype 52 also sampled at Horoera (pop. 16). Haplotypes from Bay of Plenty and Hawke Bay populations were not shared with neighbouring regions, apart from one Hawke Bay haplotype that was also sampled in the Chatham Islands. Four haplotypes were endemic to the Chatham Islands. Population differentiation was moderate to strong between populations with shared haplotypes (Table 2.2).

A small number of haplotypes were widely distributed (Fig. 2.2a). Haplotype 01 was sampled in Northland (pops. 9 and 10), Bay of Plenty (pop. 15), Riversdale (pop. 24), the west coast of the North Island (all populations except 4), the Chatham Islands (pops. 31 and 30) and in all South Island populations. Haplotype 01 was the only haplotype sampled from the South Island. Three haplotypes were shared between Northland and East Coast populations. Haplotype 21 was sampled frequently in both regions. Also shared were haplotype 52 (pop. 4 and a single sample from pop. 16) and haplotype 67 (pops. 1 and 19). Haplotype 67 appears to be derived from the *C. angustifolium* lineage and is 14 mutational steps removed from the closest *C. maschalocarpum* haplotype. Three Chatham Island haplotypes were shared with mainland New Zealand: haplotype 39 was sampled in Otanga (pop. 12) and Waitangi (pop. 31) only; haplotype 08 was shared with two Hawke Bay sites (pops. 20 and 21); and the common southern haplotype 01 was sampled at two Chatham Islands sites (pops. 30 and 31).

Haplotype and nucleotide diversity (Table 2.1) was highest in three Bay of Plenty populations: Sailors Grave, Otanga and Maraehako Bay (pops. 11, 12 and 13 respectively), and in two Chatham Island populations: Waitangi and Point Dorset (pops. 31 and 30). Diversity was lowest in the South Island (a single haplotype sampled in four populations).

Biogeographic regions

SAMOVA analyses reached maximum F_{ct} at $K = 14$ ($F_{ct} = 0.811425$). At $K = 14$ populations were partitioned into five groups and nine singletons (Fig. 2.3). The largest group of ten populations, dominated by the common haplotype 01, included all South Island sites, Riversdale (pop. 24), Point Dorset (pop. 30) and all west coast North Island

Fig. 2.2. (opposite) (a) Distribution of mitochondrial haplotypes of *Carpophyllum maschalocarpum*. Area of pie charts is proportional to sample size. Locations are described in Table 2.1. (b) Statistical parsimony network inferred by TCS software (95% confidence limits). Area of circles is proportional to number of haplotypes sampled. Small black circles represent extinct or unsampled haplotypes. Dashed line indicates an alternative but unlikely connection.

Table 2.2. Pairwise Φ_{st} estimates from mtDNA spacer data (below diagonal) and minimum round coast distances (km, above diagonal) for Bay of Plenty populations of *Carpophyllum maschalocarpum*. Significance shown as: ** = $p \leq 0.01$ and * = $p \leq 0.05$ after sequential Bonferroni correction for multiple tests. N. S. = No significant difference. Population numbers refer to Table 2.1/Fig. 2.2.

	Sailors Grave	Maketu	Opape	Maraehako Bay	Otanga
Sailors Grave (pop. 11)	-	103	179	190	215
Maketu (pop. 14)	0.54931**	-	87	117	151
Opape (pop. 15)	0.46280**	0.84906**	-	48	86
Maraehako Bay (pop. 13)	0.42232**	0.68024**	N. S.	-	38
Otanga (pop. 12)	0.12573*	0.26357**	0.11334*	0.12548**	-

sites except Ahipara (pop. 4). This group was unchanged for all iterations of $K > 2$. Chatham Island populations did not group together despite their proximity. Only Waitangi and Manukau Reef (pops. 31 and 32) were retained in the same group above $K=3$ and these two populations separated into singletons at $K = 13$ and above. Wharekauri (pop. 29) grouped with Northland populations and two populations south of East Cape, then formed a singleton at $K = 8$. Point Dorset (pop. 30) remained grouped with South Island and southern North Island populations.

AMOVA based on biogeographic regions and provinces proposed by Shears *et al.* (2008) explained considerably less between-group variance than the maximum obtained by SAMOVA (Table 2.3). The low haplotype diversity in South Island and southern North Island samples precluded showing the North/South disjunction found in other studies. A biogeographic break at East Cape was not shown in *C. maschalocarpum*, rather the data showed connectivity between two populations south of East Cape (pops. 16 and 19) and Northland.

Fig. 2.3. Grouping of populations of *Carpophyllum maschalocarpum* inferred by SAMOVA to maximise F_{ct} . Shaded areas show groups inferred with $K = 14$. Division into two groups shown was used for mismatch distribution analyses (Fig. 2.4).

Departures from neutrality

Neutrality tests compare estimates of θ using different parameters. Data from demographically stable populations using neutral markers should estimate similar values of θ . Discrepancies in estimates of θ indicate departure from neutrality, with different tests sensitive to different underlying factors (Fu 1997).

Tajima's D value (Tajima 1989) was negative ($D = -0.7109$), which suggests population expansion, but not significant ($P = 0.268$). Fu and Li's D^* test (Fu & Li 1993) was significant ($D^* = -2.3973$, $P = 0.016$). Fu and Li's F^* test (Fu & Li 1993) was also significant ($F^* = -2.1275$, $P = 0.024$). Fu's F_s value, which is sensitive to population

Table 2.3: Hierarchical partitioning of molecular variance estimated by AMOVA from *Carpophyllum maschalocarpum* mtDNA spacer data. (a) Variation explained by grouping in biogeographic regions proposed by Shears *et al.* (2008); and (b) groupings at maximum F_{ct} inferred by SAMOVA.

a. Groupings by Biogeographic Regions

<i>Source of variation</i>	<i>d.f.</i>	<i>Sum of squares</i>	<i>Variance</i>	<i>Percentage of variation</i>
Among groups	6	603.55	$V_a = 1.026$	33.05
Among populations	22	626.90	$V_b = 1.402$	45.14
Within populations	552	373.84	$V_c = 0.677$	21.81
Fixation indices: $F_{sc} = 0.674$, $F_{st} = 0.782$, $F_{ct} = 0.331$				

b. Groupings by SAMOVA (Fig. 2.3)

<i>Source of variation</i>	<i>d.f.</i>	<i>Sum of squares</i>	<i>Variance</i>	<i>Percentage of variation</i>
Among groups	13	804.64	$V_a = 1.652$	81.14
Among populations	15	6.94	$V_b = 0.004$	0.21
Within populations	550	208.93	$V_c = 0.380$	18.65
Fixation indices: $F_{sc} = 0.011$, $F_{st} = 0.814$, $F_{ct} = 0.811$				

expansion (Fu 1997), was strongly negative, and highly significant ($F_s = -25.2996$, $P = 0.0002$). F_s tests were also carried out on data split into two groups as indicated by SAMOVA. Group 1 (North Island west coast/South Island/Riversdale/Port Dorset) and Group 2 (all other samples) both gave negative and significant F_s values ($F_s = -11.7910$, $P < 0.0001$ and $F_s = -23.1464$, $P = 0.0002$ respectively).

Isolation by distance

Correlation between geographic distance and genetic distance was low but significant ($r^2 = 0.10388$, $P < 0.0001$). Recalculating distances to exclude east-west distances gave

similar results ($r^2 = 0.10820$, $P < 0.0001$). Stronger correlation ($r^2 = 0.20768$, $P < 0.0001$) was obtained by excluding Chatham Islands samples from the analysis.

Mismatch analysis

Mismatch analyses showed an approximation to the expected distribution under a model of sudden expansion (Fig. 2.4). Separation of data into two groups as indicated by SAMOVA resulted in two distinct distributions: an L-shaped distribution in Group 1 (North Island west coast/South Island/Riversdale/Port Dorset) and a unimodal distribution in Group 2 (all other populations). No group showed significant departure from the expected distribution under a model of sudden expansion (Group 1: $P = 0.355$; Group 2: $P = 0.881$; combined populations: $P = 0.624$). M (the parameter for gene flow where $M = 2Nm$ and N is the deme size and m is the migration rate) is low in Group 1 ($M = 0.035$) and high in Group 2 ($M = 12.156$).

Fig. 2.4 Mismatch distributions: Group 1 (South Island/North Island West Coast, Riversdale and Point Dorset populations): black bars; Group 2 (all other populations): Grey bars; Combined populations: open bars.

2.5 Discussion

Patterns of gene flow that arise from multiple dispersal modes are complex as dispersal patterns might overlap, especially when each mode is facilitated by different physical processes and influenced by separate historical events. In these data three main patterns are evident: (1) Often high genetic differentiation between populations, including adjacent populations; (2) Connectivity between some distant populations, with only slight overall isolation by distance; and (3) Generally low southern diversity, with the notable exception of the Chatham Islands. These patterns are consistent with fucalean morphological and life history characteristics that limit dispersal at the gamete/zygote stage, but facilitate occasional long distance dispersal (Thiel & Haye 2006). Historical climatic changes also appear to have influenced the distribution of *C. maschalocarpum*.

Genetic differentiation

Genetic differentiation between many populations of *Carpophyllum maschalocarpum* was in the order of that expected for fucalean algae (Coleman & Brawley 2005b, Hoarau *et al.* 2007, Tatarenkov *et al.* 2007, Coleman & Kelaher 2009). Strong genetic differentiation was found between some adjacent populations that were separated by habitat discontinuities, even where distances were relatively short (for example, Cape Wiwiki (pop. 6) and Bland Bay (pop. 7), 39 kilometres apart but separated by the estuarine Bay of Islands). These discontinuities would be expected to provide a barrier to gametic or zygotic dispersal, but not to dispersal of floating thalli. Despite the abundance of floating thalli (Kingsford 1992), it appears that this mode of dispersal seldom contributes to gene flow.

Explanations for limited gene flow by floating thalli include dioecy, small target areas for immigrant floating algae (e.g., Maketu a short, 1 km, rocky peninsula flanked by long stretches of soft sediment coast) and density blocking (Deysher & Norton 1982, Hewitt 1996, Austerlitz *et al.* 2000), where relatively rare immigrants seldom contribute alleles to established stands of algae. High regional haplotype endemism might reflect isolation of water masses, for example limited current transport into embayments such as Hawke Bay and Bay of Plenty (Ridgway 1960, 1962, Stanton *et al.* 1997). Direct studies of transport of floating algae have not been undertaken in New Zealand. A high prevalence of westerly winds might mean surface floating species are usually exported and lost from nearshore habitat.

Long distance dispersal

Shared haplotypes in Northland and North Island east coast populations, and North Island and Chatham Island populations, show some haplotypes have dispersed over long distances. Haplotype 21 is common in Northland and East Coast populations, and it is improbable that this haplotype arose independently in two areas. Further, haplotype 67, which clusters with *C. angustifolium*, was only sampled at Hooper Point (pop. 1) and Kaiti Beach (pop. 19). Kaiti is well outside of the range of *C. angustifolium*, which does not extend south of East Cape (Morton & Miller 1968 and my observations). It appears this haplotype entered the *C. maschalocarpum* gene pool by hybridisation and dispersed to southern populations (Hodge *et al.* 2010). This connection, and the connection between the mainland and the Chatham Islands, is congruent with contemporary currents and surface winds. Northland and the East Coast are connected by the East Auckland/East Cape Current. This current does not extend to the inner area of the Bay of Plenty (Heath 1985, Stanton *et al.* 1997) and my data suggest Bay of Plenty populations occupy an isolated eddy. Despite their spatial separation, the outer populations (Sailors Grave-pop. 11 and Otanga-pop. 12) are more connected to each other than to inner populations. Haplotype 28 occurs in high frequencies in these populations, but was absent from the other Bay of Plenty populations.

Four lineages are present in Chatham Island populations, with haplotypes in each lineage the same or closely related to haplotypes found in North and South Island populations, suggesting at least four dispersal events between mainland New Zealand and the Chatham Islands. High connectivity between the mainland and the Chatham Islands is surprising as many Chatham Island marine algae are endemic (Nelson *et al.* 1991, Schiel *et al.* 1995), several common mainland species are absent (e.g., the non-buoyant laminarian species *Ecklonia radiata* and *Lessonia variegata*) (Nelson *et al.* 1991, Nelson 1994), and studies in other organisms have shown genetic isolation in Chatham Island populations (Goldstien *et al.* 2009). My data suggest a very different pattern for algae capable of dispersal by floating, with prevailing westerly winds and currents facilitating eastward dispersal from the mainland to the Chatham Islands (Chiswell & Booth 2008). Removing the East-West component of geographic distance between populations did not affect correlation with genetic connectivity. These data

suggest wind and currents facilitate long distance transport to the Chatham Islands, while North to South movement proceeds mostly by stepping stone dispersal.

The North/South disjunction found in other New Zealand studies (Apte & Gardner 2002, Sponer & Roy 2002, Waters & Roy 2004, Ayres & Waters 2005, Goldstien *et al.* 2006) was evident in the sudden reduction in haplotype diversity in southern populations, but this occurs further north than the break found in previous studies. A barrier to dispersal by upwelling or current transport of propagules offshore around the north of the South Island has been suggested, but evidence for this is equivocal (Ross *et al.* 2009). I suggest historical climate change explains the pattern in these data better than a dispersal barrier and discuss this below. Bioregions proposed by Shears *et al.* (2008) are partly reflected in the distribution of *C. maschalocarpum* haplotypes (Table 2.3). In particular, disjunctions just south of Ahipara and south of Hawke Bay (pops. 4 and 20/21, Fig. 2.3) in these data are congruent with boundaries of bioregions in Shears *et al.* (2008). Conversely, the connection between Northland and the East coast, and the relative isolation of the Bay of Plenty and Hawke Bay populations of *C. maschalocarpum* is not evident in data obtained by Shears *et al.* (2008). It is not clear whether this arises from sampling differences, the stochastic nature of floating dispersal or factors unique to *C. maschalocarpum*.

This pattern of population differentiation and connectivity is consistent with a species that is generally restricted in dispersal, but with intermittent long distance dispersal by floating (Thiel & Haye 2006). Leapfrog dispersal, where more distant populations show greater connectivity than adjacent populations, is a predicted outcome of intermittent floating dispersal (Thiel & Haye 2006) and has been found in a variety of organisms (Snyder & Gooch 1973, Bockelmann *et al.* 2003, Colson & Hughes 2004). In *C. maschalocarpum*, leapfrog dispersal appears to be overlaid on patterns of low dispersal or stepping stone dispersal.

Low southern diversity and post-LGM recolonisation

The abundant centre model (Eckert *et al.* 2008) predicts greatest genetic diversity in the centre of a species' range, with a decline toward peripheral populations. My data are broadly consistent with this model. Diversity is highest in the Bay of Plenty in North-East New Zealand and generally declines with distance from this region. But there are two departures from this pattern: (1) Haplotypically diverse populations in the far north

(pops. 1, 2 & 5); and (2) high haplotypic diversity in the Chatham Islands. I invoke both contemporary factors and historical factors to explain these departures.

Far north diversity appears to have hybrid origin, with gene flow between *C. angustifolium* and *C. maschalocarpum*. Far north populations have been identified as either species by different workers (e.g., *C. maschalocarpum* by Shears & Babcock 2007, *C. angustifolium* by Hay & Grant 2003). Hybrids have been reported between *Carpophyllum* species based on intermediate morphology (Lindauer *et al.* 1961, Dromgoole 1973, Shears & Babcock 2007) and molecular data (Hodge *et al.* 2010), and hybridisation in fucalean species has been widely reported (Scott & Hardy 1994, Kim *et al.* 1997, Coyer *et al.* 2002, Engel *et al.* 2005).

The pattern of high northern diversity and low southern diversity suggests relatively recent southward expansion with warming of the ocean waters following the Last Glacial Maximum (LGM). Water temperature is a major determinant of species range in many macroalgae (Lüning 1990, Adey & Steneck 2001). Low diversity at high latitudes is often reported in species that have recolonised high latitudes during post-glacial warming (Coyer *et al.* 2003, Marko 2004, Hoarau *et al.* 2007, Maggs *et al.* 2008, Fraser *et al.* 2009). The present day distribution of *Carpophyllum maschalocarpum* extends south to Fiordland (46° S) on the South Island's west coast (Nelson *et al.* 2002), where warm currents push in from the Tasman Sea, but only as far south as Banks Peninsula (43° 45' S) on the east coast, where cold currents push up from the south (Adams 1994). This southern limit is correlated with the 15°C Sea Surface Temperature (SST) isobar for the warmest month of the year (Fig. 2.1). While there are few direct data on thermal tolerances in *C. maschalocarpum*, the correlation between southern limit and SST on both coasts suggests this is a low temperature limit. Records of specimens of *C. maschalocarpum* from driftlines on Stewart Island and Sub-Antarctic Islands (Dromgoole 1973) show thalli are transported further south, but populations have not established.

Climate reconstructions of the LGM (Barrows & Juggins 2005) place the 15°C warmest month SST isobar north of Taranaki on the New Zealand's west coast and north of Hawke Bay on the east coast (Fig. 2.1). These temperatures would have restricted *C. maschalocarpum* to the northern half of the North Island during the LGM. Recolonisation southwards down the North Island west coast, through the reopened

Cook Strait (Lewis *et al.* 1994) and down the east coast of the South Island would explain the low diversity and the different patterns of population expansion indicated by mismatch distributions.

Genetic diversity in northern hemisphere algae has been shown to retain the signature of past climate events – in particular the last glaciation, which removed algae from higher latitudes (reviewed by Maggs *et al.* 2008). Similar patterns have been shown for marine species in the southern hemisphere (Hickey *et al.* 2009) including algae (Fraser *et al.* 2009). The distribution of haplotype diversity in my data is strikingly similar to patterns of the same marker in *Fucus serratus* from Europe (Hoarau *et al.* 2007), and is similar to distributions interpreted as showing post-glacial range expansion for a number of species (Hewitt 1999, Provan *et al.* 2005, Maggs *et al.* 2008).

I offer two explanations for the presence of four endemic haplotypes in the Chatham Islands. Either this is further evidence for the ease of dispersal to the Chatham Islands, with these haplotypes either extinct or not sampled on the main islands of New Zealand, and with dispersal pathways facilitating co-dispersal of closely related haplotypes (e.g., haplotypes 08, 09 and 10); or Chatham Island populations of *C. maschalocarpum* have persisted for a considerable time, allowing the accumulation of unique mutations. At present summer water temperatures around the Chatham Islands are elevated as seasonal north/south movement of the sub-tropical convergence is constrained by an area of shallow bathymetry, the Chatham Rise (Stanton 1997). Paleoclimate studies indicate warm summer water temperatures around the Chatham Islands might have persisted during the LGM (Nelson *et al.* 2000, Fenner *et al.* 1992). Hoarau *et al.* (2007) estimated a mutation rate of 2% to 3.4%/Myr for this mitochondrial spacer in *Fucus*. While I would be cautious in applying this rate directly to *C. maschalocarpum*, a much faster mutation rate would be required if four endemic haplotypes arose following post-LGM recolonisation.

The pattern of local population differentiation, but with specific pathways for gene flow between distant populations, overlaid on a historical pattern of retreat and recolonisation is consistent with the intermittent and stochastic dynamics of long distance dispersal by floating (Thiel & Haye 2006), and might be a general pattern for buoyant algae (Fraser *et al.* 2009) and organisms rafting on these algae. An important finding is that connectivity between distant populations can be high, even where local

connectivity is often low. This highlights the importance of sampling across species' ranges when investigating dispersal, as large scale processes driving long distance dispersal might not be apparent in studies at local scales.

2.6 Acknowledgements

I thank Wendy Nelson, Kate Neill, Tracy Farr, Nick Shears, Peter Martin, Erasmo Macaya, Fiona Hodge and Chatham Islands Department of Conservation staff for help with collections, and Peter Ritchie for help with early drafts of the manuscript. Dr Martin Thiel provided many useful comments. I also thank the New Zealand Department of Conservation for funding assistance.

2.7 References

- ADAMS, N. M. (1994). *Seaweeds of New Zealand: An illustrated guide*. Christchurch, New Zealand: Canterbury University Press.
- ADEY, W. H. & STENECK, R. S. (2001). Thermogeography over time creates biogeographic regions: A temperature/space/time-integrated model and an abundance weighted test for benthic marine algae. *Journal of Phycology*, 37, 677–698.
- APTE, S. & GARDNER, J. P. A. (2002). Population genetic variation in the New Zealand greenshell mussel, *Perna canaliculus*: SSCP and RFLP analyses of mitochondrial DNA. *Molecular Ecology*, 11, 1617–1628.
- AUSTERLITZ, F., MARIETTE, S., MACHON, N., GOUYON P-H. (2000). Effects of colonization processes on genetic diversity: Differences between annual plants and tree species. *Genetics*, 154, 1309–1321.
- AYRES K. L. & WATERS, J. M. (2005). Marine biogeographic disjunction in central New Zealand. *Marine Biology*, 147, 1045–1052.
- BARROWS, T. T. & JUGGINS, S. (2005). Sea-surface temperatures around the Australian margin and Indian Ocean during the Last Glacial Maximum, *Quaternary Science Review*, 24, 1017–1047.
- BASSAM, B. J., CAETANO-ANOLLES, G. & GRESSHOFF, P. M. (1991). Fast and sensitive silver staining of DNA in polyacrylamide gels. *Analytical Biochemistry*, 196, 80–83.
- BOCKELMANN, A. C., REUSCH, T. B. H., BIJLSMA, R. & BAKKER, J. P. (2003). Habitat differentiation vs. isolation by distance: The genetic population structure of *Elymus athericus* in European salt marshes. *Molecular Ecology*, 12, 505–515.
- BOHONAK, A. J. (1999). Dispersal, gene flow and population structure. *Quarterly Review of Biology*, 74, 21–45.
- CHAPMAN, A. R. O. (1995). Functional ecology of fucoid algae: Twenty-three years of progress. *Phycologia*, 34, 1–32.
- CHISWELL, S. H. & BOOTH, J. D. (2008). Sources and sinks of larval settlement in *Jasus edwardsii* around New Zealand: Where do larvae come from and where do they go? *Marine Ecology Progress Series*, 354, 201–217.
- CLAYTON, M. N. (1992). Propagules of marine macroalgae: Structure and development. *European Journal of Phycology*, 27, 219–232.
- CLEMENT, M., POSADA, D., CRANDALL, K. A. (2000). TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, 9, 1657–1659.

- COLEMAN, M. A. & BRAWLEY, S. H. (2005a). Spatial and temporal variability in dispersal and population genetic structure of a rockpool alga. *Marine Ecology Progress Series*, 300, 63–77.
- COLEMAN, M. A. & BRAWLEY, S. H. (2005b). Are life history characteristics good predictors of genetic diversity and structure? A case study of the intertidal alga *Fucus spiralis* (Heterokontophyta; Phaeophyceae). *Journal of Phycology*, 41, 753–762.
- COLEMAN, M. A. & KELAHER, B. P. (2009). Connectivity among fragmented populations of a habitat-forming alga, *Phyllospora comosa* (Phaeophyceae, Fucales) on an urbanised coast. *Marine Ecology Progress Series*, 381, 63–70.
- COLSON, I. & HUGHES, R. N. (2004). Rapid recovery of genetic diversity of dogwhelk (*Nucella lapillus* L.) populations after local extinction and recolonization contradicts predictions from life-history characteristics. *Molecular Ecology*, 13, 2223–2233.
- COYER, J. A., PETERS, A. F., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2002). Hybridization of the marine seaweeds *Fucus serratus* and *Fucus evanescens* (Heterokontophyta: Phaeophyceae) in a 100-year old zone of secondary contact. *Proceedings of the Royal Society of London B Series*, 269, 1829–1834.
- COYER, J. A., PETERS, A. F., STAM, W. T. & OLSEN, J. L. (2003). Post ice-age recolonization and differentiation of *Fucus serratus* L. (Phaeophyceae; Fucaceae) populations in Northern Europe. *Molecular Ecology*, 12, 1817–1829.
- DELFT, E. M. (1939). Studies in the Fucales of New Zealand I. The genus *Carpophyllum* Grev. *Journal of Botany, London*, 77, 129–138.
- DETHIER, M. N., McDONALD, K. & STRATHMANN, R. R. (2003). Colonization and connectivity of habitat patches for coastal marine species distant from source populations. *Conservation Biology*, 17, 1024–1035.
- DEYSHER, L. & NORTON, T. A. (1982). Dispersal and colonisation of *Sargassum muticum* (Yendo) Fensholt. *Journal of Experimental Marine Biology and Ecology*, 56, 179–195.
- DROMGOOLE, F. I. (1973). *A contribution to the biology of the genus Carpophyllum Grev.* Unpublished PhD. thesis. University of Auckland, Auckland, New Zealand.
- DUPANLOUP, I., SCHNEIDER, S. & EXCOFFIER, L. (2002). A simulated annealing approach to define the genetic structure of populations. *Molecular Ecology*, 11, 2571–2581.
- ECKERT, C. G., SAMIS, K. E. & LOUGHEED, S. C. (2008). Genetic variation across species' geographical ranges: the central-marginal hypothesis and beyond. *Molecular Ecology*, 17, 1170–1188.
- ENGEL, C. R., DAGUIN, C. & SERRÃO, E. (2005). Genetic entitries and mating systems in hermaphroditic *Fucus spiralis* and its close dioecious relative *F. vesiculosus* (Fucaceae, Phaeophyceae). *Molecular Ecology*, 14, 2033–2046.
- EXCOFFIER, L. (2004). Patterns of DNA sequence diversity and genetic structure after a range expansion: Lessons from the infinite island model. *Molecular Ecology*, 13, 853–864.
- EXCOFFIER, L., LAVAL, G. & SCHNEIDER, S. (2005). ARLEQUIN, version 3.0: An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online*, 1, 47–50.
- FENNER, J., CARTER, L. & STEWART, R. (1992). Late Quaternary paleoclimate and paleoceanographic change over northern Chatham Rise, New Zealand. *Marine Geology*, 108, 383–404.
- FRASER, C. L., NIKULA, R., SPENCER, H. G. & WATERS, J. M. (2009). Kelp genes reveal effects of subantarctic sea ice during the Last Glacial Maximum. *Proceedings of the National Academy of Sciences of the U.S.A.*, 106, 3249–3253.
- FU, X. Y. (1997). Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics*, 147, 915–925.
- FU, X. Y. & LI, W. H. (1993). Statistical tests of neutrality of mutations. *Genetics*, 133, 693–709.
- GOLDSTIEN, S. J., GEMMELL, N. J. & SCHIEL, D. R. (2006). Comparative phylogeography of coastal limpets across a marine disjunction in New Zealand. *Molecular Ecology*, 15, 3259–3268.

- GOLDSTIEN, S. J., GEMMELL, N. J. & SCHIEL, D. R. (2009). Colonisation and connectivity by intertidal limpets among New Zealand, Chatham and Sub-Antarctic Islands. I. Genetic connections. *Marine Ecology Progress Series*, 388, 111–119.
- HAY, B. & GRANT, C. (2003). *Marine Resources in Tai Tokerau*. Auckland, New Zealand: James Henare Maori Research Centre, University of Auckland.
- HEATH, R. A. (1985). A review of the physical oceanography of the seas around New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 19, 79–124.
- HERNÁNDEZ-CARMONA, G., HUGHES, B. & GRAHAM, M. H. (2006). Reproductive longevity of drifting kelp *Macrocystis pyrifera* (Phaeophyceae) in Monterey Bay, USA. *Journal of Phycology*, 42, 1199–1207.
- HEWITT, G. M. (1996). Some genetic consequences of ice ages, and their role in divergence and speciation. *Biological Journal of the Linnean Society*, 58, 247–276.
- HEWITT, G. M. (1999). Post-glacial re-colonisation of European biota. *Biological Journal of the Linnean Society*, 68, 87–112.
- HICKEY, A. J. R., LAVERY, S. D., HANNAN, D. A., BAKER, C. S. & CLEMENTS, K. D. (2009). New Zealand triplefin fishes (family Tripterygiidae): contrasting population structure and mtDNA diversity within a marine species flock. *Molecular Ecology*, 18, 680–696.
- HIGGINS, S. I., NATHAN, R. & CAIN, M. L. (2003). Are long distance dispersal events in plants usually caused by non-standard means of dispersal? *Ecology*, 84, 1945–1956.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- HODGE, F., BUCHANAN, J. & ZUCCARELLO, G. C. (2010). Hybridization between the endemic brown algae *Carpophyllum maschalocarpum* and *C. angustifolium* (Fuciales): Genetic and morphological evidence. *Phycological Research*, 58, 239–247.
- KENDRICK, G. A. & WALKER, D. I. (1991). Dispersal distances for propagules of *Sargassum spinuligerum* (Sargassaceae, Phaeophyta) measured directly by vital staining and venturi suction sampling. *Marine Ecology Progress Series*, 79, 133–138.
- KENDRICK, G. A. & WALKER, D. I. (1995). Dispersal of propagules of *Sargassum* spp. (Sargassaceae: Phaeophyta): Observations of local patterns and consequences for recruitment and population structure. *Journal of Experimental Marine Biology and Ecology*, 192, 273–288.
- KIM, K. Y., O'LEARY, S. J. & GARBARY, D. J. (1997). Artificial hybridization between *Ascophyllum nodosum* and *Fucus vesiculosus* (Phaeophyceae) from Nova Scotia, Canada. *Canadian Journal of Botany*, 75, 1133–1136.
- KINGSFORD, M. J. (1992). Drift algae and small fish in coastal waters of northeastern New Zealand. *Marine Ecology Progress Series*, 80, 41–55.
- KINLAN, B. P., GAINES, S. D. & LESTER, S. E. (2005). Propagule dispersal and the scales of marine community processes. *Diversity and Distributions*, 11, 139–148.
- KUMAR, S., TAMURA, K. & NEI, M. (2004). MEGA3: Integrated Software for Molecular Evolutionary Genetics Analysis and Sequence Alignment. *Briefings in Bioinformatics*, 5, 150–163.
- LEWIS, K. B., CARTER, L. & DAVEY, F. J. (1994). The opening of Cook Strait: Interglacial tidal scour and aligning basins at a subduction to transform plate edge. *Marine Geology*, 116, 293–312.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae *Nova Hedwigia*, 3, 129–350.
- LÜNING, K. (1990). *Seaweeds: Their environment, biogeography, and ecophysiology*. New York, Wiley.
- MACAYA, E., BOLTAÑA, S., BUSCHMANN, A., HINOJOSA, I., MACCHIAVELLO, J., VALDIVIA, N. ET AL. (2005). Presence of sporophylls in floating kelp rafts of *Macrocystis* spp. (Phaeophyceae) along the Chilean Pacific Coast. *Journal of Phycology*, 41, 913–922.

- MAGGS, C. A., CASTILHO, R., FOLTZ, D., HENZLER, C., JOLLY, M. T., KELLY, J. *ET AL.* (2008). Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology*, 89, S108–S122.
- MARKO, P. B. (2004). ‘What’s larvae got to do with it?’ Disparate patterns of post-glacial population structure in two benthic marine gastropods with identical dispersal potential. *Molecular Ecology*, 13, 597–611.
- MARKO, P. B., HOFFMAN, J. M., EMME, S. A., MCGOVERN, T. M., KEEVER, C. C. & COX, L. N. (2010). The ‘Expansion-Contraction’ model of Pleistocene biogeography: rocky shores suffer a sea change? *Molecular Ecology*, 19, 146–169.
- MCKENZIE, P. F. & BELLGROVE, A. (2008). Dispersal of *Hormosira banksii* (Phaeophyceae) via detached fragments: Reproductive viability and longevity. *Journal of Phycology*, 44, 1108–1115.
- MORTON, J. E. & MILLER M. C. (1968). *The New Zealand Sea Shore*. London-Auckland: Collins.
- MUHLIN, J. F., ENGEL, C. R., STESSEL, R., WEATHERBEE, R. A. & BRAWLEY, S. H. (2008). The influence of coastal topography, circulation patterns and rafting in structuring populations of an intertidal alga. *Molecular Ecology*, 17, 1198–1210.
- NAYLOR, M. (1954). The female conceptacle and young sporeling of the four New Zealand species of *Carpophyllum*. *Transactions and Proceedings of the Royal Society of New Zealand*, 82, 1–6.
- NELSON, C. S., HENDY, I. L., NEIL, H. L., HENDY, C. H. & WEAVER, P. P. E. (2000). Last glacial jetting of cold waters through the Subtropical Convergence zone in the Southwest Pacific off eastern New Zealand, and some geological implications. *Palaeogeography Palaeoclimatology Palaeoecology*, 156, 103–121.
- NELSON, W. A. (1994). Distribution of macroalgae in New Zealand – an archipelago in space and time. *Botanica Marina*, 37, 221–233.
- NELSON, W. A., ADAMS, N. M. & HAY, C. H. (1991). *Marine Algae of the Chatham Islands*. (National Museum of New Zealand Miscellaneous Series No. 23). Wellington, New Zealand: National Museum of New Zealand.
- NELSON, W. A., VILLOUTA, E., NEILL, K. F., WILLIAMS, G. C., ADAMS, N. M. & SLIVSGAARD, R. (2002). Marine macroalgae of Fiordland, New Zealand. *Tuhinga*, 13, 117–152.
- NORTON, T. A. (1992). Dispersal by macroalgae. *British Phycological Journal*, 27, 293–301.
- LOUDON-LE SECQ, M.-P., KLOAREG, B., LOISEAUX-DE GÖER, S. (2002). The mitochondrial genome of the brown alga *Laminaria digitata*: a comparative analysis. *European Journal of Phycology*, 37, 163–172.
- LOUDON-LE SECQ, M.-P., LOISEAUX-DE GÖER, S., STAM, W. T. & OLSEN, J. L. (2006). Complete mitochondrial genomes of the three brown algae (Heterokonta: Phaeophyceae) *Dictyota dichotoma*, *Fucus vesiculosus* and *Desmarestia viridis*. *Current Genetics*, 49, 47–58.
- PAKKER, H., BREEMAN, A. M., PRUD’HOMME VAN REINE, W. F., VAN OPPEN, M. J. H. & VAN DEN HOEK, C. (1996). Temperature responses of tropical to warm-temperate Atlantic seaweeds. I. Absence of ecotypic differentiation in amphi-Atlantic tropical Canary Islands species. *European Journal of Phycology*, 31, 123–132.
- PROVAN, J., WATTIER, R. A. & MAGGS, C. A. (2005). Phylogeographic analysis of the red seaweed *Palmaria palmata* reveals a Pleistocene marine glacial refugium in the English Channel. *Molecular Ecology*, 14, 793–803.
- RAY, N., CURRAT, M. & EXCOFFIER, L. (2003). Intra-deme molecular diversity in spatially expanding populations. *Molecular Biology and Evolution*, 20, 76–86.
- REED, D. C., AMSLER, C. D. & EBELING, A. W. (1992). Dispersal in kelps – Factors affecting spore swimming and competence. *Ecology*, 73, 1577–1585.
- REED, D. C., LAUR, D. R. & EBELING, A. W. (1998). Variation in algal dispersal and recruitment: The importance of episodic events. *Ecological Monographs*, 58, 321–335.
- RIDGWAY, N. M. (1960). Surface water movements in Hawke Bay, New Zealand. *New Zealand Journal of Geology and Geophysics*, 3, 253–261.

- RIDGWAY, N. M. (1962). Nearshore surface currents in southern Hawke Bay, New Zealand. *New Zealand Journal of Geology and Geophysics*, 5, 545–566.
- ROGERS, A. R. & HARPENDING, H. (1992). Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution*, 9, 552–569.
- ROSS, P. M., HOGG, I. D., PILDITCH, C. A. & LUNDQUIST, C. J. (2009). Phylogeography of New Zealand's coastal benthos. *New Zealand Journal of Marine and Freshwater Research*, 43, 1009–1027.
- SCHIEL, D. R., ANDREW, N. L. & FOSTER, M. S. (1995). The structure of subtidal algae and invertebrate assemblages at the Chatham Islands, New Zealand. *Marine Biology*, 123, 355–367.
- SCOTT, G. W. & HARDY, F. G. (1994). Observations of the occurrence of hybrids between two sympatric species of fucoid algae. *Cryptogamie, Algologie*, 15, 297–305.
- SHEARS, N. T. & BABCOCK, R. C. (2007). *Quantitative description of New Zealand's shallow sub-tidal reef communities*. (Science for Conservation 280). Wellington, New Zealand: New Zealand Department of Conservation.
- SHEARS, N. T., SMITH, F., BABCOCK, R. C., DUFFY, C. A. & VILLOUTA, E. (2008). Evaluation of biogeographic classification schemes for conservation planning: application to New Zealand's coastal marine environment. *Conservation Biology*, 22, 467–481.
- SMITH, S. D. A. (2002). Kelp rafts in the Southern Ocean. *Global Ecology & Biogeography*, 11, 67–69.
- SNYDER, T. P. & GOOCH, J. L. (1973). Genetic differentiation in *Littorina saxatilis* (Gastropoda). *Marine Biology*, 22, 177–182.
- SOTKA, E. E., WARES, J. P., BARTH, J. A., GROSEBERG, R. K. & PALUMBI, S. R. (2004). Strong genetic clines and geographical variation in gene flow in the rocky intertidal barnacle *Balanus glandula*. *Molecular Ecology*, 13, 2143–2156.
- SPONER, R. & ROY, M. S. (2002). Phylogeographic analysis of the brooding brittle star *Amphipholis squamata* (Echinodermata) along the coast of New Zealand reveals high cryptic genetic variation and cryptic dispersal potential. *Evolution*, 56, 1954–1967.
- STANTON, B. R. (1997). Sea level and sea surface temperature variability at the Chatham Islands, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 31, 525–536.
- STANTON, B. R., SUTTON, P. J. H. & CHISWELL, S. M. (1997). The East Auckland Current, 1994–95. *New Zealand Journal of Marine and Freshwater Research*, 31, 537–549.
- SUNNUCKS, P., WILSON, A. C. C., BEHEREGARAY, L. B., ZENGER, K., FRENCH, J. & TAYLOR, A. C. (2000). SSCP is not so difficult: the application and utility of single-stranded conformation polymorphism in evolutionary biology and molecular biology. *Molecular Ecology*, 9, 1699–1710.
- TAJIMA, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123, 585–595.
- TATARENKOV, A., JÖNSSON, R. B., KAUTSKY, L. & JOHANNESSON, K. (2007). Genetic structure in populations of *Fucus vesiculosus* (Phaeophyceae) over spatial scales from 10 m to 800 km. *Journal of Phycology*, 43, 675–685.
- THIEL, M. & GUTOW, L. (2005a). The ecology of rafting in the marine environment. II. The rafting organisms and community. *Oceanography and Marine Biology*, 43, 279–418.
- THIEL, M. & GUTOW, L. (2005b). The ecology of rafting in the marine environment. I. The floating substrata. *Oceanography and Marine Biology*, 42, 181–263.
- THIEL, M. & HAYE, P. A. (2006). The ecology of rafting in the marine environment. III. Biogeographical and evolutionary consequences. *Oceanography and Marine Biology*, 44, 323–429.
- VAN DEN HOEK, C. (1987). The possible significance for long range dispersal for the biogeography of seaweeds. *Helgoländer wissenschaftliche Meeresuntersuchungen*, 41, 261–72.
- WARES, J. P. & CUNNINGHAM, C. W. (2001). Phylogeography and historical ecology of the North Atlantic intertidal. *Evolution*, 55, 2455–2469.

- WATERS, J. M. & ROY, M. S. (2004). Phylogeography of a high dispersal New Zealand sea-star: does upwelling block gene flow? *Molecular Ecology*, *13*, 2797–2806.
- ZUCCARELLO, G. C. & LOKHORST, G. M. (2005). Molecular phylogeny of the genus *Tribonema* (Xanthophyceae) using *rbcL* gene sequence data: monophyly of morphologically simple algal species. *Phycologia*, *44*, 384–392.
- ZUCCARELLO, G. C., WEST, J. A., KAMIYA, M. & KING, R. J. (1999). A rapid method to score plastid haplotypes in red seaweeds and its use in determining parental inheritance of plastids in the red alga *Bostrychia* (Ceramiales). *Hydrobiologia*, *401*, 207–214.

Chapter 3
Patterns of hybridisation in *Carpophyllum*
(Phaeophyceae, Fucales)

3. 1 Abstract

Hybrid zones can vary in time and space, even where the same species are involved. *Carpophyllum maschalocarpum* (Phaeophyceae, Fucales) and *C. angustifolium* hybridise in sites in north-eastern New Zealand where the two species are broadly sympatric, with *C. angustifolium* limited to more wave-exposed habitats. Patterns of hybridisation were investigated by sampling across this zone of sympatry, from the northern tip of New Zealand to near the south-eastern limit of *C. angustifolium*. Specimens were assigned to species using morphological criteria. Hybrids have an intermediate morphology, specifically, a stipe width intermediate between the two parental species. Ribotypes and haplotypes were determined for ITS2 and a mtDNA spacer. In the southern range of *C. angustifolium*, most hybrids have heterozygous ITS2 and ribotypes associated with both parent species, and are interpreted as F1 hybrids, with a low frequency of backcrossing. Mitochondrial haplotypes were species specific in southern populations, but one haplotype found only in *C. angustifolium* and hybrids clustered with haplotypes found only in *C. maschalocarpum*, suggesting previous mitochondrial introgression in *C. angustifolium*. In the northern range of *C. angustifolium*, shared ITS ribotypes suggest backcrosses are more common, and all specimens in some populations resemble hybrids with parent plants apparently absent. Unlike *Fucus*, where asymmetrical hybridisation results from different mating systems between sister species, *Carpophyllum* is dioecious and either species can act as the maternal parent in the production of hybrids. High frequencies of *C. angustifolium*-associated mtDNA haplotypes in hybrids probably arise from imbalances in population sizes of parent species.

Abbreviations:

CTAB, Cetyltrimethylammonium Bromide; ITS, Internal Transcribed Spacer; SSCP, Single Stranded Conformational Polymorphism; TBE, Tris-Borate-EDTA.

3.2 Introduction

Natural interspecific hybridisation events are natural experiments that can provide insight into evolutionary processes that lead to divergence and speciation (Hewitt 1988). Hybridisation can enable gene flow between lineages (Arnold 2006, Mallet 2005), reversing species divergence, and can produce novel phenotypes and initiate new species (Riesberg 1997, 2006). Gametic contact between potentially hybridising species is usually restricted in space creating hybrid zones, where genetically distinct populations overlap (Barton & Hewitt 1985, 1989, Arnold 1993, Gardner 1997). Hybrid zones can arise from secondary contact following natural range expansion (Hewitt 2001, Neiva *et al.* 2010) or anthropogenic introductions (Coyer *et al.* 2002a, 2006b, Rhymer & Simberloff 1996). Early studies assumed hybrid zones were an ephemeral stage in species divergence, occurring when development of reproductive barriers was incomplete (Howard *et al.* 2003). Subsequent theoretical work suggested hybrid zones might be stable, either maintained by a balance of selection against hybrids and dispersal into the zone (Bigelow 1965, Barton & Hewitt 1985), or by greater fitness of hybrids in some habitats (Moore 1977). Recently, a more complex picture has emerged, with variable hybrid genotypes (Arnold & Hodges 1995) interacting with patchy and dynamic environmental conditions to create a mosaic hybrid zone, with multiple potential evolutionary outcomes (Butlin *et al.* 2008, Jiggins & Mallet 2000, Schwenk *et al.* 2008). Consequently, all hybrid zones are not all alike, even where the same species are involved (Riginos & Cunningham 2005). In the marine environment, hybridisation is probably common (Gardner 1997) but has been extensively studied in only a few systems (Rawson & Hilbish 1998, Rawson *et al.* 1999, Coyer *et al.* 2004, Lewis 1996, Coyer *et al.* 2007). Fuclean brown algae are a useful system for studying hybridisation as they have a simple life history, with no alternation of generations and predominantly sexual reproduction (Chapman 1995). Various scenarios lead to hybridisation in *Fucus*: anthropogenic secondary contact (Coyer *et al.* 2002a, 2006b, 2007); marginal habitat (Wallace *et al.* 2004, Coyer *et al.* 2006a); and natural sympatry (Billard *et al.* 2005, Engel *et al.* 2005, Neiva *et al.* 2010). Species integrity in *Fucus* appears to be maintained by selection against hybrids (Coyer *et al.* 2007) and contrasting mating systems (Billard *et al.* 2005, Engel *et al.* 2005), but some populations show extensive

introgression (Neiva *et al.* 2010), and possible incipient hybrid speciation (Coyer *et al.* 2010 in press).

In New Zealand, species of the endemic genus *Carpophyllum* Greville typically dominate the shallow rocky sub-tidal (Adams 1994, Shears & Babcock 2007). Hybridisation, inferred from observations of morphologically intermediate specimens where two species are in contact, has been reported between *Carpophyllum maschalocarpum* Turner (Greville) and the three other species in the genus (Lindauer *et al.* 1961, Dromgoole 1973). *Carpophyllum maschalocarpum* is a common species, growing in dense stands around the Chatham Islands and most of mainland New Zealand (Adams 1994). A sister species, *Carpophyllum angustifolium* J. Agardh, has a more restricted range, extending from the northern tip of New Zealand's North Island to just short of East Cape (Adams 1994). These species are broadly sympatric, but *C. angustifolium* is absent from sheltered areas and replaces *C. maschalocarpum* in sites exposed to strong wave action (Shears & Babcock 2007). Morphologically, they can be distinguished by their main axes, broad and flattened in *C. maschalocarpum* and narrow and terete in *C. angustifolium* (Adams 1994).

Hodge *et al.* (2010, Appendix 8) combined ribosomal DNA and morphological data to show that hybridisation occurred between *Carpophyllum maschalocarpum* and *C. angustifolium* J. Agardh from two populations in the eastern Bay of Plenty, New Zealand. Specimens were morphologically assigned to species, primarily by stipe width (stipe < 2mm = *C. angustifolium*, stipe 2–5 mm = hybrid, stipe >5mm = *C. maschalocarpum*). Parent species had homozygous ITS2, with one of two species-specific ribotypes. In contrast, ITS2 electropherograms from morphologically intermediate specimens showed a pattern of double peaks consistent with the presence of both ribotypes in each individual. A small number of individuals had morphological characters consistent with *C. angustifolium*, but had hybrid (heterozygous) ITS2 sequences. Hodge *et al.* (2010) suggested these were backcrosses, possibly indicating asymmetrical introgression.

Hybrids have also been reported from Auckland localities (Dromgoole 1973) and Shears & Babcock (2007) reported what they considered to be a slender form of *C. maschalocarpum* from sites in the far north of New Zealand. These specimens somewhat resembled *C. angustifolium* and were found in wave

exposed locations, but where the authors considered *C. angustifolium* was absent. As hybrids collected from the Bay of Plenty (Hodge *et al.* 2010) could be described as resembling “slender” *C. maschalocarpum*, I consider it likely that specimens observed by Shears & Babcock (2007) are also hybrids. Hay & Grant (2003) reported only *C. angustifolium* from the northern tip of the North Island, whereas Shears & Babcock (2004) reported only *C. maschalocarpum*. This situation is unsatisfactory and the taxonomic status of these populations needs to be clarified.

Here I investigate evidence for hybridisation across the range of *C. angustifolium*, by comparing populations from Northland and the Bay of Plenty, New Zealand. In addition to ribotyping by ITS2, a mitochondrial intergenic spacer (Hoarau *et al.* 2007) was used to determine the maternal species of putative hybrid specimens. The utility of this spacer was shown in an extensive survey of *C. maschalocarpum* and *C. angustifolium* (Chapter 2) where haplotypes formed three clusters, two associated with *C. angustifolium* and one associated with *C. maschalocarpum*. The aim of this study is to determine if other hybridising populations of *Carpophyllum* follow the same patterns as those investigated by Hodge *et al.* (2010), and how species boundaries are maintained in *Carpophyllum*.

3.3 Materials and Methods

Sampling

Carpophyllum maschalocarpum specimens were collected from around New Zealand, and *C. angustifolium* and putative *C. maschalocarpum* × *C. angustifolium* hybrids from northern New Zealand (Table 3.1 and Fig. 3.2). More intensive sampling of parent species and putative hybrids, identified by morphological characters, was done at four sites: the Poor Knights Islands (N=36) and Leigh (N=29) in Northland, and Maraehako Bay (N=75) and Otanga (N=66) in the eastern Bay of Plenty. In addition, morphologically intermediate thalli were collected at Wekarua (N=7) and Hooper Point (N=14), both Northland sites where parental forms were not found.

Thalli were collected haphazardly, at least 1 metre apart. Tips of thalli for DNA extraction were dried with silica gel. In addition, one or more typical thalli from each species/site were prepared as voucher specimens. *A priori* species assignments were made using morphological characters developed by Hodge *et al.*

(2010), primarily a wide (>5mm) flattened primary axis in *C. maschalocarpum* and a narrow terete axis in *C. angustifolium*. Putative hybrids had a flattened primary axis intermediate in width (2–5 mm).

Molecular methods

DNA was extracted from approximately 2–5 mg of dry tissue in a modified CTAB buffer following procedures described in Zuccarello & Lokhorst (2005), with the addition of 1% polyvinylpolypyrrolidone to the extraction buffer. The ITS2 rDNA region and a variable mitochondrial spacer region were amplified and screened for variation by SSCP and sequencing. Amplification and sequencing of ITS2 (448–450 bp) used KP5 (forward) and KG4 (reverse) primers (Lane *et al.* 2006). Two further primers, ITS86F (5'–ACAGCTTCGGGTTCGATCT–3') and ITS424R (5'–ACCGGTCTCTCTCCGGTATT–3'), were developed from preliminary *Carpophyllum* ITS2 sequences to amplify a 337–339 bp fragment for SSCP analyses. This shorter fragment produced clearer SSCP profiles while retaining all known variable positions.

A 222 bp mitochondrial DNA fragment, containing 78 bp of the 3' end of the 23S rRNA gene and 144 bp of spacer was amplified from Frasers Landing (Poor Knights Islands) and Maraehako Bay specimens with primers from Chapter 2. In addition, mtDNA sequences from Hooper Point and Wekarua specimens (Chapter 2) are included in the dataset.

Table 3.1. Exemplars of specimens sampled with ITS2 ribotypes and locality information.

ITS2 ribotype	Specimen field number	Herbarium number ⁵	Species	Site	Latitude/ Longitude	GenBank Accession number
B	A220	WELT ASG290#1	<i>Carpophyllum maschalocarpum</i>	Bland Bay, Bay of Islands, Northland	35.342°S 174.366°E	JF313148
G×N	A397	WELT A031674a	<i>Carpophyllum angustifolium</i>	Cavalli Passage, Northland	35.051°S 173.943°E	JF313160
G	A398	WELT A031674b	<i>Carpophyllum angustifolium</i>	Cavalli Passage, Northland	35.051°S 173.943°E	JF313161
I	A404	WELT A031673	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Cavalli Passage, Northland	35.051°S 173.943°E	JF313162
B	A561	PKI008	<i>Carpophyllum maschalocarpum</i>	Frasers Landing, Poor Knights Islands	35.483°S 174.745°E	JF313165
G	A563	WELT A031326	<i>Carpophyllum angustifolium</i>	Frasers Landing, Poor Knights Islands	35.483°S 174.745°E	JF313166

⁵ Specimens are awaiting accession at WELT. Herbaria number will be inserted when the accession process is complete.

Table 3.1 (Continued)

ITS2 ribotype	Specimen field number	Herbarium number	Species	Site	Latitude/ Longitude	GenBank Accession number
D	A885	-	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Hooper Point, Northland	34.416°S 172.854°E	JF313175
B	A905	-	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Hooper Point, Northland	34.416°S 172.854°E	JF313176
B	A725	-	<i>Carpophyllum maschalocarpum</i>	Otanga, Bay of Plenty	37.550°S 178.161°E	JF313168
A	A1000	-	<i>Carpophyllum maschalocarpum</i>	Otanga, Bay of Plenty	37.550°S 178.161°E	JF313178
B	A758	BOP339	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	JF313169
B	A766	-	<i>Carpophyllum maschalocarpum</i>	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	JF313170
B	A808	BOP350	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	JF313171

Table 3.1 (Continued)

ITS2 ribotype	Specimen field number	Herbarium number	Species	Site	Latitude/ Longitude	GenBank Accession number
B	A809	BOP351	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	JF313172
B	A639	-	<i>Carpophyllum maschalocarpum</i>	Matts Crack, Poor Knights Islands	35.476°S 174.741°E	JF313167
G	A378	WELT A031669	<i>Carpophyllum angustifolium</i>	Flat Island, Northland	34.984°S 173.865°E	JF313155
G	A379	WELT A031669	<i>Carpophyllum angustifolium</i>	Flat Island, Northland	34.984°S 173.865°E	JF313156
D	A423	WELT A031670	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	North Cape, Northland	34.416°S 173.053°E	JF313164
C	A935	WELT A031313	<i>Carpophyllum angustifolium</i>	Sailor's Grave, Coromandel	36.960°S 175.845°E	JF313177
I	A387	WELT A031671	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Stephenson Island, Northland	34.965°S 173.777°E	JF313157

Table 3.1 (Continued)

ITS2 ribotype	Specimen field number	Herbarium number	Species	Site	Latitude/ Longitude	GenBank Accession number
D	A388	WELT A031672	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Stephenson Island, Northland	34.965°S 173.777°E	JF313158
G	A390	WELT A031672	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Stephenson Island, Northland	34.965°S 173.777°E	JF313159
G	A408	WELT A031666	<i>Carpophyllum angustifolium</i>	Stephenson Island, Northland	34.965°S 173.777°E	JF313163
B	A341	-	<i>Carpophyllum maschalocarpum</i>	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	JF313149
B	A342	-	<i>Carpophyllum maschalocarpum</i>	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	JF313150
F	A345	-	<i>Carpophyllum angustifolium</i>	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	JF313151
F	A869	WELT A031320	<i>Carpophyllum maschalocarpum</i>	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	JF313173

Table 3.1 (Continued)

ITS2 ribotype	Specimen field number	Herbarium number	Species	Site	Latitude/ Longitude	GenBank Accession number
B	A870	-	<i>Carpophyllum angustifolium</i>	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	JF313174
H	A350	WELT A031668	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Wekarua, Northland	34.945°S 173.653°E	JF313152
D	A353	WELT A031667	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Wekarua, Northland	34.945°S 173.653°E	JF313153
E	A358	WELT A031667	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Wekarua, Northland	34.945°S 173.653°E	JF313154
B×G	A562	WELT A031326	<i>Carpophyllum angustifolium</i>	Fraser's Landing, Poor Knights Islands	35.483°S 174.745°E	-
H×L	A349	WELT A031320	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Waterfall Reef, Leigh, Northland	36.269°S 174.798°E	-
B×G	A810	BOP352	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	-

Table 3.1 (Continued)

ITS2 ribotype	Specimen field number	Herbarium number	Species	Site	Latitude/ Longitude	GenBank Accession number
B×G	A814	BOP356	<i>Carpophyllum maschalocarpum</i> × <i>C. angustifolium</i> hybrid	Maraehako Bay, Bay of Plenty	37.672°S 177.797°E	-

The PCR mix contained 1µl genomic DNA, 0.5 U Taq DNA polymerase (New England Biolabs), 1X Thermopol reaction buffer (NEB), 7.5 pmoles each primer, 200 nM dNTP, 5% DMSO and 0.01% BSA. All PCR amplifications used a touchdown protocol with an initial denaturation step of 94°C for 4 min, followed by 5 cycles of 1 min at 94°C, 1 min at 55°C (−1°C/cycle) and 1 min at 72°C, followed by 30 cycles of 1 min at 94°C, 1 min at 50°C and 1 min at 72°C, with a final extension of 72°C for 10 min. Amplified products were checked for length and yield on 1% agarose gels stained with ethidium bromide.

SSCP screening

3µl PCR product was mixed with 9 µL 98% formamide, 10 mM NaOH, 0.025% bromophenol blue, 0.025% xylene cyanol, denatured at 100°C for 5 min, then snap cooled on ice before loading. Gels were 225 mm long and 0.75 mm thick (BioRad D-Code System) and contained 10% Mutation Detection Acrylamide (Sigma A5934, Sigma Aldrich, St Louis, USA) for ITS2 or 9% 37.5:1 acrylamide/bis-acrylamide (Sigma Aldrich) for the mitochondrial spacer, in 0.5X TBE buffer with the addition of 10% glycerol. Electrophoresis was carried out for 16–18 hrs at 8W in 0.5X TBE buffer at 4°C. After electrophoresis, gels were silver stained following protocols in Bassam *et al.* (1991) and banding patterns scored by eye.

Sequences were obtained from random samples of each ITS2 ribotype or mitochondrial haplotype indicated by SSCP analyses, as well as any samples that showed ambiguous SSCP patterns. In addition, sequences were obtained from other populations where low sampling did not warrant use of SSCP (Table 3.1). PCR products were cleaned with ExoSAP-IT (USB, Cleveland, Ohio) enzymes and sequenced commercially (Macrogen Inc., Korea).

ITS2 Cloning

Four hybrid specimens were selected for cloning, two from Leigh and one each from Maraehako Bay and Otanga, to determine if both parental ribotypes could be obtained. The complete ITS1–5.8S–ITS2 cistron was amplified using KP1 and KG4 primers. Purified PCR product was cloned using the pGEM-T Easy vector kit (Promega Corporation, Madison, Wisconsin USA) following the manufacturer's protocols.

Transformed colonies were sub-cultured and plasmid preparations carried out by alkali lysis (Sambrook *et al.* 1989). Inserts were sequenced commercially using the pUC M13 forward and reverse primers.

Data analysis

Homozygous sequences were aligned using ClustalW in MEGA 3.1 (Kumar *et al.* 2004) and checked by eye (Appendix 2). A statistical parsimony map was inferred from ITS2 data using TCS 1.21 (Clement *et al.* 2000) from 83 sequences from parental specimens, and six cloned sequences from hybrid specimens.

Heterozygous ITS2 sequences were assumed to result from inheritance of only two ribotypes. Electropherograms were examined and assigned the most likely parental ribotypes required to produce the observed heterozygosity (Table 3.1, Fig. 3.2). Three heterozygous sequences could not be explained as progeny of sampled homozygous ribotypes. Parental ribotypes of these could be inferred from heterozygous sequences as these all differed from frequently sampled ribotypes by a single heterozygous position. These inferred parental ribotypes were included in the statistical parsimony analysis. Mitochondrial spacer haplotypes were all identical to haplotypes obtained in a previous study (Chapter 2). Haplotypes were assigned to species according to clusters inferred by statistical parsimony in that study.

3.4 Results

Morphology

Specimens from Bay of Plenty sites (Maraehako Bay and Otanga), Leigh and the two Poor Knights Islands sites were able to be assigned to three morphological classes (*C. maschalocarpum*, *C. angustifolium* or putative hybrids) based on stipe width (Hodge *et al.* 2010). All three classes were sampled at these sites. All specimens sampled from Wekarua and Hooper Point showed intermediate (2–5 mm) stipe width and no specimens from these sites were assigned to parental species (Table 3.1, Fig. 3.1)..

SSCP data

SSCP successfully distinguished mitochondrial spacer haplotypes in all populations. Variation in the longer ITS2 fragment was less well resolved by SSCP, but ribotypes

could be distinguished among populations of low ribotype diversity (Bay of Plenty, Poor Knights Islands). Assigning ribotypes from SSCP profiles was difficult in Northland populations where ribotype diversity was higher and differences were often single base changes only. I therefore rely on sequence data for most Northland populations.

ITS2

Nine ITS2 ribotypes were sampled by direct sequencing from homozygous specimens, three of these were common (Fig. 3.1). Ribotype G was associated with *C. angustifolium*, and was widespread, from Otanga to Stephenson Island. Ribotypes D and B were widespread and mainly associated with *C. maschalocarpum*. Ribotype D is predominantly northern, from Leigh to Hooper Point; ribotype B was found throughout New Zealand. Two ribotypes, A and E, were found in single specimens only. Other uncommon ribotypes appeared to be fairly local: Ribotype C was found only in *C. angustifolium* from Sailor's Grave; ribotype I from parental species and hybrids from North Cape to Cavalli Passage in Northland; ribotype H from *C. angustifolium* from Wekarua and one hybrid from the Poor Knights Islands; and ribotype F from *C. angustifolium* from Leigh and the Alderman Islands.

Four ribotypes (L, M, N and O, enclosed by dashed lines in Fig. 3.1) were not sampled in homozygous specimens but were obtained by inferring parental ribotypes from heterozygous specimens. Ribotypes M, N and O were each inferred from single specimens that differed from common ribotypes by one heterozygous position. Ribotype L was inferred from sequences from six heterozygous specimens from Leigh (Fig. 3.2) and two heterozygous specimens sequenced from North Cape (Table 3.1). Ribotype L was also found in cloned sequences from a hybrid specimen from Leigh (Table 3.2).

Statistical parsimony analysis did not cluster ribotypes with morphological species assignments (Fig. 3.1). Ribotypes G and H were predominantly found in *C. angustifolium* or hybrids and were separated by two substitutions, but ribotypes F, I and

Fig. 3.1. Statistical parsimony map inferred from 83 homozygous ITS2 sequences and six cloned sequences from *Carpophyllum maschalocarpum*, *C. angustifolium* and putative hybrids. Shaded circles are predominantly *C. angustifolium*; clear circles are predominantly *C. maschalocarpum*. Black circles represent ribotypes sampled only by cloning. Dashed circles indicate ribotypes not found in homozygotes but are parental ribotypes required to produce heterozygotic sequences sampled in some specimens. Area of circles is proportional to number of samples with that haplotype. Connections are made at 95% probability.

C were also predominantly found in *C. angustifolium* but were connected by single changes to ribotypes sampled extensively in *C. maschalocarpum*. The most common *C. maschalocarpum* ribotypes (B and D) were separated by three changes, and by ribotypes only sampled from cloned specimens.

Hybrids

With three exceptions, all specimens from Bay of Plenty and Poor Knights Islands sites that had intermediate morphologies also had heterozygous ITS2 (Fig. 3.2). In heterozygous specimens, double peaks were visible at particular sites in electropherograms. Both sequences could be read from these electropherograms. One ribotype was identical to ribotypes from *C. maschalocarpum* specimens from the same population, and the other ribotype was identical to ribotypes from *C. angustifolium* specimens from the same population. At Leigh, six hybrid specimens were heterozygous but one ribotype (ribotype L in Fig. 3.1) was not sampled in any non-hybrid specimens. Ribotype L differed from a common Northland *C. maschalocarpum*-associated ribotype by one substitution (Fig. 3.1). Three specimens with hybrid morphology from Maraehako Bay were homozygous for ribotype B, the ribotype found in all *C. maschalocarpum* specimens from this site. Specimens from Wekarua and Hooper Point were all assigned to hybrids based on morphology but nearly all specimens were homozygous for ITS2, except for two heterozygous specimens from Wekarua. All specimens sampled at these two sites had ribotypes associated with *C. maschalocarpum*, apart from two Wekarua specimens that had the *C. angustifolium*-associated Ribotype H (Fig. 3.1).

At Frasers Landing (Poor Knights Islands) and Leigh, two specimens morphologically assigned to *C. maschalocarpum* and seven specimens assigned to *C. angustifolium* also had heterozygous ITS2. These specimens all possessed the common *C. maschalocarpum*-associated ribotype B (Fig. 3.1) and *C. angustifolium*-associated ribotypes F or G.

ITS cloning

Eight to eleven clones containing PCR products were sequenced for each specimen selected for cloning (Table 3.2; Appendix 3). Single base changes that appear in one cloned sequence only, and were not detected in any directly sequenced sample, are

regarded as copy errors from PCR, and are excluded from further analysis. Multiple copies of expected parental ribotypes were obtained from all cloned specimens. In addition, two ribotypes (L and P, shown as black circles in Fig. 3.1) obtained from one specimen from Leigh (A347 in Table 3.2) were not sampled by direct sequencing of homozygous specimens. Both these ribotypes were predicted as unsampled ribotypes by statistical parsimony analysis using homozygous sequences, and ribotype L was inferred as one ITS2 variant in heterozygous specimens from Leigh and North Cape.

Table 3.2: Frequency of ribotypes from cloned ITS2 sequences from heterozygous specimens of *Carpophyllum*. Ribotype designation is shown in Fig.3.1.

Specimen/site	Ribotype						
	B	D	F	G	H	L	P
A980 Otanga	2	-	-	9	-	-	-
A810 Maraehako Bay	3	-	-	5	-	-	-
A868 Leigh	4	1	3	-	-	-	-
A347 Leigh	-	-	-	-	5	3	1

Table 3.3. Mitochondrial haplotype frequency from *Carpophyllum maschalocarpum*, *C. angustifolium* and hybrids from northern New Zealand. Haplotypes associated with *C. angustifolium* are shaded. Other haplotypes were associated with *C. maschalocarpum*. Haplotype numbers are from Chapter 2, Fig. 2.2.

Site	MorphoSpecies	Haplotype	N
Maraehako Bay	<i>C. angustifolium</i>	48	3
		23	17
	<i>C. maschalocarpum</i>	14	15
		16	2
		17	1
		18	1
		19	2
		38	2
		71	1
		72	1
	Hybrids	14	3
		15	3
		16	3
		23	14
		48	3
Fraser's Landing	<i>C. angustifolium</i>	45	7
	<i>C. maschalocarpum</i>	21	6
	Hybrids	21	6
Hooper Point	Hybrids	21	10
		34	34
		65	1
		66	1
		67	1
Wekarua	Hybrids	30	7
		33	1
		47	7
		50	4
		67	1

Mitochondrial DNA

Mitochondrial DNA haplotypes associated with both *C. angustifolium* and *C. maschalocarpum* (Chapter 2) were found in morphologically intermediate specimens from the Bay of Plenty and Hooper Point (Table 3.3). At Frasers Landing (Poor Knights Islands) all hybrids had *C. maschalocarpum*-associated mtDNA haplotypes. At all other sites the ratios of *C. angustifolium*-associated haplotypes:*C. maschalocarpum*-associated haplotypes were > 0.5 . Departure from a 0.5 ratio was significant at Hooper Point (Binomial Test, $P = 0.01$) and Wekarua ($P < 0.01$).

Morphologically intermediate specimens from the most northern sites (Wekarua and Hooper Point) had predominantly *C. maschalocarpum*-associated ITS2 ribotypes but *C. angustifolium*-associated mtDNA haplotypes. At Wekarua, five out of seven specimens sequenced had *C. maschalocarpum*-associated ITS2 ribotypes, but all specimens had *C. angustifolium*-associated mtDNA haplotypes. At Hooper Point, all 14 specimens sequenced showed *C. maschalocarpum*-associated ITS2 ribotypes, but ten of these had *C. angustifolium*-associated mtDNA haplotypes.

3.5 Discussion

These data suggest hybridisation with *C. maschalocarpum* is occurring across the range of *C. angustifolium*, but patterns of hybridisation differ between localities. Three patterns are evident: (1) *Carpophyllum* populations from the Bay of Plenty and the Poor Knights Islands show strong congruence between intermediate morphology and ITS heterozygosity. I interpret this as interspecies crossing resulting in morphologically distinct hybrids with few backcrosses. (2) At Leigh, specimens with intermediate morphology and heterozygous ITS2 were also present, but four (out of ten) specimens assigned morphologically to *C. angustifolium* had heterozygous ITS2, with one ribotype inferred from electropherograms associated with *C. maschalocarpum*. These data suggest backcrosses between hybrids and *C. angustifolium*, resulting in *C. maschalocarpum*-associated ITS2 ribotypes being found in specimens resembling *C. angustifolium*. (3) At Hooper Point and Wekarua, all specimens were morphologically similar to hybrids found in other populations, and parental species were not seen, and are assumed to be rare or absent. Specimens were predominantly homozygous at ITS2, with *C. maschalocarpum*-associated ribotypes, but the majority of samples had *C.*

Fig. 3.2. Geographical patterns of ITS2 ribotypes in *Carpophyllum maschalocarpum*, *C. angustifolium* and hybrids from northern New Zealand populations. Pie charts show proportion of ribotypes sampled at each site.

angustifolium mtDNA haplotypes. These data suggest a population with hybrid origins, but where ITS homogenisation has taken place.

In the Bay of Plenty, hybrids are found predominantly in areas of moderate wave exposure, where *C. maschalocarpum* begins to be replaced by the more wave exposure-tolerant *C. angustifolium* (Hodge 2009). These hybrid zones are consistent with Barton & Hewitt's (1985) tension model, where hybrids are selected against relative to parents, but hybrid zones are maintained by migration into the zone. Hybrid populations at Hooper Point and Wekarua appear to persist without obvious sources of immigrants, but also without competition from parent populations. Either local environmental conditions favour hybrid morphologies (Moore 1977) or hybrid populations have established through founder effects (Neiva *et al.* 2010). Further research is needed to clarify this.

These ITS2 data from the Bay of Plenty are broadly in agreement with Hodge *et al.* (2010) who sampled *Carpophyllum* from Maraehako Bay and the adjoining Uncles Bay. I extended ITS2 data by including a third, more distant site (Otanga, near Lottin Point). Two ITS2 ribotypes dominated Bay of Plenty sites, although an additional ITS2 ribotype (A) was sampled in one Otanga specimen. With the exception of three specimens from Maraehako Bay, morphology is a reliable predictor of ITS2 ribotype in these populations: in homozygous specimens one ribotype was associated with *C. angustifolium*, the other with *C. maschalocarpum*. Specimens with intermediate morphology from these sites were heterozygous, with both parental ribotypes, and could be interpreted as F1 hybrids.

Hodge *et al.* (2010) found six specimens that were assigned to *C. angustifolium* on the basis of morphology, but had heterozygous ITS2, and suggested these were backcrosses between F1 hybrids and *C. angustifolium*. No specimens with *C. maschalocarpum* morphology were found with *C. angustifolium*-associated ribotypes, suggesting backcrossing might be asymmetrical. No thalli with heterozygous ribotypes and parental morphology were found in the Bay of Plenty in this study. Instead, three specimens had both an intermediate hybrid morphology and homozygous ITS2, with the *C. maschalocarpum*-associated ribotype B. I interpret these as F2 or later progeny of hybrid parents, where ITS2 DNA has homogenised to a single ribotype. ITS homogenisation, where a single, randomly selected ITS variant is fixed within the

individual (Elder & Turner 1995), is expected to remove the signature of hybridisation. However, the rate of homogenisation in brown algae is not known. Curiously, these three specimens had a unique mitochondrial spacer haplotype (haplotype 15 in Chapter 2), which was not sampled in any other population in this study (or in Chapter 2). Possible explanations are that this is a quirk of stochastic sampling and this haplotype is more widespread, or these might represent an incipient hybrid lineage, such as is found in the low-shore variety of *Fucus spiralis* (Coyer *et al.* 2010).

Biparental inheritance of ribosomal DNA provides a useful tool for the detection of hybridisation (Coleman 2002, Carine *et al.* 2007), but its value is limited (Álvarez & Wendel 2003) because of homogenisation. ITS heterozygosity is expected in F1 hybrids, but the signature of backcrosses will disappear if F1 hybrids homogenise to the same ribotype as the parental species involved in the backcross. Hence backcrosses will be underestimated by this method, especially where ribotype diversity is low, such as the Bay of Plenty. In addition, direct sequencing might not reveal the intra-individual diversity of ITS where homogenisation is incomplete and some ribotypes are retained in low copy number. One solution is to use cloning to estimate intra-individual ITS diversity. In my data, Ribotype L was predicted by statistical parsimony analysis as an intermediate haplotype, and inferred from heterozygous specimens, but was only confirmed in one cloned specimen, from Leigh. This specimen also provided a single cloned sequence of ribotype P, also predicted by statistical parsimony analysis. PCR artifacts can be a problem during cloning, but it is unlikely that copy errors would produce five copies of predicted ribotypes. This suggests backcrossing might be more frequent in Bay of Plenty populations, but other markers, such as microsatellites need to be developed to accurately estimate rates.

Patterns of hybridisation in Bay of Plenty populations are relatively straightforward. A more complicated pattern was found in Northland sites. At Frasers Landing and Leigh, three and four specimens (out of seven and ten samples, respectively) specimens had *C. angustifolium* morphology but heterozygous ITS2, with one ribotype that is usually associated with *C. maschalocarpum*. Two specimens from Leigh had *C. angustifolium* morphology and were homozygous, but had *C. maschalocarpum*-associated ribotypes (Fig. 3.1). Conversely, only a single specimen from each site had *Carpophyllum maschalocarpum* morphology and heterozygous ITS2,

and only one *C. maschalocarpum* specimen (from Leigh) was homozygous with a *C. angustifolium*-associated ribotype. I interpret this mis-match between morphology and ITS sequence(s) as backcrosses. Backcrosses mostly had *C. angustifolium* morphology, a result also found by Hodge *et al.* (2010), suggesting hybrids are more likely to backcross with *C. angustifolium*.

Maternal mitochondrial inheritance seems to be ubiquitous in oogamous algae, such as the Fucales, although rare heteroplasmy does occur (Coyer *et al.* 2002b, Coyer *et al.* 2004, Hoarau *et al.* 2009). There was no evidence of heteroplasmy in my samples, and I consider mitochondrial data to reflect maternal lineages. My mitochondrial data show that either *Carpophyllum* species can act as the maternal parent in hybridisation events, but in all populations except the Poor Knights Islands, hybrids had a higher frequency of *C. angustifolium* mtDNA (Table 3.3). The simplest explanation for this is a greater frequency of motile *C. maschalocarpum* male gametes (sperm) available for contact with immotile *C. angustifolium* female gametes (eggs). *Carpophyllum angustifolium* is usually restricted to the most wave exposed areas, often on the exposed parts of headlands, whereas *C. maschalocarpum* occupies a wide zone, from moderately exposed to moderately sheltered areas and is more abundant at most sites (Shears & Babcock 2007). This should result in a greater likelihood of *C. maschalocarpum* sperm being carried into stands of *C. angustifolium* than *vice versa*, and a greater potential for hybrids with *C. angustifolium* mtDNA. In contrast, hybrids from Frasers Landing had exclusively *C. maschalocarpum* mtDNA. This pattern might arise stochastically from low sampling at this site, but in an extensive survey of macroalgal abundance in New Zealand, the Poor Knights Islands were the only locality where *C. angustifolium* occurred more frequently than *C. maschalocarpum* (Shears & Babcock 2007). This might reverse the usual imbalance in gamete availability.

Haplotype 23 was common in *C. angustifolium* and hybrids from Maraehako Bay (Table 3.3). This haplotype was not sampled in *C. maschalocarpum* in this study or a previous study (Chapter 2), so I consider it indicative of a *C. angustifolium* maternal parent. However, statistical parsimony analysis in Chapter 2 places haplotype 23 in a cluster with *C. maschalocarpum* haplotypes, and just one change from a common haplotype found in *C. maschalocarpum* from Northland and the east coast of New Zealand's North Island. This suggests this haplotype or its ancestor has introgressed into

C. angustifolium populations in the past, and is now either extinct or rare in *C. maschalocarpum*. Founder effects might account for its high frequency at Maraehako Bay. A similar pattern is found in Europe, where a mitochondrial haplotype associated with *F. vesiculosus* is found in *F. ceranoides* throughout the latter species' northern range. This extensive introgression is believed to result from repeated founder effects (Neiva *et al.* 2010)

The high frequency of *C. angustifolium*-associated mtDNA haplotypes but mostly *C. maschalocarpum*-type ITS ribotypes in morphologically intermediate specimens from the most northern populations of Wekarua and Hooper Point suggest that these populations have hybrid origin, rather than being a wave-exposed ecotype of *C. maschalocarpum* as suggested by Shears & Babcock (2007). As no parent specimens were found at these sites, they appear to be self-sustaining populations of hybrid origin that have undergone ITS homogenisation, but retain hybrid morphology.

Mitochondrial spacer haplotypes are usually diagnostic for *Carpophyllum* species (Chapter 2). In contrast, only some ITS2 ribotypes had strong species associations, and most had exceptions (Table 3.1, Fig. 3.1). Most mitochondrial haplotypes also formed species-specific clusters (Chapter 2), whereas ITS2 ribotypes did not. It is not clear from these data whether ITS paraphyly is a consequence of gene transfer by hybridisation, or if this results from incomplete lineage sorting, with ancestral polymorphisms retained in both species. These results agree with the prediction that lineage sorting will proceed at a faster rate in mitochondrial DNA compared to nuclear DNA, due to a smaller effective population size in the former (Avice 2000). However, this prediction is probabilistic and cannot be applied to individual lineages with certainty (Hudson & Turelli 2003).

Carpophyllum populations in the Bay of Plenty are strongly segregated by habitat (Hodge 2009), with *C. maschalocarpum* occupying relatively sheltered embayments, but being replaced by *C. angustifolium* on the exposed headlands. Hybrids are predominantly found in a zone between the most exposed and sheltered areas. Species integrity in the Bay of Plenty might be maintained by segregation by habitat and limited gamete dispersal, with episodic selection against hybrids, possibly by physical removal of hybrids during storm events and overgrowth by *C. maschalocarpum* during calm periods. Habitat segregation is a factor in maintaining

species integrity in hybridising *Fucus* (Coyer *et al.* 2007, 2010). Northland sites appeared to have less sheltered habitat than Bay of Plenty sites. This might limit habitat segregation and increase the frequency of hybridisation events and backcrosses. Unlike *Fucus*, all *Carpophyllum* species are dioecious, which is expected to increase the frequency of hybridisation.

Three main findings arise from my data: (1) Hybridisation between *C. angustifolium* and *C. maschalocarpum* is widespread; (2) Patterns of hybridisation differ across the range of *C. angustifolium*; and (3) Initial hybridisation events appear symmetrical, but limited data suggests backcrosses are more likely between hybrids and *C. angustifolium*. Further ecological research and additional genetic markers are needed to determine if hybrids are under selection and if this varies between sites. While mitochondrial introgression from *C. angustifolium* into *C. maschalocarpum* appears rare (Chapter 2), the high frequency of haplotype 23 in *C. angustifolium* from Maraehako Bay suggests introgression in *C. angustifolium* might be more common. Further mitochondrial DNA data for *C. angustifolium* populations is needed to elucidate this pattern.

3.6 Acknowledgements

I thank Nick Shears, Wendy Nelson, Franz Smith, Kate Neill, Tracy Farr, Fiona Hodge and Peter Martin for assistance with collections.

3.7 References

- ADAMS, N. M. (1994). *Seaweeds of New Zealand: An illustrated guide*. Christchurch, New Zealand: Canterbury University Press.
- ÁLVAREZ, I. & WENDEL, J. F. (2003). Ribosomal ITS sequences and plant phylogenetic inference. *Molecular Phylogenetics and Evolution*, 29, 417–434.
- ARNOLD, J. (1993). Cytonuclear disequilibria in hybrid zones. *Annual Review of Ecology and Systematics*, 24, 521–554.
- ARNOLD, M. L. (2006). *Evolution through Genetic Exchange*. Oxford, United Kingdom: Oxford University Press.
- ARNOLD, M. L. & HODGES, S. A. (1995). Are natural hybrids fit or unfit relative to their parents? *Trends in Ecology and Evolution*, 10, 67–71.
- AVISE, J. C. (2000). *Phylogeography: The History and Formation of Species*. Cambridge MA: Harvard University Press.
- BARTON, N. H. & HEWITT, G. M. (1985). Analysis of hybrid zones. *Annual Reviews of Ecology and Systematics*, 16, 113–148.
- BARTON, N. H. & HEWITT, G. M. (1989). Adaption, speciation and hybrid zones. *Nature*, 341, 497–503.

- BASSAM, B. J., CAETANO-ANOLLES, G. & GRESSHOFF, P. M. (1991). Fast and sensitive silver staining of DNA in polyacrylamide gels. *Analytical Biochemistry*, 196, 80–83.
- BIGELOW, R. S. (1965). Hybrid zones and reproductive isolation. *Evolution*, 19, 449–458.
- BILLARD, E., DAGUIN, C., PEARSON, G., SERRÃO, E., ENGEL, C. & VALERO, M. (2005). Genetic isolation between three closely related taxa: *Fucus vesiculosus*, *F. spiralis* and *F. ceranoides* (Phaeophyceae). *Journal of Phycology*, 41, 900–905.
- BILLARD, E., SERRÃO, E., PEARSON, G. A., ENGEL, C. R., DESTOMBE, C. & VALERO, M. (2005). Analysis of sexual phenotype and prezygotic fertility in natural populations of *F. spiralis*, *F. vesiculosus* (Fucaceae, Phaeophyta) and their putative hybrids. *European Journal of Phycology*, 40, 397–407.
- BUTLIN, R. K., GALINDO, J. & GRAHAME, J. W. (2008). Sympatric, parapatric or allopatric: the most important way to classify speciation? *Philosophical Transactions of the Royal Society of London B Series*, 363, 2997–3007.
- CARINE, M. A., ROBBA, L., LITTLE, R., RUSSELL, S. & SANTOS GUERRA, A. (2007). Molecular and morphological evidence for hybridization between endemic Canary Island *Convolvulus*. *Botanical Journal of the Linnean Society*, 154, 187–204.
- CHAPMAN, A. R. O. (1995). Functional ecology of fucoid algae: Twenty-three years of progress. *Phycologia*, 34, 1–32.
- CLEMENT, M., POSADA, D. & CRANDALL, K. A. (2000). TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, 9, 1657–1659.
- COLEMAN, A. W. (2002). Comparison of *Eudorina/Pleodorina* ITS sequences of isolates from nature with those from experimental hybrids. *American Journal of Botany*, 89, 1523–1530.
- COYER, J. A., HOARAU, G., COSTA, J. F., HOGERDIJK, B., BILLARD, E., VALERO, M., PEARSON, G. A. & OLSEN, J. L. (2010). Evolution and species diversification within the intertidal brown macroalgae *Fucus spiralis*/*F. vesiculosus* species complex in the North Atlantic. *Molecular Phylogenetics and Evolution*, doi:10.1016/j.ympev.2010.11.015.
- COYER, J. A., HOARAU, G., PEARSON, G. A., SERRAO, E. A., STAM, W. T. & OLSEN, J. L. (2006a). Convergent adaptation to a marginal habitat by homoploid hybrids and polyploid ecads in the seaweed genus *Fucus*. *Biology Letters*, 2, 405–408.
- COYER, J. A., HOARAU, G., SKAGE, M., STAM, W. T. & OLSEN, J. L. (2006b). Origin of *Fucus serratus* (Heterokontophyta; Fucaceae) populations in Iceland and the Faroes: a microsatellite based assessment. *European Journal of Phycology*, 41, 235–246.
- COYER, J. A., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2004). Geographically specific heteroplasmy of mitochondrial DNA in the seaweed, *Fucus serratus* (Heterokontophyta: Phaeophyceae, Fucales). *Molecular Biology*, 13, 1323–1326.
- COYER, J. A., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2007). Hybridization and introgression in a mixed population of the intertidal seaweeds *Fucus evanescens* and *F. serratus*. *Journal of Evolutionary Biology*, 20, 2322–2333.
- COYER, J. A., PETERS, A. F., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2002a). Hybridization of the marine seaweeds *Fucus serratus* and *Fucus evanescens* (Heterokontophyta: Phaeophyceae) in a 100-year-old zone of secondary contact. *Proceedings of the Royal Society of London B Series*, 269, 1829–1834.
- COYER, J. A., PETERS, A. F., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2002b). Inheritance patterns of ITS1, chloroplasts and mitochondria in artificial hybrids of the seaweeds *Fucus serratus* and *F. evanescens* (Phaeophyceae). *European Journal of Phycology*, 37, 173–178.
- DROMGOOLE, F. I. (1973). *A contribution to the biology of the genus Carpophyllum Grev.* Unpublished PhD. thesis. Auckland, New Zealand: University of Auckland.
- ELDER, J. F. & TURNER, B. J. (1995). Concerted evolution of repetitive DNA sequences in eukaryotes. *The Quarterly Review of Biology*, 70, 297–320.

- ENGEL, C. R., DAGUIN, C. & SERRÃO, E. (2005). Genetic entities and mating systems in hermaphroditic *Fucus spiralis* and its close dioecious relative *F. vesiculosus* (Fucaceae, Phaeophyceae). *Molecular Ecology*, 14, 2033–2046.
- GARDNER, J. P. A. (1997). Hybridization in the Sea. *Advances in Marine Biology*, 31, 1–78.
- HAY, B. & GRANT, C. (2003). *Marine Resources in Tai Tokerau: Prepared for the James Henare Māori Research Centre's Capacity Building for Sustainable Māori Development research programme*. Auckland, New Zealand: James Henare Māori Research Centre.
- HEWITT, G. M. (1988). Hybrid zones – natural laboratories for evolutionary studies. *Trends in Ecology and Evolution*, 3, 158–167.
- HEWITT, G. M. (2001). Speciation, hybrid zones and phylogeography – or seeing genes in space and time. *Molecular Ecology*, 10, 537–549.
- HOARAU, G., COYER, J. A. & OLSEN, J. L. (2009). Paternal leakage of mitochondrial DNA in a *Fucus* (Phaeophyceae) hybrid zone. *Journal of Phycology*, 45, 621–624.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- HODGE, F. J. (2009). *Hybridisation in the brown algae Carpophyllum: Investigating morphology, distribution and wave exposure*. Unpublished masters thesis. Wellington, New Zealand: Victoria University of Wellington.
- HODGE, F., BUCHANAN, J. & ZUCCARELLO, G. C. (2010). Hybridization between the endemic brown algae *Carpophyllum maschalocarpum* and *C. angustifolium* (Fucales): Genetic and morphological evidence. *Phycological Research*, 58, 239–247.
- HOWARD, D. J., BRITCH, S. C., BRASWELL, W. E. & MARSHALL, J. L. (2003). Evolution in hybrid zones. In Singh, R. K. & Uyenoyama, M. K. (Eds.), *The Evolution of Population Biology* (pp. 297–314). Cambridge, United Kingdom: Cambridge University Press.
- HUDSON, R. R. & TURELLI, M. (2003). Stochasticity overrules the “three-times rule”: Genetic drift, genetic draft, and coalescence times for nuclear loci versus mitochondrial DNA. *Evolution*, 57, 182.
- JIGGINS, C. D. & MALLET, J. (2000). Bimodal hybrid zones and speciation. *Trends in Ecology and Evolution*, 15, 250–255.
- KUMAR, S., TAMURA, K. & NEI, M. (2004). MEGA3: Integrated Software for Molecular Evolutionary Genetics Analysis and Sequence Alignment. *Briefings in Bioinformatics*, 5, 150–163.
- LANE, C. E., MAYES, C. M., DRUEHL, L. & SAUNDERS, G. W. (2006). A multi-gene molecular investigation of the kelp (Laminariales, Phaeophyceae) resolves competing phylogenetic hypotheses and supports substantial taxonomic re-organization. *Journal of Phycology*, 42, 493–512.
- LEWIS, R. J. (1996). Hybridization of brown algae: Compatibility and speciation. In Chaudhary, B. R. & Agrawal, S. B., *Cytology, Genetics and Molecular Biology of Algae* (pp. 275–289). Amsterdam: SPB Academic Publishing.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae. *Nova Hedwigia*, 3, 129–350.
- MALLET, J. (2005). Hybridization as an invasion of the genome. *Trends in Ecology and Evolution*, 20, 229–237.
- MOORE, W. S. (1977). An evaluation of narrow hybrid zones in vertebrates. *The Quarterly Review of Biology*, 52, 263–277.
- NEIVA, J., PEARSON, G. A., VALERO, M. & SERRÃO, E. A. (2010). Surfing the wave on a borrowed board: range expansion and spread of introgressed organellar genomes in the seaweed *Fucus ceranoides* L. *Molecular Ecology*, 19, 4812–4822.
- RAWSON, P. D. & HILBISH, T. J. (1998). Asymmetric introgression of mitochondrial DNA among European populations of Blue Mussels (*Mytilus spp.*). *Evolution*, 52, 100–108.

- RAWSON, P. D., AGRAWAL, V. & HILBISH, T. J. (1999). Hybridization between the blue mussels *Mytilus galloprovincialis* and *M. trossulus* along the Pacific coast of North America: evidence for limited introgression. *Marine Biology*, 134, 201–211.
- RHYMER, J. M. & SIMBERLOFF, D. (1996). Extinction by hybridization and introgression. *Annual Review of Ecology and Systematics*, 27, 83–109.
- RIESEBERG, L. H. (1997). Hybrid origins of plant species. *Annual Review of Ecology and Systematics*, 28, 359–389.
- RIESEBERG, L. H. (2006). Hybrid speciation in wild sunflowers. *Annals of the Missouri Botanical Garden*, 93, 34–48.
- RIGINOS, C. & CUNNINGHAM, C. W. (2005). Local adaptation and species segregation in two mussel (*Mytilus edulis* × *Mytilus trossulus*) hybrid zone. *Molecular Ecology*, 14, 381–400.
- SAMBROOK, J., FRITSCH, E.F. & MANIATIS, T. (1989). *Molecular cloning: a laboratory manual* (2nd ed.). Cold Spring Harbor NY: Cold Spring Harbor Press.
- SCHWENK, K., BREDE, N. & STREIT, B. (2008). Introduction: Extent, processes and evolutionary impact of interspecific hybridization in animals. *Philosophical Transactions of the Royal Society B Series*, 363, 2805–2811.
- SHEARS, N. T. & BABCOCK, R. C. (2004). *Community composition and structure of shallow subtidal reefs in northeastern New Zealand* (Science for Conservation 245). Wellington, New Zealand: New Zealand Department of Conservation.
- SHEARS, N. T. & BABCOCK, R. C. (2007). *Quantitative description of New Zealand's shallow sub-tidal reef communities* (Science for Conservation 280). Wellington, New Zealand: New Zealand Department of Conservation.
- ZUCCARELLO, G. C. & LOKHORST, G. M. (2005). Molecular phylogeny of the genus *Tribonema* (Xanthophyceae) using *rbcL* gene sequence data: monophyly of morphologically simple algal species. *Phycologia*, 44, 384–392.

Chapter 4

Barcoding brown algae: Does *cox1* identify species of *Cystophora* (Fucales, Phaeophyceae)?

4.1 Abstract

Many brown macroalgal species can be confused because of limited morphological differentiation and plasticity. DNA barcoding has been proposed as an alternative to morphological species identification. In barcoding, short (c. 600 bp) mitochondrial DNA sequences are analysed by distance methods and discontinuities between intraspecific and interspecific mtDNA variation are used as indicators of species boundaries. The utility of *cox1* DNA barcoding was evaluated by comparing 108 sequences from 13 species of *Cystophora*, an Australasian genus of brown algae. A 615 base pair fragment of the *cox1* gene was adequate to assign most specimens to species and resolved some morphologically difficult specimens, but molecular distance between some species was low – often only slightly greater than intraspecific distances. Morphologically ambiguous specimens of *C. subfarcinata*, *C. retorta* and *C. siliquosa* could be identified by *cox1* sequences. Neither *cox1* nor ITS data supported the maintenance of *C. congesta* or *C. distenta* as independent species. I suggest *cox1* alone might be too conserved for reliable barcoding in the brown algae.

4.2. Introduction

DNA BARCODING uses phenetic differences between short mtDNA sequences from a single region for species identification (Hebert *et al.* 2003, 2004; Ratnasingham & Hebert 2007). Proponents of DNA barcoding have suggested that the technique can fill gaps in taxonomic knowledge, document biodiversity loss, facilitate ecological studies and allow objective delimitation of species (Hebert *et al.* 2003). The approach attracted controversy (Will & Rubinoff 2004; Rubinoff *et al.* 2006), in part because both proponents and detractors have failed to distinguish barcoding from DNA taxonomy, species discovery, and molecular phylogenetics. In practice, establishing databases of species tagged sequences has required use of additional markers and morphological and ecological data to assign species names to specimens, effectively engaging in integrative taxonomy (Dayrat 2005), and early suggestions that the approach might supplant traditional taxonomy have abated (Schindel & Miller 2005; Rubinoff 2006). Large

DNA barcoding programmes have been established, particularly in animal groups (Frézal & Leblois 2008; Janzen *et al.* 2009; Golding *et al.* 2009; Ward *et al.* 2009).

Most barcoding studies have delimited species using a discontinuity between intraspecific and interspecific genetic distances (either uncorrected p-distance or Kimura 2-parameter). Hebert *et al.* (2004) proposed a species level threshold of 10-times mean intraspecific variation. Using mean distances inflates discontinuities and the smallest distance should be used (Meier *et al.* 2008). Ten-times thresholds cannot be rigidly applied, as genetic distance between congeners is variable, depending on the time since speciation, generation time, the strength of different speciation processes, population size and connectivity, and mutation and repair rate variation between lineages (Ferguson 2002; Will & Rubinoff 2004; Kumar 2005; Hickerson *et al.* 2006). One solution is to test species divergence using coalescence models (Papadopoulou *et al.* 2008; Baker *et al.* 2009). Character-based species delimitation has been proposed as an alternative to distance methods (DeSalle *et al.* 2005), but its application, so far, is limited (Rach *et al.* 2008).

Spatial heterogeneity of intraspecific variation (Avice 2000; Hoarau *et al.* 2007; Maggs *et al.* 2008) is problematic for barcoding as even geographically widespread sampling might not encompass the most variable populations (Zuccarello *et al.* 2006), and the limited sampling of conspecifics in many barcoding studies might seriously underestimate intraspecific variation and overestimate the number of fixed differences between species. This will produce misleading estimates of genetic distance between species. Alternatively, where genetic structure is high, extensive sampling might result in demes being identified as species (Trewick 2008).

As a technical method of assigning specimens to species, either character-based or distance-based barcoding procedures might assist species identification if a preliminary study shows sufficient interspecific distance or fixed characters to separate closely related (i.e. sister) species, based on adequate sampling over the geographical range of the species; and if interspecific distances or fixed characters are corroborated in a preliminary study with morphological or other characters (i.e. integrated taxonomy, Dayrat 2005).

Most barcoding studies aim to evaluate the efficacy of a marker to delineate species within a group, and to establish a reference database of species tagged

sequences to identify samples from subsequent studies. Specimens are assigned to species according to existing taxonomic criteria; the congruence of disjunctions in genetic variation with *a priori* species assignments is assessed; and, if necessary, taxonomic revisions are made on the basis of corroborated information. I suggest four criteria for a conservative evaluation of a locus for species identification: (1) does the marker resolve discrete clusters within a genus? (2) Are these clusters concordant with species boundaries indicated by other taxonomic information (e.g., morphology, anatomy, ecology, chemistry, physiology, mating studies)? (3) Is sampling sufficient to represent intraspecific variation across the range of each species? (4) Does sampling include closely related species?

The 5' end of the cytochrome oxidase 1 mtDNA gene (*cox1*) is the preferred locus for barcoding in vertebrates and arthropods (Hebert *et al.* 2004; Linares *et al.* 2008; Ward *et al.* 2008; Golding *et al.* 2009). This region is too conserved at the species level in plants and green algae, and various combinations of regions have been proposed as potential barcodes for terrestrial plants (Chase *et al.* 2007; Kress & Erickson 2007; Lahaye *et al.* 2008; Ford *et al.* 2009; Seberg & Petersen 2009) or green algae (Presting 2006; Engelmann *et al.* 2009). Barcoding has been attempted in red algae (Saunders 2005, 2008, 2009; Robba *et al.* 2006; Yang *et al.* 2008), dinoflagellates (Litaker *et al.* 2007), diatoms (Evans *et al.* 2007; Moniz & Kaczmarska 2009, 2010), and brown algae (Lane *et al.* 2007; Kucera & Saunders 2008; McDevit & Saunders 2009). The *cox1* region has been proposed as a suitable locus for barcoding red and brown algae (Saunders 2005; Robba *et al.* 2006).

To evaluate *cox1* as a locus for species identification in brown algae, I compare a barcoding approach to phylogenetic and morphological species delimitation in the genus *Cystophora* J. Agardh (Sargassaceae, Fucales), with an emphasis on New Zealand representatives. *Cystophora* is an ideal genus for testing the barcoding approach as a thorough morphological taxonomy is available (Womersley 1964, 1987), the genus is relatively speciose (23 species listed in Womersley 1964), and includes species with highly disjunct morphologies and others that are morphologically similar.

Barcoding of brown algae by *cox1* has had limited success. Lane *et al.* (2007) attempted a *cox1* barcoding study of *Alaria* species from the north-east Pacific. Intergeneric variation was relatively low and there was no clear threshold between inter-

clade and intra-clade variation. In addition, clusters of *cox1* mitotypes were not congruent with *a priori* morphological species assignments, nor with clades inferred from nrDNA internal transcribed spacer (ITS) sequences. ITS variation was relatively high, but did not resolve species level groups. Lane *et al.* (2007) suggest that widespread introgressive hybridisation precludes molecular delimitation of *Alaria* species in the study region. Kucera & Saunders (2008) used *cox1* sequences to delineate samples of *Fucus* into three clusters, but it is not clear whether these results support wider use of *cox1* in brown algae, as sampling did not extend across the species' ranges, some morphological assignments were not congruent with sequence data and one cluster included two species. McDevit & Saunders (2009) reported *cox1* barcodes for 29 species of brown algae, but did not assess intraspecific or intrageneric diversity. Most genera were represented by a single species and sampling of congeners or conspecifics was geographically restricted. This study shows adequate *cox1* variation to separate genera, but does not address the separation of sister species.

These studies leave the efficacy of *cox1* barcoding of brown algae uncertain. This study aims to further evaluate *cox1* species identification in brown algae by comparing *cox1* species delimitation of *Cystophora* species against morphological criteria and combined *cox1* and ITS phylogenetic analysis.

4.3 Materials and Methods

Cystophora species and close relatives were collected from 51 sites in New Zealand and southern Australia (Table 4.1, Fig. 4.1). Thalli were pressed as voucher specimens. Clean apices were detached and rapidly dried and stored in silica gel for subsequent DNA extraction. Morphological identifications used characters specified by Womersley (1964, 1987). Key diagnostic characters are listed in Table 4.2.

DNA was extracted from approximately 2–5 mg of dry tissue using a modified CTAB buffer procedure (Zuccarello & Lokhorst 2005), with the addition of 1% polyvinylpyrrolidone to the extraction buffer.

PCR and primers

A 615 bp fragment from the 5' end of the *cox1* mitochondrial gene was amplified using primer pairs GAZF2 and GAZR2 (Lane *et al.* 2007) or GAZF2 and a new primer,

fcox808R (5'-TAAACTTCGGGGTGTCCAAA-3'). The latter primer was designed from alignments of the *Fucus*, *Laminaria* and *Desmarestia* mitochondrial genomes (Oudot-Le Seq *et al.* 2002, 2006). Amplification of the ITS1-5.8S-ITS2 cistron used either AFP2(F) (Peters & Burkhardt 1998) or P1 (Tai *et al.* 2001) forward primers and KG4 reverse primer (Lane *et al.* 2006).

Cox1 amplification used a PCR routine with an initial denaturation step of 94°C for 4 min, then 35 cycles of 1 min at 94°C, 30 seconds at 50°C and 1 min at 72°C, with a final extension of 72°C for 7 min. ITS amplifications used a touchdown PCR routine with an initial denaturation step of 94°C for 4 min, then 5 cycles of 1 min at 94°C, seconds at 55°C (−1°C/cycle) and 1 min at 72°C, followed by 30 cycles of 1 min at 94°C,

Fig. 4.1. Map showing sampling locations (for details see Table 4.1). **A.** Southern Australia. **B.** New Zealand.

45 seconds at 50°C and 1 min at 72°C, with a final extension of 72°C for 10 min. The PCR mix for both regions contained 1µl genomic DNA, 0.5 U Taq DNA polymerase (New England Biolabs) 1× ThermoPol reaction buffer (NEB), 7.5 pmoles each primer,

200 nM dNTP and 0.01% BSA. Amplified products were checked for length and yield on 1% agarose gels stained with ethidium bromide.

PCR products were cleaned with ExoSAP-IT (USB, Cleveland, Ohio) enzymes and sequenced commercially (Macrogen Inc., Seoul, Korea). Sequencing *cox1* fragments used the same primers as amplification. ITS sequencing used the additional internal primers KP5 and KIR1 (Lane *et al.* 2006).

Data analysis

Cox1 fragments were aligned using the ClustalW algorithm in MEGA3 (Tamura *et al.* 2007). Alignment was straightforward with no gaps (Appendix 4). A neighbour-joining tree was inferred in PAUP*4.10 (Swofford 2002), with the parameters estimated from MODELTEST 3.7 (Posada & Crandall 1998) (HKY85 + Γ +I, transition/transversion ratio = 3.6467, kappa = 7.489067, nucleotide frequencies: A = 0.1948, C = 0.1909, G = 0.2191, T = 0.3952, invariable sites = 0.6392, Γ = 0.9596). Confidence limits of branches were tested with 1000 bootstrap replications.

ITS sequences were aligned using the ClustalW algorithm in MEGA3, and the alignment edited manually (Appendix 5). A partition homogeneity test for incongruity between *cox1* and ITS datasets was executed in PAUP* (1000 replicates). Maximum-parsimony trees were inferred from the combined ITS and *cox1* sequences (2074 bp) in PAUP* with the parameters estimated from MODELTEST (HKY+I+ Γ , transition/transversion ratio = 2.5289, nucleotide frequencies: A = 0.1801, C = 0.2663, G = 0.3050, T = 0.2486, invariable sites = 0.5809, Γ = 0.7626). 1000 maximum likelihood trees were inferred using GARLI 1.0 (Zwickl 2006) using the same parameters, and a consensus tree was calculated in PAUP*. A Bayesian tree for the combined data was inferred using MrBayes 3.1.2 (Huelsenbeck & Ronquist 2001) with ITS and *cox1* regions partitioned, and each *cox1* codon position partitioned.

Table 4.1. Voucher numbers of specimens used in analyses, collection locations, latitude and longitude and GenBank accession numbers of sequences.

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
				<i>cox1</i>	ITS
WELT A031507	<i>C. congesta</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	GU289263	GU289363
WELT A031510	<i>C. congesta</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	GU289264	GU289364
WELT A031509	<i>C. congesta</i>	Port Campbell VIC	38°37'14.84"S, 142°59'32.69"E	GU289281	-
WELT A031512	<i>C. congesta</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	GU289330	-
WELT A031511	<i>C. congesta</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	GU289285	GU289374
WELT A031508	<i>C. congesta</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	GU289286	GU289375
WELT A030927	<i>C. congesta</i>	The Sirens, Wellington NZ	41°20'56.91"S, 174°45'51.05"E	GU289316	GU289388
WELT A031340	<i>C. distenta</i>	Manukau Reef, Chatham Islands	44° 2'4.90"S, 176°19'44.31"W	GU289317	GU289389
WELT A030933	<i>C. distenta</i>	Mapoutahi, Otago Peninsula NZ	45°44'5.57"S, 170°37'4.53"E	GU289338	-
WELT A031343	<i>C. distenta</i>	Owenga, Chatham Islands	44° 2'0.30"S, 176°19'58.80"W	GU289318	GU289390
WELT A031497	<i>C. monilifera</i>	Eden NSW	37° 4'16.80"S, 149°54'21.85"E	GU289255	GU289356
WELT A031500	<i>C. monilifera</i>	Huskisson, NSW	35° 2'34.31"S, 150°40'38.77"E	GU289248	GU289352
WELT A031501	<i>C. monilifera</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	GU289266	-
WELT A031498	<i>C. monilifera</i>	Ulladulla, NSW	35°21'37.65"S, 150°29'5.55"E	GU289251	-

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
				<i>cox1</i>	ITS
WELT A031499	<i>C. monilifera</i>	Ulladulla, NSW	35°21'37.65"S 150°29'5.55"E,	GU289306	GU289387
WELT A031496	<i>C. moniliformis</i>	Blackmans Beach, Tasmania	43° 0'27.26"S, 147°19'44.06"E	GU289291	GU289379
WELT A031495	<i>C. moniliformis</i>	Jervis Bay, NSW	35° 8'11.52"S 150°44'28.69"E	GU289249	GU289353
WELT A031493	<i>C. platylobium</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289257	GU289358
WELT A031494	<i>C. platylobium</i>	Duttons Beach, Portland VIC	38°19'42.74"S, 141°36'22.27"E	GU289279	-
WELT A030929	<i>C. platylobium</i>	Porpoise Bay, Otago NZ	46°39'39.37"S, 169° 6'29.32"E	GU289237	GU289348
WELT A031492	<i>C. polycystidea</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289261	GU289361
WELT A031489	<i>C. polycystidea</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289262	GU289362
WELT A031491	<i>C. polycystidea</i>	Coles Bay, Tasmania	42° 7'35.52"S, 148°17'32.09"E	GU289298	GU289383
WELT A031490	<i>C. polycystidea</i>	Port Fairy VIC	38°23'42.65"S, 142°14'35.99"E	GU289284	GU289373
WELT ASZ243A	<i>C. retorta</i>	Cape Jaffa SA	36°56'30.94"S, 139°40'30.93"E	GU289309	-
WELT ASZ243B	<i>C. retorta</i>	Cape Jaffa SA	36°56'30.94"S, 139°40'30.93"E	GU289310	-
WELT A031546	<i>C. retorta</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289258	-
WELT A031545	<i>C. retorta</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289259	GU289359

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
WELT A031547	<i>C. retorta</i>	Coles Bay, Tasmania	42° 7'35.52"S, 148°17'32.09"E	GU289297	-
WELT A031550	<i>C. retorta</i>	Point Roadknight VIC	38°25'40.72"S, 144°11'7.78"E	GU289273	-
WELT A031548	<i>C. retorta</i>	Point Roadknight VIC	38°25'40.72"S, 144°11'7.78"E	GU289275	GU289369
WELT A031549	<i>C. retorta</i>	Squeaky Beach, Wilsons Promontory VIC	39° 1'45.89"S, 146°18'24.24"E	GU289278	-
WELT A030919	<i>C. retroflexa</i>	Akaroa, Banks Peninsula, NZ	43°49'49.58"S, 172°54'30.03"E	GU289246	-
WELT A031488	<i>C. retroflexa</i>	Blackmans Beach, Tasmania	43° 0'27.26"S, 147°19'44.06"E	GU289288	GU289377
WELT A031534	<i>C. retroflexa</i>	Coles Bay, Tasmania	42° 7'35.52"S, 148°17'32.09"E	GU289296	-
WELT A031537	<i>C. retroflexa</i>	Coles Bay, Tasmania	42° 7'35.52"S, 148°17'32.09"E	GU289324	-
WELT A031531	<i>C. retroflexa</i>	Depot Beach NSW	35°37'37.91"S, 150°19'37.27"E	GU289252	-
WELT A030917	<i>C. retroflexa</i>	Island Bay, Wellington, NZ	41°20'51.22"S, 174°46'9.77"E	GU289329	-
WELT A030934	<i>C. retroflexa</i>	Kaiti Beach, Gisborne NZ	38°40'55.97"S, 178° 1'45.79"E	GU289333	-
WELT A030918	<i>C. retroflexa</i>	Mahia, East Coast Road, Eastland NZ	39° 5'14.33"S, 177°55'46.40"E	GU289331	-
WELT A031533	<i>C. retroflexa</i>	Point Lonsdale VIC	38°17'20.52"S, 144°36'4.42"E	GU289269	-

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
WELT A030928	<i>C. retroflexa</i>	Porpoise Bay, Otago NZ	46°39'39.37"S, 169° 6'29.32"E	GU289340	-
WELT A031536	<i>C. retroflexa</i>	Port Fairy VIC	38°23'42.65"S, 142°14'35.99"E	GU289342	-
WELT A030914	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	GU289325	-
WELT A030913	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	GU289326	-
WELT A030926	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	GU289232	GU289345
A085	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	GU289233	GU289346
WELT A031535	<i>C. retroflexa</i>	Sorrento, VIC	38°20'5.35"S, 144°44'33.96"E	GU289268	-
WELT A031530	<i>C. retroflexa</i>	Sorrento, VIC	38°20'5.35"S, 144°44'33.96"E	GU289267	GU289366
WELT A030900	<i>C. retroflexa</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	GU289238	GU289349
WELT A030932	<i>C. retroflexa</i>	Tokomaru Bay, Eastland, NZ	38° 6'35.83"S, 178°21'18.27"E	GU289328	-
WELT A030930	<i>C. sp. cf. scalaris</i>	Ahuriri Street, Napier NZ	39°28'42.28"S, 176°54'12.89"E	GU289312	-
WELT A030919	<i>C. scalaris</i>	Akaroa, Banks Peninsula NZ	43°49'49.58"S, 172°54'30.03"E	GU289245	-
WELT A031346	<i>C. scalaris</i>	Kaingaroa, Chatham Islands	43°43'51.20"S, 176°16'2.26"W	GU289303	GU289385
WELT ASG207	<i>C. scalaris</i>	Manukau Reef, Chatham Islands	44° 2'4.90"S, 176°19'44.31"W	GU289241	GU289350

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
WELT A030922	<i>C. scalaris</i>	Ocean Beach, Jackson Bay NZ	43°57'54.44"S, 168°36'21.70"E	GU289243	GU289351
WELT A031345	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	GU289302	-
WELT A030909	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	GU289335	-
WELT A030908	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	GU289336	-
WELT A030910	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	GU289337	-
WELT A031345	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	GU289319	GU289391
WELT A030925	<i>C. scalaris</i>	Pukerua Bay NZ	41° 1'28.36"S, 174°54'5.54"E	GU289235	-
A596*	<i>C. scalaris</i>	Ripapa Island, Banks Peninsula NZ	43°37'7.12"S, 172°45'13.27"E	GU289244	-
WELT A030921	<i>C. scalaris</i>	Riversdale, Wairarapa NZ	41° 6'24.49"S, 176° 4'10.73"E	GU289339	-
B176*	<i>C. scalaris</i>	Auckland Islands, NZ	50°41'5.44"S, 166°17'55.93"E	GU289301	-
WELT A031342	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	GU289313	-
WELT A031341	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	GU289314	-
WELT A031344	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	GU289315	-

* Vouchers for these sites were lost as a result of a chemical spillage accident.

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
A318*	<i>C. scalaris</i>	Tumbledown Bay, Banks Peninsula NZ	43°51'27.57"S, 172°46'13.01"E	GU289242	-
A154*	<i>C. scalaris</i>	Waipapa Point, Kaikoura NZ	42°12'40.90"S, 173°52'28.64"E	GU289236	GU289347
WELT A030911	<i>C. scalaris</i>	Waitangi, Chatham Islands	43°56'41.58"S, 176°33'41.78"W	GU289334	-
WELT A031502	<i>C. siliquosa</i>	Duttons Beach, Portland VIC	38°19'42.74"S, 141°36'22.27"E	GU289307	-
WELT A031506	<i>C. siliquosa</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	GU289265	GU289365
WELT A031505	<i>C. siliquosa</i>	Point Roadknight VIC	38°25'40.72"S, 144°11'7.78"E	GU289274	-
WELT A031503	<i>C. siliquosa</i>	Port Campbell VIC	38°37'14.84"S, 142°59'32.69"E	GU289280	GU289371
WELT A031504	<i>C. siliquosa</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	GU289322	GU289392
WELT A028492	<i>C. sp. cf. C. scalaris</i>	Wilson Bay, Coromandel NZ	36°53'22.06"S, 175°25'32.12"E	GU289320	-
WELT A031526	<i>C. subfarcinata</i>	Bicheno, Tasmania	41°52'29.89"S, 148°18'40.76"E	GU289293	GU289381
WELT A031518	<i>C. subfarcinata</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289260	GU289360
WELT A031521	<i>C. subfarcinata</i>	Dora Bay, Tasmania	41°15'16.35"S, 148°19'4.67"E	GU289295	GU289382

Table 4.1. (continued)

Herbarium voucher	Species	Location	Latitude, Longitude	GenBank Accession numbers	
------------------------------	----------------	-----------------	--------------------------------	--------------------------------------	--

number

WELT	<i>C.</i>	Duttons Beach,	38°19'42.74"S,	GU289276	-
A031522	<i>subfarcinata</i>	Portland VIC	141°36'22.27"E		
WELT	<i>C.</i>	Duttons Beach,	38°19'42.74"S,	GU289341	-
A031520	<i>subfarcinata</i>	Portland VIC	141°36'22.27"E		
WELT	<i>C.</i>	Lonsdale VIC	38°17'5.30"S,	GU289323	-
A031528	<i>subfarcinata</i>		144°36'54.74"E		
WELT	<i>C. retroflexa</i>	Mimosa Rocks,	36°35'2.07"S,	GU289254	GU289355
A031532		Arragunnu NSW	150° 3'15.18"E		
WELT	<i>C.</i>	Observation Head,	35°43'45.50"S,	GU289253	GU289354
A031523	<i>subfarcinata</i>	Batemans Bay NSW	150°12'30.26"E		
WELT	<i>C.</i>	Point Roadknight	38°25'40.72"S,	GU289272	-
A031519	<i>subfarcinata</i>	VIC	144°11'7.78"E		
WELT	<i>C.</i>	Point Roadknight	38°25'40.72"S,	GU289271	GU289368
A031523	<i>subfarcinata</i>	VIC	144°11'7.78"E		
WELT	<i>C.</i>	Port Campbell	38°37'14.84"S,	GU289343	-
A031525	<i>subfarcinata</i>	VIC	142°59'32.69"E		
WELT	<i>C.</i>	Port Campbell	38°37'14.84"S,	GU289287	GU289376
A031514	<i>subfarcinata</i>	VIC	142°59'32.69"E		
WELT	<i>C.</i>	Port Fairy VIC	38°23'42.65"S,	GU289344	-
A031517	<i>subfarcinata</i>		142°14'35.99"E		
WELT	<i>C.</i>	Port Fairy VIC	38°23'42.65"S,	GU289282	GU289372
A031516	<i>subfarcinata</i>		142°14'35.99"E		
WELT	<i>C.</i>	Port Fairy VIC	38°23'42.65"S,	GU289289	GU289378
A031524	<i>subfarcinata</i>		142°14'35.99"E		
WELT	<i>C.</i>	Squeaky Beach,	39° 1'45.89"S,	GU289321	-
A031527	<i>subfarcinata</i>	Wilsons Promontory VIC	146°18'24.24"E		
WELT	<i>C.</i>	Squeaky Beach,	39° 1'45.89"S,	GU289277	GU289370
A031515	<i>subfarcinata</i>	Wilsons Promontory VIC	146°18'24.24"E		

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
WELT A031529	<i>C.</i> <i>subfarcinata</i>	Ulladulla, NSW	35°21'37.65"S, 150°29'5.55"E	GU289250	-
WELT A031544	<i>C. torulosa</i>	Cape Paterson VIC	38°40'29.26"S, 145°37'2.65"E	GU289308	-
WELT A031543	<i>C. torulosa</i>	Coles Bay, Tasmania	42° 7'35.52"S, 148°17'32.09"E	GU289294	-
WELT A031540	<i>C. torulosa</i>	Dora Bay, Tasmania	41°15'16.35"S, 148°19'4.67"E	GU289305	-
WELT A031538	<i>C. torulosa</i>	Eaglehawk Neck, Tasmania	43° 0'32.91"S, 147°55'59.87"E	GU289292	GU289380
WELT A030912	<i>C. torulosa</i>	Maraehako Bay, Bay of Plenty NZ	37°40'19.74"S, 177°47'47.80"E	GU289300	-
WELT A030915	<i>C. torulosa</i>	Nugget Point, Otago NZ	46°26'26.91"S, 169°48'1.78"E	GU289240	-
WELT A030920	<i>C. torulosa</i>	Pihama, Taranaki, NZ	39°31'16.48"S, 173°54'54.98"E	GU289304	GU289386
WELT A031539	<i>C. torulosa</i>	Point Roadknight VIC	38°25'40.72"S, 144°11'7.78"E	GU289270	GU289367
WELT A031541	<i>C. torulosa</i>	Port Fairy VIC	38°23'42.65"S, 142°14'35.99"E	GU289283	-
WELT A030916	<i>C. torulosa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	GU289327	-
WELT A030924	<i>C. torulosa</i>	Pukerua Bay NZ	41° 1'28.36"S, 174°54'5.54"E	GU289234	-
WELT A031542	<i>C. torulosa</i>	Squeaky Beach, Wilsons Promontory VIC	39° 1'45.89"S, 146°18'24.24"E	GU289256	GU289357

Table 4.1. (continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	GenBank Accession numbers	
WELT A031347	<i>C. torulosa</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	GU289239	-
WELT A030931	<i>Landsburgia quercifolium</i>	Pihama, Taranaki, NZ	39°31'16.48"S, 173°54'54.98"E	GU289299	GU289384

Table 4.2. Key morphological characters used in species assignments.

Species	Primary axis	Primary branching	Branching of laterals	Receptacles	Vesicles
<i>C. congesta</i>	Fairly straight, compressed, 4–6(–10) mm broad × 2–4 mm thick	Alternately distichously branched from face of the main axes, Secondary axes usually 20– 50 mm long forming crowded tufts	Irregularly branched on all sides	Slightly compressed, 10–30 mm long × 1–1.5 (–2) mm wide, usually with sterile apical awn	Subspherical, 3– 7(–10) mm diameter, occasionally absent
<i>C. distenta</i>	Strongly zig- zag, flattened, 2–5 mm broad × 1.5–2 mm thick	Alternately distichously branched from face of the main axes	Alternately distichously branched with rounded axils	Compressed, torulose with rows of prominent conceptacles, 10– 40(–50) mm long × 1.5–3 mm broad	Spherical to subspherical, 4– 10 mm diameter
<i>C. expansa</i>	Fairly straight, flattened, 2– 5(–10) mm broad × 1–2(– 4) mm thick	Alternately distichously branched from the face of the main axes	Tristichously branched from a three- sided rhachis	Terete to slightly compressed, slender, 0.5–1(– 1.5) mm long × 0.5–1(–1.5) mm diameter	Small, elongate- ellipsoid, 3–5(–7) long × 1–2 mm broad
<i>C. monilifera</i>	Flattened, 3– 8(–15) mm broad × 1–2 mm thick, with narrow edges	Alternately distichously branched from the face of the main axes	Tristichously branched from a subterete rhachis	Terete, slender, (5–)10–20(–30) long × 0.5–1 mm diameter, constricted between conceptacles	Subspherical to ovoid, 3–6(–8) mm diameter
<i>C. moniliformis</i>	Straight, strongly compressed, 5–20 mm broad × 2–3(– 7) mm thick	Alternately distichously branched from the edge of the main axes	Alternately distichously branched, fasciculate	Moniliform to torulose, 5–15(– 40) long × 0.5–1 mm diameter	Absent

Table 4.2 (Continued)

Species	Primary axis	Primary branching	Branching of laterals	Receptacles	Vesicles
<i>C. platylobium</i>	Strongly compressed, flexuous, (3–)5–10(–15) mm broad × 2–4(–5) mm thick	Alternately distichously branched from the edge of the main axes	Alternately distichously branched, complanate, once pinnate	Strongly flattened, lanceolate to elliptic-lanceolate, basally stalked, 10–30 mm long × 3–8 (–10) mm broad	Spherical to ovoid, 5–15 mm diameter
<i>C. polycystidea</i>	Compressed, quadrangular, 2–7 mm broad × 2–5 mm thick	Alternately distichously branched from the face of the main axes	Radially and spirally branched on all sides	Distantly moniliform, 5–20 mm long and 0.4–0.7(–1) mm diameter at conceptacles	Numerous, small, elongate, tapering at both ends, in clusters, (2–)3–5(–7) mm long × (1–)2–3(–4) mm diameter
<i>C. retorta</i>	Relatively straight, strongly compressed, 2–5(–7) mm broad × 1–2 mm thick	Alternately distichously branched from the face of the main axes	Alternately distichously branched, becoming subdichotomous with broadly rounded axils	Terete to slightly compressed, (10–)20–50(–80) long × 1–2 mm diameter, conceptacles bisexual, often separated by considerable sterile tissue	Usually absent
<i>C. retroflexa</i>	Straight to zig-zag, compressed, 3–10 mm broad × 2–5 mm thick, narrow edged	Alternately distichously branched from the face of the main axes	Terete, irregularly branched on all sides	Compressed, smooth to slightly torulose, 20–50(–60) mm long × 1–2(–2.5) mm broad, apex attenuate with sterile awn, ostiole in axils of conceptacle swelling.	Ovoid to subspherical, 4–10 mm long × 3–6 mm broad
<i>C. scalaris</i>	Zig-zag, compressed, 2.5–15(–20) mm broad × 2–3 mm thick	Alternately distichously branched from face of main axes	Alternately distichously branched, complanate	Ovoid to compressed, smooth to torulose, 1.5–2.5 mm wide × 1 mm thick	Spherical to ovoid 2–7(–10) mm diameter
<i>C. siliquosa</i>	Straight, quadrangular to square in section, 2–5 mm broad and thick	Alternately distichously branched from the face of the main axes	Alternately distichously branched, becoming subdichotomous with rounded axils	Slightly to markedly compressed, 30–80 mm long × 1–2 mm broad, conceptacles unisexual and thalli dioecious	Absent

Table 4.2 (Continued)

Species	Primary axis	Primary branching	Branching of laterals	Receptacles	Vesicles
<i>C. cuspidata</i>	Fairly straight, compressed, 3–6 mm broad × 1–2 thick	Alternately distichously branched from the face of the main axes	Usually irregularly alternately (sometime sub-distichously) and closely branched.	Simple or branched, (10–)20–60(–80) mm long with prominent swollen conceptacles closely arranged in 2–3 rows in the basal half or more of receptacles, and here 1.5–4 mm broad	Absent
<i>C. subfarcinata</i>	Slightly flexuous, compressed, 2–7 mm broad × 1–2(–3) thick	Densely and distichously branched from the face of the main axes	Irregularly alternately branched on all sides	Often branched, (5–)10–30(–50) mm long × 1–2 mm thick at conceptacles, with prominent scattered conceptacles separated by sterile tissue with a sterile awn, ostiole apical on conceptacle swelling.	Present or absent, elongate ovoid to sub spherical, 2–04(–6) mm long × 2–3(–4) mm diameter
<i>C. torulosa</i>	Compressed, 3–10 mm broad × 2–5 mm thick, straight to slightly flexuous	Alternately densely branched from the face of the main axes	Densely, irregularly alternately subdistichously branched	Terete to slightly three sided, swollen with a blunt rounded apex, 10–50(–70) mm long × (1–)2–3(–4) mm diameter	Subspherical to ovoid, 3–8(–10) long × 3–6(–9) mm diameter

4.4 Results

Amplification and species sampling

The *cox1* region was amplified from 108 specimens of 13 species of *Cystophora* and five specimens of putative outgroup genera (*Caulocystis*, *Carpoglossum* and *Landsburgia*) (Table 4.1). I was unable to amplify the *cox1* region of *Cystophora expansa* (Areschoug) Womersley with any primer combination and suspect a primer mismatch. The GAZF2–fcox808R primer combination successfully amplified most *Cystophora* species, whereas the GAZF2–GAZR2 combination was successful when amplifying *Caulocystis*, *Carpoglossum*, *Landsburgia* and other outgroup species.

The ITS1–5.8S–ITS2 cistron was amplified from 47 specimens of *Cystophora* and from *Landsburgia quercifolia* (Hooker f. et Harvey) Harvey. Preliminary trees positioned *Landsburgia* as the genus most closely related to *Cystophora* and *L. quercifolia* was designated as an outgroup in subsequent analyses.

Molecular distance

A distance tree inferred from *cox1* sequences showed clusters that were generally consistent with morphological species assignments. Sequences from four specimens of *Cystophora* were incongruent with morphological assignments (indicated by arrows in Fig. 4.2). DNA from these specimens was subsequently re-extracted and sequenced, with the same results. In two of these specimens (WELT A031539 and WELT A031523) morphological identification was unambiguous and this identification is retained. Two specimens (WELT A028492 and WELT A030930) were initially assigned to *C. scalaris* J. Agardh but grouped with *C. torulosa* (R. Brown) J. Agardh and *C. retroflexa* (Labillardière) J. Agardh respectively. These specimens were re-examined and as identity is ambiguous I designated them as *Cystophora* sp. cf. *scalaris*.

Uncorrected interspecific pairwise differences between *cox1* sequences in *Cystophora* varied from 0 to 49 (0–8%). Intraspecific distances varied from 0 to 4 (0–0.65%). Kimura 2-parameter distances ranged from 0 to 8.6% between species and 0 to 0.7% within species (Table 4.2, Fig. 4.2). In general, species with wide geographic ranges (e.g., *C. retorta* (Mertens) J. Agardh, *C. retroflexa*) showed greater intraspecific variation than geographically restricted species (e.g., *C. monilifera* J. Agardh).

Fig. 4.3. ML topology of *Cystophora* species inferred using combined ITS1–5.8S–ITS2 and *cox1* sequences, support is shown at nodes as maximum parsimony/maximum likelihood/Bayesian posterior probabilities. Specimen labels show GenBank accession numbers for ITS sequences only.

Fig. 4.2. (Previous page). Distance tree inferred in PAUP* by neighbour-joining from 108 *cox1* sequences of *Cystophora* and one outgroup. Confidence is shown at nodes based on 1000 bootstrap replicates. Arrows indicate specimens where morphology and sequence position were incongruent.

Table 4.3. Number of changes in *cox1* sequences between species (above diagonal) and K2P distances (below diagonal)

	<i>C. congesta</i>	<i>C. monilifera</i>	<i>C. moniliformis</i>	<i>C. platylobium</i>	<i>C. polycystidea</i>	<i>C. retorta</i>	<i>C. retroflexa</i>	<i>C. scalaris</i>	<i>C. siliquosa</i>	<i>C. subfarcinata</i>	<i>C. torulosa</i>
<i>C. congesta</i>	0–2	9–11	47–48	16–18	11–13	16–20	0–3	8–11	18–21	3–6	10–13
<i>C. monilifera</i>	0.015–0.018	0 (0.00)	49	18	14	19–22	10–11	13–14	20–21	8–9	13–14
<i>C. moniliformis</i>	0.082–0.084	0.086	2 (0.003)	41–43	51	40–43	47–48	46	37–40	47–49	48–49
<i>C. platylobium</i>	0.027–0.030	0.030	0.071–0.075	0 (0.00)	19	6–9	17–18	16–17	8–9	15–16	15–16
<i>C. polycystidea</i>	0.018–0.022	0.023	0.090	0.032	0 (0.00)	20–23	12–13	15–16	23–24	10–11	13–14
<i>C. retorta</i>	0.029–0.034	0.032–0.037	0.069–0.075	0.010–0.015	0.035–0.039	0–4 (0.00–0.07)	17–20	16–19	7–10	16–20	16–20
<i>C. retroflexa</i>	0.00–0.005	0.017–0.018	0.082–0.084	0.027–0.030	0.020–0.22	0.028–0.032	0–3 (0.00–0.005)	8–10	19–21	4–6	11–13
<i>C. scalaris/distenta</i>	0.013–0.018	0.022–0.023	0.080	0.028–0.027	0.025–0.027	0.027–0.032	0.013–0.017	0–1 (0.00–0.002)	18–20	9–11	8–10
<i>C. siliquosa</i>	0.030–0.035	0.034–0.035	0.064–0.069	0.013–0.015	0.039–0.041	0.012–0.015	0.032–0.034	0.030–0.034	0–1 (0.00–0.002)	19–21	19–21
<i>C. subfarcinata</i>	0.002–0.010	0.013–0.015	0.082–0.086	0.025–0.027	0.017–0.018	0.027–0.034	0.007–0.010	0.015–0.018	0.032–0.035	0–2 (0.00–0.003)	9–11
<i>C. torulosa</i>	0.017–0.022	0.022–0.023	0.084–0.086	0.025–0.027	0.032–0.035	0.029–0.034	0.018–0.022	0.015–0.017	0.034–0.035	0.015–0.018	0–1 (0.00–0.002)

Distances between closely related species were often low, in the same order as intraspecific distances. In some cases these species were easily distinguished by morphology (e.g., *C. retorta* and *C. platylobium* (Mertens) J. Agardh), in other cases they were not (*C. siliquosa* J. Agardh and *C. retorta*, *C. retroflexa* and *C. subfarcinata* (Mertens) J. Agardh).

Phylogeny

A partition homogeneity test showed significant congruence between ITS and *cox1* data sets ($P = 0.999$). Analyses of combined data produced trees with good support for most branches, but did not separate *C. congesta/retroflexa* and *C. scalaris/distenta*, and did not show support for the position of *C. platylobium* (Fig. 4.3).

Cystophora moniliformis (Esper) Womersley & Nizamuddin is sister to all other *Cystophora* species. The early branching position of *C. moniliformis* is consistent with the rather anomalous form of this species (a straight, dorsiventrally flattened main axis, with fine laterals developing from the sides of the main axes). The remaining species formed two groups, with *C. platylobium* sister to a group containing *C. retorta* and *C. siliquosa*, but with only weak support, and this group is sister to a larger group containing all other species. Species level clades were well supported within this latter group, but the higher level branching was only weakly supported (61–68%).

4.5 Discussion

Few studies have attempted *cox1* DNA barcoding of brown algae. These studies have either not discriminated species (Lane *et al.* 2007), not sampled across species ranges, risking underestimating intraspecific divergence (Kucera & Saunders 2008), or have sampled a limited number of congeners (McDevit & Saunders 2009). While using DNA sequences to identify algae is an established technique (e.g., Sasaki *et al.* 2003; Lane & Saunders 2005; Fox & Swanson 2007), a prerequisite for extending this case-by-case molecular identification into a DNA barcoding system is to firmly establish the utility of one or more markers for delimiting a wide range of species. I discuss the utility of *cox1* barcoding in *Cystophora* according to the criteria outlined in the introduction:

1. Does the marker resolve discrete clusters within a genus? Yes, a 615 bp fragment from the *cox1* was sufficient to resolve clusters, albeit with little bootstrap support, and in some cases, with low molecular distances between clusters.

2. Are clusters concordant with recognised species boundaries? Most specimens collected in this study could be assigned to species using *cox1* sequences alone. In two cases, closely related species could not be separated by *cox1* or combined *cox1*–ITS (*C. distenta*–*C. scalaris*, *C. retroflexa*–*C. congesta*). In several cases, interspecific variation between morphologically distinct species was only slightly greater than intraspecific variation. This is unsatisfactory if distance methods alone are used to define species boundaries.

3. Is sampling sufficient to represent intraspecific variation across the range of each species? Sampling of species that occur in New Zealand was reasonably intensive and widespread, which strengthens confidence in *cox1* as an estimator of species identity, estimates of intraspecific variation in Australian species is likely to be low because of limited sampling of Australian populations. Further sampling of these species might unearth novel haplotypes that are difficult to assign to species.

4. Is sampling of congeners sufficient to show resolution of closely related species?

Some western Australian *Cystophora* species and some deepwater species were not sampled. I was unable to amplify *C. expansa* with any primer combination.

Cystophora expansa is morphologically similar to *C. monilifera*, and I am unable to evaluate *cox1* for discriminating these species. However species sampled included several pairs of sister species, an important consideration for evaluating a barcoding marker.

Distance methods

McDevit & Saunders (2009) suggested that the ten-fold threshold between intraspecific and interspecific variation proposed by Hebert *et al.* (2004) for separating species with *cox1* could be applied to brown algae. In *Cystophora*, *C. moniliformis* was highly divergent from most other species, with pairwise distances between all other species exceeding the ten-fold threshold. Species in which low intraspecific variation was detected were also separated from most other congeners by this threshold. However, species with moderate intraspecific variation (*C. retorta*, *C. retroflexa* and *C. subfarcinata*) often showed distances to closely related congeners of around twice intraspecific distances. A lower threshold must be applied if this method

is used to separate species in *Cystophora*. The appeal of distance thresholds is that species can be efficiently demarcated from large datasets using sequence data alone and automated using algorithms. It is generally recognised that a single threshold value cannot be applied to all groups of organisms (Vogler & Monaghan 2006), but establishing an operational threshold requires extensive sampling to accurately estimate intraspecific marker variation. Mitochondrial DNA variation in brown algae is often highly spatially structured (Hoarau *et al.* 2007; Fraser *et al.* 2009; Uwai *et al.* 2009; Chapter 2) so sampling has to be intensive within populations and, ideally, across the range of the species. Only Lane *et al.* (2007) have applied barcoding methods to an intensively sampled collection of brown algae and they failed to delineate species. Kucera & Saunders (2008) used multiple samples of *Fucus* species, but only sampled in Canada where their most clearly delineated species, *F. serratus*, is a recent introduction (Lyons & Scheibling 2009). While I sampled across the range of several *Cystophora* species, it is likely that intraspecific variation is underestimated.

Species level differences in intraspecific variation in *Cystophora* did not correlate with sampling effort. *Cox1* variation was high in *C. subfarcinata* (n = 16) and *C. retroflexa* (n = 20) which were extensively sampled, but was also high in *C. retorta*, where sampling was limited (n = 8). Conversely low intraspecific variation was found in *C. scalaris* (n = 19) and *C. torulosa* (n = 12), but collections span mainland Australia, Tasmania and New Zealand. Intraspecific *cox1* variation in brown algae appears to vary widely. Kucera & Saunders (2008) found low (0–0.3%) *cox1* intraspecific variation in *Fucus*, albeit based on limited geographical sampling, whereas a worldwide study of *Durvillea antarctica* (Fraser *et al.* 2009) found relatively high *cox1* diversity (uncorrected distances of up to 5.4%). Similar problems have been found when using *cox1* to demarcate well sampled red algal species (Sherwood *et al.* 2008; Yang *et al.* 2008).

Geographically widespread sampling and the inclusion of sister species are expected to reduce variation thresholds between species, especially if the marker used is not evolving and sorting sufficiently quickly to demarcate species formed in recent radiations (Shaffer & Thomson 2007). McDevit & Saunders' (2009) survey of interspecies *cox1* variation of brown algae did not include sister species or extensive intraspecies sampling (apart from *Fucus* sequences from Kucera & Saunders (2008)), and their support for a ten-times threshold in brown algae is likely to be an artefact of

low sampling of conspecifics and congeners rather than an indicator of the efficacy of *cox1* barcoding.

Character based DNA barcoding is an alternative to distance methods (DeSalle *et al.* 2005; Rach *et al.* 2008). This has the advantage of defining species by synapomorphies, rather than phenetic clustering, and is concordant with phylogenetic species concepts, but would also require wider geographical sampling of closely related species to ensure characters nominated are conserved in species.

Taxonomic oversplitting

Four *Cystophora* species were not delimited in any molecular analyses. No evidence was found for a distinct obligate epiphyte species equivalent to *C. cymodoceae*; no specimens could be unequivocally assigned to *C. cuspidata*, and specimens assigned to *C. congesta* and *C. distenta* grouped with other species in molecular analyses.

Failure to separate species by molecular markers can be attributed to three processes: (1) insufficient time for stochastic lineage sorting processes (coalescence) to sort the marker to reciprocal monophyly; (2) ongoing gene flow by hybridisation; or (3) taxonomic over-splitting of species. In these cases taxonomic oversplitting (Dayrat 2005) is a probable cause.

Cystophora retroflexa (Labillardière) J. Agardh and *C. congesta* Womersley & Nizamuddin ex Womersley are morphologically similar and these two species were not separated in any molecular analysis (Figs. 4.2, 4.3). Rather, haplotypes and ribotypes from *C. congesta* and *C. retroflexa* were intermixed. Womersley (1964, 1987) separated these species by vegetative characters, but accepted that intergrades occurred. *Cystophora retroflexa* has longer (20–60 mm) receptacles and a more openly branched habit. *Cystophora congesta* has shorter (15–30 mm) receptacles and tufted laterals and ramuli, and usually ridges on the primary axes. Seven specimens were assigned to *C. congesta*, including three from the type locality (Robe, SA) but some specimens assigned to *C. retroflexa* (e.g., WELT A031533) tended toward a *C. congesta*-like morphology.

Cystophora scalaris and *C. distenta* did not separate in any molecular analysis. *Cystophora distenta* was established by J. Agardh based on material from the Chatham Islands (Agardh 1870). The species has been reported from Wellington, New Zealand southwards and the Chatham Islands (Lindauer *et al.* 1961). It is similar to *C. scalaris* but with longer, broader and more torulose receptacles, but dimensions

given by Womersley (1964) overlap (Table 4.2). Three specimens collected from the Chatham Islands and the South Island of New Zealand were considered sufficiently morphologically distinct to assign to *C. distenta*, other specimens showed intermediate morphologies and were assigned to *C. scalaris*.

I consider that morphological characters used to separate *C. distenta*/*C. scalaris* and *C. retroflexa*/*C. congesta* are continuous and in the absence of molecular divergence oversplitting is probable. These results should be regarded as provisional as other closely related species of brown algae cannot be separated using mitochondrial sequences, due to ongoing gene flow or incomplete lineage sorting, but can be resolved with microsatellite data (Bergström *et al.* 2005; Billard *et al.* 2005; Pereyra *et al.* 2009).

The southern Australian species *Cystophora subfarcinata* and *C. cuspidata* are separated according to the arrangement and development of conceptacles (Womersley 1987). *Cystophora cuspidata* has receptacles that are broader (1.5–4 mm) with the lower conceptacles less scattered and arranged in two or three rows. *Cystophora subfarcinata* has narrower ((5–)10–30(–50) mm long $\times$ 1–2 mm) receptacles with usually scattered conceptacles. These characters intergrade and some specimens (e.g., WELT A031517) could have been assigned to either species. Specimens resembling *C. cuspidata* did not cluster separately from *C. subfarcinata* in the molecular analyses. I agree with Womersley (1987) that *C. cuspidata* is probably only a form of *C. subfarcinata*, but I was unable to collect samples from the type locality of *C. cuspidata* (Encounter Bay, SA), and specimens sampled did not have conceptacles as prominent as those illustrated by Womersley 1987 (Fig. 149 H, p. 403). Therefore I do not reduce *C. cuspidata* to synonymy with *C. subfarcinata* but recommend further work on these species.

Revisions in species assignments

Some specimens were morphologically intermediate between *C. retroflexa* and *C. subfarcinata*, or between *C. retorta* and *C. siliquosa*, and preliminary morphological assignments were incongruent with *cox1* sequence clusters. These specimens were re-examined and species assignments were revised using characters discussed below.

Cystophora retroflexa and *C. subfarcinata* can resemble one another when young, infertile or stunted, but fertile specimens could be distinguished using the shape of the conceptacle. In *C. subfarcinata* the conceptacle is spherical or

subspherical; in *Cystophora retroflexa* the conceptacles are elongate. Some specimens that were initially assigned to *C. retroflexa* clustered with *C. subfarcinata* in molecular analyses. Careful examination of conceptacles in these specimens allowed us to reassign them to *C. subfarcinata*, despite other characters being consistent with *C. retroflexa*.

Cystophora siliquosa and *C. retorta* share complanately branched laterals, rounded axils and relatively straight main axes. Presence of vesicles is a diagnostic character for *C. retorta*, but this is of limited value as vesicles are absent on some specimens (Womersley 1964, 1987). I could not distinguish species by the shape of the main axis, as suggested by Womersley (1964) (see Table 4.2), and found microscopic examination of conceptacles to determine monoecy/dioecy was required to confidently assign specimens to species.

In both these species pairs, low morphological divergence and plasticity made species assignments difficult. In these cases *cox1* sequences provided a useful indication that more careful morphological examination was required.

Other markers

These results, together with similar low variation in previous studies (Lane *et al.* 2007; Kucera & Saunders 2008) suggests while *cox1* has some utility for species identification in brown algae, it is too conserved to serve as a basis for a single marker barcoding system. Mitochondrial markers with sufficient resolution to discriminate closely related brown algae are rare, and low genetic divergence between recognised species of brown algae appears common (Van Oppen *et al.* 1993; Coyer *et al.* 2001, 2006; Stiger *et al.* 2000). A small proportion of errors (either failing to separate sister species, or recognising demes as species) might be acceptable in some applications of barcoding, where the trade-off is reduced time and cost (e.g., Moniz & Kaczmarska 2009), but if barcoding is to contribute usefully to brown algal taxonomy other candidate markers should be evaluated.

Screening the mitochondrial genome of *Saccharina* species (Yotsukura *et al.* 2009) found mitochondrial regions that are more variable than *cox1* and these could be evaluated for species discrimination. In addition, the ITS1–5.8S–ITS2 cistron, or for ease of amplification, ITS2 alone, might be a useful addition to *cox1* for species identification (see Coleman & Mai 1997 in Chlorophyta, Guillemin *et al.* 2008 and Hu *et al.* 2009 in Rhodophyta, Litaker *et al.* 2007 in dinoflagellates, Moniz &

Kaczmarska 2009, 2010 in diatoms). The use of ITS sequences for barcoding has been criticised because length differences between species create alignment problems (McDevit & Saunders 2009). However if barcoding is limited to species assignment (Ratnasingham & Hebert 2007) this is not problematic, as inability to produce an alignment can be regarded as strong evidence for separating species (Müller *et al.* 2009) and alignment methods that better handle indels (Steinke *et al.* 2005) or alignment-free methods of sequence comparison (Chu *et al.* 2006, 2009; Kuksa & Pavlovic 2009) can be used.

In conclusion, molecular data did not support the separate species status of *C. distenta* and *C. congesta*, and further work is required to confirm the separate status of *C. cuspidata* and *C. cymodoceae*. Other species were well supported by both morphological and molecular data. These results suggest that *cox1* barcoding might have utility in brown algae where delimiting closely related or sister species is not an issue (for example many studies identifying invasive algae, or studies where generic identity of cryptic life stages is sufficient). Species identification by *cox1* might also be useful in selected groups of brown algae or in restricted geographical areas, but reliance on this marker will require preliminary sampling across the area of interest with adequate sampling and corroboration by other data.

4.6 Acknowledgements

I thank the New Zealand Department of Conservation and Education NZ Study Abroad Scheme for funding assistance; Wendy Nelson, Tracy Farr, Kate Neill and the Department of Conservation Chatham Islands staff for assistance with collections and Alan Millar (National Herbarium of New South Wales) and Jenn Dalen (Museum of New Zealand Te Papa Tongarewa herbarium) for assistance with herbaria access.

4.7 References

- AGARDH, J. G. (1870). Om Chatham-öarnes Alger. *Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar, Stockholm*, 5, 435–456.
- AVISE, J. C. (2000). *Phylogeography: The History and Formation of Species*. Cambridge MA: Harvard University Press.
- BAKER, A. J., TAVERES, E. S. & ELBOURNE, R. F. (2009). Countering criticisms of single mitochondrial DNA gene barcoding in birds. *Molecular Ecology Resources*, 9, (Suppl. 1) 257–268.
- BERGSTRÖM, L., TATARENKOV, A., JOHANNESSON, K. JÖNSSON, R. B. & KAUTSKY, L. (2005). Genetic and morphological identification of *Fucus radicans* sp. nov. (Fucales, Phaeophyceae) in the brackish Baltic Sea. *Journal of Phycology*, 41, 1025–1038.

- BILLARD, E., DAGUIN, C., PEARSON, G., SERRÃO, E., ENGEL, C. & VALERO, M. (2005). Genetic isolation between three closely related taxa: *Fucus vesiculosus*, *F. spiralis* and *F. ceranoides* (Phaeophyceae). *Journal of Phycology*, *41*, 900–905.
- CHASE, M. W., COWAN, R. S., HOLLINGSWORTH, P. M., VAN DEN BERG, C., MADRIÑAN, S., *ET AL.* (2007). A proposal for a standardised protocol to barcode all land plants. *Taxon*, *56*, 295–299.
- CHU, K. H., LI, C. P. & QI, J. (2006). Ribosomal RNA as molecular barcodes: a simple correlation analysis without sequence alignment. *Bioinformatics*, *22*, 1690–1701.
- CHU, K. H., XU, M. & LI, C. P. (2009). Rapid DNA barcoding analysis of large datasets using the composition vector method. *BMC Bioinformatics*, *10*, (Suppl. 14), S8.
- COLEMAN, A. W. & MAI, J. C. (1997). Ribosomal DNA ITS-1 and ITS-2 sequence comparisons as a tool of predicting genetic relatedness. *Journal of Molecular Evolution*, *45*, 168–177.
- COYER, J. A., SMITH, G. J. & ANDERSEN, R. A. (2001). Evolution of *Macrocystis* spp. (Phaeophyceae) as determined by ITS1 and ITS2 sequences. *Journal of Phycology*, *37*, 574–585.
- COYER, J. A., HOARAU, G., OUDOT-LE SECQ, M-P., STAM, W. T. & OLSEN, J. L. (2006). A mtDNA-based phylogeny of the brown algal genus *Fucus* (Heterokontophyta; Phaeophyta). *Molecular Phylogenetics and Evolution*, *39*, 209–222.
- DAYRAT, B. (2005). Towards integrative taxonomy. *Biological Journal of the Linnean Society*, *85*, 407–415.
- DESALLE, R., EGAN, M. G. & SIDDALL, M. (2005). The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, *360*, 1905–1916.
- ENGELMANN, J. C., RAHMANN, S., WOLF, M., SCHULTZ, J., FRITZILAS, E., *ET AL.* (2009). Modelling cross-hybridization on phylogenetic DNA microarrays increases the detection power of closely related species. *Molecular Ecology Resources*, *9*, 83–93.
- EVANS, K. M., WORTLEY, A. H. & MANN, D. G. (2007). An assessment of potential diatom “barcode” genes (*cox1*, *rbcL*, 18S and ITS rDNA) and their effectiveness in determining relationships in *Sellaphora* (Bacillariophyta). *Protist*, *158*, 349–364.
- FERGUSON, J. W. H. (2002). On the use of genetic divergence for identifying species. *Biological Journal of the Linnean Society*, *75*, 509–516.
- FORD, C. S., AYRES, K. L., TOOMEY, N., HAIDER, N., VAN ALPHEN STAHL, J., KELLY, L. J., *ET AL.* (2009). Selection of candidate coding DNA barcoding regions for use on land plants. *Botanical Journal of the Linnean Society*, *159*, 1–11.
- FOX, C. H. & SWANSON, A. K. (2007). Nested PCR detection of microscopic life-stages of laminarian macroalgae and comparison with adult forms along intertidal height gradients. *Marine Ecology Progress Series*, *332*, 1–10.
- FRASER, C. L., NIKULA, R., SPENCER, H. G. & WATERS, J. (2009). Kelp genes reveal effects of subantarctic sea ice during the Last Glacial Maximum. *Proceedings of the National Academy of Sciences of the U.S.A.*, *106*, 3249–3253.
- FRÉZAL, L. & LEBLOIS, R. (2008). Four years of DNA barcoding: Current advances and prospects. *Infection, Genetics and Evolution*, *8*, 727–736.
- GOLDING, G. B., HANNER, R. & HEBERT, P. D. N. (2009). Preface. *Molecular Ecology Resources*, *9*, (Suppl. 1) iv–vi.
- GUILLEMIN, M.-L., AIT AKKI, S., GIVERNAUD, T., MOURADI, A., VALERO, M., DESTOMBE, C. (2008). Molecular characterisation and development of rapid molecular methods identify species of Gracilariaceae from the Atlantic coast of Morocco. *Aquatic Botany*, *89*, 324–330.
- HEBERT, P. D. N., CYWINSKA, A., BALL, S. L. & DE WAARD, J. R. (2003). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B Series*, *270*, 313–321.
- HEBERT, P. D. N., STOECKLE, M. Y., ZEMLAK, T. S. & FRANCIS, C. M. (2004). Identification of birds through DNA barcodes. *PLoS Biology*, *2*, e312. doi: 10.1371/journal.pbio.0020312.

- HICKERSON, M. J., MEYER, C. P. & MORITZ, C. (2006). DNA barcoding will often fail to discover new animal species over broad parameter space. *Systematic Biology* 55: 729–739.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- HU, Z. M., GUIRY, M. D. & DUAN, D. L. (2009). Using the ribosomal internal transcribed spacer (ITS) as a complement marker for species identification of red macroalgae. *Hydrobiologia*, 635, 279–287.
- HUELSENBECK, J. P. & RONQUIST, F. (2001). MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17, 754–755.
- JANZEN, D. H., HALLWACHS, W., BLANDIN, P., BURNS, J. M., CADIOU, J.-M., CHACON, I., ET AL. (2009). Integration of DNA barcoding into an ongoing inventory of complex tropical biodiversity. *Molecular Ecology Resources*, 9, (Suppl. 1), 1–26.
- KRESS, W. J. & ERICKSON, D. L. (2007). A two-locus global DNA barcode for land plants: The coding *rbcL* gene complements the non-coding *trnH/psbA* spacer region. *PLoS ONE*, 2, e508. doi 10.1371/journal.pone.0000508.
- KUCERA, H. & SAUNDERS, G. W. (2008). Assigning morphological variants of *Fucus* (Fucales, Phaeophyceae) in Canadian waters to recognized species using DNA barcoding. *Botany*, 86, 1065–1079.
- KUKSA, P. & PAVLOVIC, V. (2009). Efficient alignment-free DNA barcode analytics. *BMC Bioinformatics*, 10, (Suppl. 14), S9.
- KUMAR, S. (2005). Molecular clocks – four decades of evolution. *Nature Reviews Genetics*, 6, 654–662.
- LAHAYE, R., VAN DER BANK, M., BOGARIN, D., WARNER, J., PUPULIN, F., GIGOT, G., ET AL. (2008). DNA barcoding the floras of biodiversity hotspots. *Proceedings of the National Academy of Sciences of the USA*, 105, 2923–2928.
- LANE, C. E. & SAUNDERS, G. W. (2005). Molecular investigation reveals epi/endophytic extragenetic kelp (Laminariales, Phaeophyceae) gametophytes colonizing *Lessoniopsis littoralis* thalli. *Botanica Marina*, 48, 426–436.
- LANE, C. E., MAYES, C. M., DRUEHL, L. & SAUNDERS, G. W. (2006). A multi-gene molecular investigation of the kelp (Laminariales, Phaeophyceae) resolves competing phylogenetic hypotheses and supports substantial taxonomic re-organization. *Journal of Phycology*, 42, 493–512.
- LANE, C. E., LINDSTROM, S. C. & SAUNDERS, G. W. (2007). A molecular assessment of northeast Pacific *Alaria* species (Laminariales, Phaeophyceae) with reference to the utility of DNA barcoding. *Molecular Phylogenetics and Evolution*, 44, 634–648.
- LINARES, M. C., SOTO-CALDERÓN, I. D., LEES, D. C. & ANTHONY, N. M. (2008). High mitochondrial diversity in geographically widespread butterflies of Madagascar: A test of the DNA barcoding approach. *Molecular Phylogenetics and Evolution*, 50, 485–495.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae. *Nova Hedwigia*, 3, 129–350.
- LITAKER, R. W., ZANDERSEA, M. W., KIBLER, S. R., REECE, K. S., STOKES, N. A., LUTZONI, F. M., ET AL. (2007). Recognizing dinoflagellate species using ITS rDNA sequences. *Journal of Phycology*, 43, 344–355.
- LYONS, D. A. & SCHEIBLING, R. E. (2009). Range expansion by invasive marine algae: rates and patterns of spread at a regional scale. *Diversity and Distributions*, 15, 1–14.
- MAGGS, C. A., CASTILHO, R., FOLTZ, D., HENZLER, C., JOLLY, M. T., KELLY, J., ET AL. (2008). Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology*, 89, S108–S122.
- MCDEVIT, D. C. & SAUNDERS, G. W. (2009). On the utility of DNA barcoding for species differentiation among brown macroalgae (Phaeophyceae) including a novel extraction protocol. *Phycological Research*, 57, 131–141.

- MEIER, R., ZHANG, G. Y. & ALI, F. (2008). The use of mean instead of smallest interspecific distances exaggerates the size of the “Barcoding Gap” and leads to misidentification. *Systematic Biology*, 57, 809–813.
- MONIZ, B. J. M. & KACZMARSKA, I. (2009). Barcoding diatoms: Is there a good marker? *Molecular Ecology Resources*, 9, (Suppl. 1), 65–74.
- MONIZ, B. J. M. & KACZMARSKA, I. (2010). Barcoding of diatoms: Nuclear encoded ITS revisited. *Protist*, 161, 7–34.
- MÜLLER, T., PHILIPPI, N., DANDEKAR, T., SCHULTZ, J. & WOLF, M. (2009). Distinguishing species. *RNA* 13, 1469–1472.
- LOUDON, M.-P., KLOAREG, B. & LOISEAUX-DE GÖER, S. (2002). The mitochondrial genome of the brown alga *Laminaria digitata*: a comparative analysis. *European Journal of Phycology*, 37, 163–172.
- LOUDON, M.-P., LOISEAUX-DE GÖER, S., STAM, W. T. & OLSEN, J. L. (2006). Complete mitochondrial genomes of the three brown algae (Heterokonta: Phaeophyceae) *Dictyota dichotoma*, *Fucus vesiculosus* and *Desmarestia viridis*. *Current Genetics*, 49, 47–58.
- PAPADOPOULOU, A., BERGSTEN, J., FUJISAWA, T., MONAGHAN, M. T., BARRACLOUGH, T. G. & VOGLER, A. P. (2008). Speciation and DNA barcodes: testing the effects of dispersal on the formation of discrete sequence clusters. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, 363, 2987–2996.
- PEREYRA, R. T., BERGSTRÖM, L., KAUTSKY, L. & JOHANNESSON, K. (2009). Rapid speciation in a newly opened postglacial marine environment, the Baltic Sea. *BMC Evolutionary Biology*, 9, 70. doi: 10.1186/1471-2148-9-70
- PETERS, A. F. & BURKHARDT, E. (1998). Systematic position of the kelp endophyte *Laminarionema elsbetiae* (Ectocarpales *sensu lato*, Phaeophyceae) inferred from nuclear ribosomal DNA sequences. *Phycologia*, 37, 114–120.
- POSADA, D. & CRANDALL, K. A. (1998). Modeltest: Testing the model of DNA substitution. *Bioinformatics*, 14, 817–818.
- PRESTING, G. G. (2006). Identification of conserved regions in the plastid genome: implications for DNA barcoding and biological function. *Canadian Journal of Botany*, 83, 1434–1443.
- RACH, J., DESALLE, R., SARKAR, I. N., SCHIERWATER, B. & HADRY, H. (2008). Character-based DNA barcoding allows discrimination of genera, species and populations in Odonata. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, 275, 237–247.
- RATNASINGHAM, S. & HEBERT, P. D. N. (2007). BOLD: The Barcode of Life Data System (www.barcodinglife.org). *Molecular Ecology Notes*, 7, 355–364.
- ROBBA, L., RUSSELL, S. J., BARKER, G. L. & BRODIE, J. (2006). Assessing the use of the mitochondrial *cox1* marker for use in DNA barcoding of red algae (Rhodophyta). *American Journal of Botany*, 93, 1101–1108.
- RUBINOFF, D. (2006). DNA barcoding evolves into the familiar. *Conservation Biology*, 20, 1548–1549.
- RUBINOFF, D., CAMERON, S. & WILL, K. (2006). A genomic perspective on the shortcomings of mitochondrial DNA for “barcoding” identification. *Journal of Heredity*, 97, 581–594.
- SASAKI, H., LINDSTROM, S. C., WAALAND, J. R. & KAWAI, H. (2003). Occurrence of the gametophyte of *Agarum clathratum* (Laminariales, Phaeophyceae) as an endophyte in *Orculifilum denticulatum* (Gigartinales, Rhodophyceae). *Phycological Research*, 51, 192–202.
- SAUNDERS, G. W. (2005). Applying DNA barcoding to red macroalgae: a preliminary appraisal holds promise for future applications. *Philosophical Transactions of the Royal Society B Series*, 360, 1879–1888.
- SAUNDERS, G. W. (2008). A DNA barcode examination of the red algal family Dumontiaceae in Canadian waters reveals substantial cryptic species diversity. 1. The foliose *Dilsea-Neodilsea* complex and *Weeksia*. *Botany*, 86, 773–789.

- SAUNDERS, G. W. (2009). Routine DNA barcoding of Canadian Gracilariales (Rhodophyta) reveals the invasive species *Gracilaria vermiculophylla* in British Columbia. *Molecular Ecology Resources*, 9, (Suppl. 1), 140–150.
- SCHINDEL, D. E. & MILLER, S. E. (2005). DNA barcoding a useful tool for taxonomists. *Nature*, 435, 17.
- SEBERG, O. & PETERSEN, G. (2009). How many loci does it take to DNA barcode a *Crocus*? *PLoS ONE*, 4, e4598. doi: 10.1371/journal.pone.0004598.
- SHAFFER, H. B. & THOMPSON, R. C. (2007). Delimiting species in recent radiations. *Systematic Biology*, 56, 896–906.
- SHERWOOD, A. R., VIS, M. L., ENTWISLE, T. J., NECCHI, O. JR. & PRESTING, G. G. (2008). Contrasting intra versus interspecies DNA sequence variation for representatives of the Batrachospermales (Rhodophyta): Insights from a DNA barcoding approach. *Phycological Research*, 56, 269–279.
- STEINKE, D., VENCES, M., SALZBURGER, W. & MEYER, A. (2005). TaxI: a software tool for DNA barcoding using distance methods. *Philosophical Transactions of the Royal Society of London B Biological Sciences*, 360, 1975–1980.
- STIGER, V., HORIGUCHI, T., YOSHIDA, T., COLEMAN, A. W. & MASUDA, M. (2000). Phylogenetic relationships of *Sargassum* (Sargassaceae, Phaeophyceae) with reference to a taxonomic revision of section Phyllocystae based on ITS-2 nrDNA sequences. *Phycological Research*, 48, 251–260.
- SWOFFORD, D. L. (2002). *PAUP*: Phylogenetic Analysis Using Parsimony (*and Other Methods)*. Version 4. Sunderland MA: Sinauer Associates.
- TAI, V., LINDSTROM, S. C. & SAUNDERS, G. W. 2001. Phylogeny of the Dumontiaceae (Gigartinales, Rhodophyta) and associated families based on SSU rDNA and internal transcribed spacer sequence data. *Journal of Phycology*, 37, 184–196.
- TREWICK, S. A. (2008). DNA Barcoding is not enough: mismatch of taxonomy and genealogy in New Zealand grasshoppers (Orthoptera: Acrididae). *Cladistics*, 24, 240–254.
- UWAI, S., KOGAME, K., YOSHIDA, G., KAWAI, H. & AJISAKA, T. (2009). Geographical genetic structure and phylogeography of the *Sargassum horneri/filicinum* complex in Japan, based on the mitochondrial *cox3* haplotype. *Marine Biology*, 156, 901–911.
- VAN OPPEN, M. J. H., OLSEN, J. L., STAM, W. T., VAN DEN HOEK, C. & WIENCKE, C. (1993). Arctic–Antarctic disjunctions in the benthic seaweeds *Acrosiphonia arcta* (Chlorophyta) and *Desmarestia viridis/willii* (Phaeophyta) are of recent origin. *Marine Biology*, 115, 381–386.
- VOGLER, A. P. & MONAGHAN, M. T. (2006). Recent advances in DNA taxonomy. *Journal of Zoological Systematics and Evolutionary Research*, 45, 1–10.
- WARD, R. D., HOLMES, B. H. & O'HARA, T. D. (2008). DNA barcoding discriminates echinoderm species. *Molecular Ecology Resources*, 8, 1202 – 1211.
- WARD, R. D., HANNER, R. & HEBERT, P. D. N. (2009). The campaign to DNA barcode all fishes, FISH-BOL. *Journal of Fish Biology*, 74, 329–356.
- WILL, K. W. & RUBINOFF, D. (2004). Myth of the molecule: DNA barcodes for species cannot replace morphology for identification and classification. *Cladistics*, 20, 47–55.
- WOMERSELY, H. B. S. (1964). The morphology and taxonomy of *Cystophora* and related genera (Phaeophyta). *Australian Journal of Botany*, 12, 53–110.
- WOMERSLEY, H. B. S. (1987). *The marine benthic flora of southern Australia. Part II*. Adelaide, Australia: South Australian Government Printing Division.
- YANG, E. C., KIM, M. S., GERALDINO, P. J. L., SAHOO, D., SHIN, J-H. & BOO, S. M. (2008). Mitochondrial *cox1* and plastid *rbcL* genes of *Gracilaria vermiculophylla* (Gracilariaceae, Rhodophyta). *Journal of Applied Phycology*, 20, 161–168.
- YOTSUKURA, N., SHIMIZU, T., KATAYAMA, T. & DRUEHL, L. D. (2009). Mitochondrial DNA sequence variation of four *Saccharina* species (Laminariales, Phaeophyceae) growing in Japan. *Journal of Applied Phycology*. doi: 10.1007/s10811-009-9452-7.

- ZUCCARELLO, G. C. & LOKHORST, G. M. (2005). Molecular phylogeny of the genus *Tribonema* (Xanthophyceae) using *rbcL* gene sequence data: monophyly of morphologically simple algal species. *Phycologia*, 44, 384–392.
- ZUCCARELLO, G. C., BUCHANAN, J. & WEST, J. A. (2006). Increased sampling for inferring phylogeographic patterns in *Bostrychia radicans*/*B. moritziana* (Rhodomelaceae, Rhodophyta) in the Eastern USA. *Journal of Phycology*, 42, 1349–1352.
- ZWICKL, D. J. (2006). *Genetic algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion*. Unpublished Ph.D. dissertation. Austin TX: The University of Texas at Austin.

Chapter 5

**Taxonomic oversplitting in the New Zealand
members of the brown algal genus *Cystophora*
(Phaeophyceae, Sargassaceae): Synonymisation of
two species**

5.1 Abstract

Six species of *Cystophora* are currently recognized in New Zealand waters. Molecular evidence supports four of these (*C. platylobium* (Mertens) J. Agardh, *C. retroflexa* (Labillardière) J. Agardh, *C. scalaris* J. Agardh and *C. torulosa* (R. Brown) J. Agardh), but data have failed to separate two currently accepted species from sister species: *Cystophora congesta* Womersley & Nizamuddin is a form of *C. retroflexa*, with short laterals and fascicled ramuli, and represents part of a range of morphological variation; *Cystophora distenta* J. Agardh is a form of *C. scalaris*, with wide, compressed receptacles, but receptacle dimensions do not separate species. I synonymise *C. congesta* with *C. retroflexa* and *C. distenta* with *C. scalaris* on the basis of available molecular and morphological data, and provide amended descriptions of these species and a key to New Zealand species of *Cystophora*.

5.2 Introduction

ALGAE OF THE GENUS *CYSTOPHORA* are a significant component of the New Zealand marine flora (Shears & Babcock 2007). *Cystophora* thalli are common in the shallow sub-tidal, rock pools and lower intertidal. They provide structure and shelter for other marine organisms (Taylor & Cole 1994, McDermott & Shima 2006), and are a significant contributor to primary productivity (Schiel 1990).

Six species of *Cystophora* are currently recognised in New Zealand (Womersley 1964, 1987, Adams 1994). *Cystophora torulosa* (R. Brown) J. Agardh and *C. retroflexa* (Labillardière) J. Agardh are widespread in New Zealand and southern Australia. *Cystophora platylobium* (Mertens) J. Agardh is found from Cook Strait southwards and in southern Australia. *Cystophora scalaris* J. Agardh is found south of Cook Strait in New Zealand only. *Cystophora congesta* Womersley & Nizamuddin was described from Robe, South Australia, in 1964 and has been occasionally reported from New Zealand (Adams 1994, Nelson *et al.* 1992). Early collections from New Zealand include specimens determined as *C. dumosa* (Greville) J. Agardh, a name rejected as a *nomen confusum* by Womersley (1964), under the then current Montreal Code (Lanjouw *et al.* 1961) Art. 69 (although reported by Womersley as Art. 66). Womersley determined herbarium material used for the original description

(Sonder 1846) consisted of a mixture of species, and synonymised some of the material with *C. monilifera*. New Zealand specimens might be referable to *C. congesta* (Adams 1994). *Cystophora distenta* J. Agardh has been reported from the Chatham Islands (Agardh 1870, Lindauer *et al.* 1961) and other southern New Zealand locations (Nelson *et al.* 1991, 1992, Adams 1994). A detailed taxonomic history of *Cystophora* is provided by Womersley (1964, 1987).

Lindauer *et al.* (1961) also report doubtful records of four Australian species (including *Cystophora cephalornithos*, now *Caulocystis*) collected from New Zealand. These appear to be many mistakes in identification or incorrect recording of localities on early collections. Difficulties arise in species identification in *Cystophora* (Adams 1994). Lindauer *et al.* (1961) suggested two reasons for this: the state of dried herbarium material, which is often poorly preserved and incomplete, and because “within the New Zealand species a degree of hybridization occurs in certain areas where some of the species overlap” (p. 295). Lindauer *et al.* (1961) only consider *C. platylobium* as distinctive, as other species intergrade. Womersley (1964) also noted that the morphology of some specimens were intermediate between species. Further confusion has arisen from unclear synonymy and inadequate descriptions. Lindauer *et al.* (1961) describe specimens under the name *Cystophora dumosa*. Womersley (1964) rejected that name, and created a new species, *C. congesta*, containing many specimens previously assigned to *C. dumosa*. Adams’ (1994, p. 141) illustration of *C. congesta* does not resemble Womersley’s description, and keys in Lindauer *et al.* (1961) and Adams (1994) use different characters from Womersley (1964).

Cystophora congesta and *C. distenta* are rarely reported outside of the taxonomic literature. Surveys of the New Zealand coastlines by Morton & Miller (1968) and Shears & Babcock (2007) recorded *Cystophora platylobium*, *C. retroflexa*, *C. scalaris* and *C. torulosa*, but not *C. congesta* or *C. distenta*. A molecular study of the genus (Chapter 4) failed to separate *C. congesta* from *C. retroflexa* and *C. distenta* from *C. scalaris*. Here I evaluate the evidence for recognising six species of *Cystophora* in New Zealand, provide additional morphological and molecular data, and formally synonymise the species *C. congesta* and *C. distenta*.

Table 5.1. Voucher numbers of specimens used in analyses, collection locations, latitude and longitude. A. *Cystophora retroflexa* and outgroups with haplotypes and GenBank accession numbers of mitochondrial spacer sequences.

Herbarium voucher number ⁷	Species	Location	Latitude, Longitude	Haplotype (see Fig. 1) and GenBank Accession number
WELT A031531	<i>C. retroflexa</i>	Depot Beach NSW	35°37'37.91"S, 150°19'37.27"E	C (JF309054)
WELT A030917	<i>C. retroflexa</i>	Island Bay, Wellington, NZ	41°20'51.22"S, 174°46'9.77"E	B (JF309069)
WELT A031510	<i>C. retroflexa</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	B (JF309057)
WELT A031532	<i>C. retroflexa</i>	Mimosa Rocks, Arragunnu NSW	36°35'2.07"S, 150° 3'15.18"E	C (JF309056)
WELT A031533	<i>C. retroflexa</i>	Point Lonsdale VIC	38°17'20.52"S, 144°36'4.42"E	D (JF309059)
WELT A031509	<i>C. retroflexa</i>	Port Campbell VIC	38°37'14.84"S, 142°59'32.69"E	C (JF309060)
WELT A030926	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43. 05"S, 174°47'14.08"E	B (JF309051)
WELT A030914	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	B (JF309066)
WELT A030913	<i>C. retroflexa</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	B (JF309067)
WELT A031511	<i>C. retroflexa</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	C (JF309061)
WELT A031508	<i>C. retroflexa</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	A (JF309062)

⁷ Field numbers are shown. These specimens have been deposited at WELT. Herbarium numbers will be added when the accession process is completed.

Table 5.1 (Continued)

Herbarium voucher number	Species	Location	Latitude, Longitude	Haplotype (see Fig. 1) and GenBank Accession number
WELT A031512	<i>C. retroflexa</i>	Robe SA	37° 9'23.33"S, 139°44'56.32"E	C (JF309070)
WELT A031530	<i>C. retroflexa</i>	Sorrento, VIC	38°20'5.35"S, 144°44'33.96"E	D (JF309058)
WELT A030927	<i>C. retroflexa</i>	The Sirens, Wellington NZ	41°20'56.91"S, 174°45'51.05"E	B (JF309063)
WELT A030932	<i>C. retroflexa</i>	Tokomaru Bay, Eastland, NZ	38° 6'35.83"S, 178°21'18.27"E	D (JF309068)
WELT A031528	<i>C. subfarcinata</i>	Lonsdale VIC	38°17'5.30"S, 144°36'54.74"E	F (JF309065)
WELT A031513	<i>C. subfarcinata</i>	Observation Head, Batemans Bay NSW	35°43'45.50"S, 150°12'30.26"E	E (JF309055)
WELT A031527	<i>C. subfarcinata</i>	Squeaky Beach, Wilsons Promontory VIC	39° 1'45.89"S, 146°18'24.24"E	F (JF309063)
WELT A031529	<i>C. subfarcinata</i>	Ulladulla, NSW	35°21'37.65"S, 150°29'5.55"E	E (JF309053)

Table 5.1. B. *Cystophora scalaris* samples with reference numbers to Fig. 5.1.

Herbarium voucher number	Species	Location	Latitude, Longitude	Reference number (Fig. 5.1)
WELT A030925	<i>C. scalaris</i>	Pukerua Bay NZ	41° 1'28.36"S, 174°54'5.54"E	1
WELT A030921	<i>C. scalaris</i>	Riversdale, Wairarapa NZ	41° 6'24.49"S, 176° 4'10.73"E	2
WELT A031338	<i>C. scalaris</i>	Evans Bay, Wellington, NZ	41°18'59.61"S, 174°48'23.25"E	3
WELT A031337	<i>C. scalaris</i>	Lyall Bay, Wellington, NZ	41°20'12.67"S, 174°47'35.14"E	4
WELT A030902	<i>C. scalaris</i>	Moa Point, Wellington, NZ	41°20'37.68"S, 174°48'31.54"E	5
WELT A030904	<i>C. scalaris</i>	Moa Point, Wellington, NZ	41°20'37.68"S, 174°48'31.54"E	6
WELT A030901	<i>C. scalaris</i>	Tarakena Bay, Wellington, NZ	41°20'39.95"S, 174°49'12.46"E	7
WELT A030903	<i>C. scalaris</i>	Princess Bay, Wellington NZ	41°20'43.05"S, 174°47'14.08"E	8
WELT A031341	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	9
WELT A031344	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	10
WELT A030905	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	11
WELT A030923	<i>C. scalaris</i>	Te Raekaihau, Wellington NZ	41°20'50.92"S, 174°47'29.01"E	12
WELT A031336	<i>C. scalaris</i>	Red rocks, Wellington, NZ	41°20'54.91"S, 174°44'30.19"E	13

Table 5.1. B. (Continued)

WELT A031334	<i>C. scalaris</i>	Red rocks, Wellington, NZ	41°20'54.91"S, 174°44'30.19"E	14
WELT A031335	<i>C. scalaris</i>	Red rocks, Wellington, NZ	41°20'54.91"S, 174°44'30.19"E	15
WELT A031346	<i>C. scalaris</i>	Kaingaroa, Chatham Islands	43°43'51.20"S, 176°16'2.26"W	16
WELT A031345	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	17
WELT A030909	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	18
WELT A030908	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	19
WELT A030910	<i>C. scalaris</i>	Point Weeding, Chatham Islands	43°56'32.08"S, 176°34'14.74"W	20
WELT A030911	<i>C. scalaris</i>	Waitangi, Chatham Islands, NZ	43°56'41.58"S, 176°33'41.78"W	21
WELT A030922	<i>C. scalaris</i>	Ocean Beach, Jackson Bay NZ	43°57'54.44"S, 168°36'21.70"E	22
WELT A031343	<i>C. scalaris</i>	Owenga, Chatham Islands	44° 2'0.30"S, 176°19'58.80"W	23
WELT A031340	<i>C. scalaris</i>	Manukau Reef, Chatham Islands	44° 2'4.90"S, 176°19'44.31"W	24
WELT A030933	<i>C. scalaris</i>	Mapoutahi, Otago Peninsula NZ	45°44'5.57"S, 170°37'4.53"E	25
WELT A030906	<i>C. scalaris</i>	Wellers Rock, Otago Harbour, NZ	45°47'51.97"S, 170°42'53.85"E	26

5.3 Materials and Methods

Cystophora specimens were collected from around New Zealand, including the Chatham Islands, and from southern Australia and Tasmania (Table 5.1). *A priori* species assignments were made on the basis of morphological characters from Womersley (1964, 1987). Thalli were pressed as herbaria specimens and one or a few ramuli were cleaned of epiphytes and the tissue rapidly dried and stored in silica gel for DNA extraction. In addition, specimens held in WELT were examined (Table 5.1).

Length, width (wider axis of compressed receptacle) and breadth (narrower axis of compressed receptacle) of receptacles of herbaria specimens of *C. scalaris* and *C. distenta* were measured with digital callipers. Ten randomly selected receptacles were measured from 26 specimens. In three specimens, selected as representing the range of receptacle size, measurements were taken of fresh receptacles. These receptacles were then air dried and remeasured, and a factor estimated by regression to determine a multiplier to convert measurements from herbarium specimens to the equivalent fresh dimensions. ANOVAs, implemented in SPSS16, were used to test for significant differences between mean dimensions. In addition to these specimens, herbaria specimens of all New Zealand species of *Cystophora* were examined from WELT and my collections and additional measurements were made to determine the range of character dimensions for revised descriptions.

DNA was extracted using a modified CTAB extraction (Zuccarello & Lokhorst 2005), with the addition of 1% polyvinylpyrrolidone to the extraction buffer. The 23S-tRNA-Lys mitochondrial spacer (Hoarau *et al.* 2007, Chapter 2) was amplified. Primers and PCR mixes and routines were described in Chapter 2. PCR products were cleaned with ExoSAP-IT (USB, Cleveland, Ohio) enzymes and sequenced commercially (Macrogen Inc., Seoul, Korea). A statistical parsimony network was inferred from mitochondrial spacer sequences from 20 specimens of *Cystophora* using TCS software (Clement *et al.* 2000), with linkages made at the 95% confidence level.

5.4 Results

Some specimens were difficult to assign to species using existing keys and morphological characters. Ambiguous specimens were assigned to the more common species. Consequently only seven specimens were assigned to *C. congesta* and four

specimens to *C. distenta* (Table 5.1). Measurements of receptacles of putative *Cystophora scalaris* and *C. distenta* specimens show a wide range of receptacle forms, often with high variation on a single thallus (Fig. 5.1). Specimens from highly exposed locations (e.g., specimens 10–13 in Fig. 5.1) show very reduced receptacle length. Neither length nor width:breadth ratios (a measure of compression of the receptacle) show clusters consistent with separate species (Fig. 5.1).

Fig. 5.1. Morphological variation from 26 specimens of *Cystophora scalaris*. Graphs show maximum, minimum median and 25% and 75% quartiles from 26 specimens. Data are arranged from northernmost (1) to southernmost (26) specimens, asterisks indicate specimens from the Chatham Islands. A. Receptacle length. B. Receptacle width: breadth ratios.

ANOVA analysis of log-transformed data showed significant ($P < 0.01$) differences between means in the data, but post-hoc tests (Tukey's HSD pairwise and

range tests) showed significant differences only existed between outlier data. For example, mean length of receptacles from specimens 5 and 18 (in Fig. 5.1a) were significantly different from specimens 8–13, but differences between both groups and intermediate specimens was not significant ($P > 0.05$). Similarly, specimens with the greatest receptacle width:breadth ratios (specimens 5 and 16 in Fig. 5.1A) showed significant differences from those specimens with the lowest ratios (specimens 12, 23 and 26 in Fig. 5.1B), but none of these specimens showed significant differences from intermediate specimens.

Molecular data

Four mitochondrial spacer haplotypes were found in *C. retroflexa* and *C. congesta* (Fig. 5.2, A–D; Appendix 6). Two haplotypes associated with *C. subfarcinata* were included as outgroups (E, F). Four *C. retroflexa*/*C. congesta* haplotypes were separated by a single change in each case. Two haplotypes were sampled in specimens of both *C. congesta* and *C. retroflexa*. Only one specimen of *C. congesta* (WELT A031508 from Robe, South Australia) had a unique haplotype (Haplotype A in Fig. 5.2).

Fig. 5.2. Statistical parsimony network inferred from 20 mitochondrial spacer sequences of *Cystophora retroflexa* (A–D) and *C. subfarcinata* (E, F). Estimated at 95% confidence. Haplotypes A–C include both *C. retroflexa* and *C. congesta* morphotypes. Numbers are sample frequencies.

5.5 Discussion

Two species of *Cystophora* found in New Zealand do not present any taxonomic problems. *Cystophora platylobium* is usually found in relatively deep water, and collections are often based on drift specimens. Attached specimens were collected in southern Otago only (at 1–2 metres depth). For this reason, sampling of this species was limited, but available data suggests this species is ecologically, morphologically and genetically (Chapter 4) distinct from other species, and New Zealand specimens are genetically very similar to Australian specimens, with identical *cox1* haplotypes (Chapter 4). *Cystophora torulosa* is also morphologically and ecologically and genetically distinct from other species (Chapter 4). No difficulty was found in assigning specimens to these species using existing keys and descriptions. These species will not be commented on further.

The remaining New Zealand species of *Cystophora* present some problems. Specimens of *Cystophora congesta* can be difficult to separate from *C. retroflexa*, and *C. scalaris* can be difficult to separate from *C. distenta*. Some confusion also appears to have arisen in distinguishing *C. scalaris* and *C. retroflexa* (Lindauer *et al.* 1961). These problems arise in part from inadequate descriptions and poor character choice in the keys provided in Lindauer *et al.* (1961) and Adams (1994), which rely on the relative zig-zaggedness of the main axes and laterals for species identification. This is confusing, as from my observations, the degree of zig-zagging in the primary and secondary axes is fairly plastic in several species of *Cystophora* and is probably related to the growth rate (Klemm & Hallam 1987). Womersley's (1964) keys separated these species by the arrangement of branching of the ramuli (complanate in *C. scalaris* and *C. distenta*, irregularly radial in *C. retroflexa* and *C. congesta*) and the size and form of the receptacles, characters which are reliable for separating *C. retroflexa* from *C. scalaris*, but still inadequate for separating both *C. retroflexa* from *C. congesta* and *C. scalaris* from *C. distenta*. Intermediate forms are common in both pairs of species and I consider the morphological characters used to split species are not discontinuous, but rather lie on a continuum of intra-specific variation. The evidence for maintaining *C. congesta* and *C. distenta* as separate species can be attributed to limited sampling, with specimens examined representing the extremes of a morphological continuum.

Fig. 5.3. Morphological variation in *Cystophora retorta*. A. congested form, zig-zag laterals [WELT A031546]; B. distended form, straight laterals [WELT A031547]. Scale bar = 100 mm.

5.5.1 *Cystophora congesta* and *C. retroflexa*

In Chapter 4 I could not separate *C. retroflexa* and *C. congesta* using ITS and *cox1* sequences. The highly variable 23S-tRNA-Lys mitochondrial spacer (Fig. 5.2) also fails to separate these species. Both this study and Chapter 4 include *C. congesta* specimens from the type locality (Robe, South Australia). While sampling in this study is limited, the mitochondrial spacer used here is highly variable in most species of brown algae (Hoarau *et al.* 2007, Chapter 2), and haplotypes found in all but one specimens assigned to *C. congesta* were also found in typical specimens of *C. retroflexa*. In addition, these species share *cox1* haplotypes and ITS ribotypes (Chapter 4). Together, this is strong evidence against genetic differentiation.

According to Womersley (1964: 87) “*C. congesta* is most closely related to *C. retroflexa*, differing in the much denser laterals and secondary axes, shorter receptacles and thicker, more rigid, primary axis. Occasional intergrades between these species occur.” Womersley then separated the species entirely on these vegetative characters, all of which are variable in other species of *Cystophora*. For example, specimens of *C. retorta* can have a thick primary axis and zig-zag laterals bearing fascicled ramuli and receptacles, and other specimen have a thin primary axis, with straight and sparsely branched laterals (Fig. 5.3). All morphological types share the scattered bisexual conceptacles that distinguish this species from its near relatives (Womersley 1964).

Some specimens could be easily assigned to *C. retroflexa* or *C. congesta* using Womersley’s criteria, but many specimens were intermediate and could be assigned to either species (Fig. 5.4). The plasticity of these characters in other species of *Cystophora*, for example in *C. siliquosa* and *C. subfarcinata* (see dimensions in Womersley 1964), the presence of intergrades and the absence of genetic differentiation suggest these characters are inadequate for inferring distinct species.

The weight of the evidence is that *C. congesta* should be regarded as either an ecotype or a developmental form of *C. retroflexa*. I consider *C. congesta* Womersley & Nizamuddin as synonymous with *C. retroflexa* (Labillardière) J. Agardh, the latter name having priority.

Fig. 5.4. Morphological variation in *Cystophora retroflexa*. **A.** Distended thallus [WELT A031534]. **B.** Intermediate thallus [WELT A031533]. **C.** Congested thallus [WELT A031508]. Scale bar = 100 mm.

Fig. 5.5. Morphological variation in receptacles of *Cystophora scalaris* A. WELT A031345A; B. WELT A031345B; C. WELT A030922; D. WELT A031346; E. WELT A031340; F. WELT A031343; G. WELT A031341; H. Examples of receptacles from a single thallus, WELT A031337). Scale bar = 20 mm.

5.5.2 Amended description

Cystophora retroflexa (Labillardière) J. Agardh *Species Genera et Ordines Algarum* 1848: 242. Womersley *Aus. J. Bot.* 1964: 89, Figs. 31, 32, pl. 10, Fig. 2. *Marine Flora of Southern Australia* 3 1987: 396, Figs. 132 I, 146B, 149B, C.

Cystophora congesta Womersley & Nizamuddin ex Womersley *Aus. J. Bot.* 1964: 86, Fig. 30, pl. 9 Fig. 2. Womersley *Marine Flora of Southern Australia* 3 1987: 394. Figs. 146A, 149A.

Figs. 4, 5.

Thallus brown, 0.3–1.5 (–2) m high, with the primary axes bearing alternate laterals and secondary axes. *Holdfast* discoid–conical, 3–15 mm diameter, with a short, terete stipe; epilithic. *Primary axes* compressed, 3–12 mm broad below and 1–4(–5) mm thick, 2–6(–10) mm broad above, narrow edged, fairly straight to zig-zag and scalariform, alternately distichously branched from the face of the axes, lower parts often denuded with prominent residues, each branch position giving rise to 1(–3) determinate laterals and 1(–2) indeterminate laterals; lower laterals usually retroflex, often with broad basal wings. *Laterals* determinate or indeterminate, determinate laterals arising from the edge of an outgrowth of the main axes, 20–100 mm long, with usually irregular radial, sometimes subtristichous or subdistichous branching, bearing ramuli, sometimes developing into receptacles, lower determinate laterals often with enlarged and flattened ramuli. Indeterminate laterals arising from the centre of outgrowths from the axes, 20–300 mm long, initially terete, becoming flattened, alternately distichously branched, becoming slightly to strongly zig-zag; bearing determinate secondary laterals and/or secondary indeterminate laterals. *Ramuli* 20–40(–50) mm long, 0.4–2(–3) mm broad. Ramuli on lower laterals sometimes enlarged and flattened. *Vesicles* ovoid, becoming subspherical when old, replacing basal (or lower 2) ramuli of laterals, stalked, mutic, 4–12(–15) mm long and 3–10(–12) mm broad.

Reproduction thalli monocious. Receptacles usually simple, occasionally branched, compressed, margins undulate to torulose when fresh, often strongly torulose when dried, 15–50 (–60) mm long and 1–2 (–4) mm broad, attenuate at apex usually with sterile terminal awn, awn rarely with scattered conceptacles, with 2 rows of conceptacles along the margins; conceptacles bisexual, ostioles along the margin.

Type locality “Cape van Diemen” (southern Tasmania).

Holotype FI (Herbarium Webbianum).

Distribution Australia: Womersley (1964) reports the species from Bondi (Sydney), NSW. I found specimens from Kiama, NSW southwards. I have not ascertained the western limit of *C. retroflexa* in Australia. Womersley (1987) reports the western limit as “from Cottesloe to Nannarup (Albany), W.A., and from Kangaroo I.”. Tasmania, New Zealand, Chatham Is., Auckland Is.

5.5.3 *Cystophora distenta* and *C. scalaris*

J. G. Agardh (1870) described *C. scalaris* and *C. distenta* from the Chatham Islands, but regarded them as contracted and distended forms of the same species, stating: “*C. scalarem* & *C. distentam* esse ejusdem specie formas – unam contractam, alteram distentam” (Agardh 1870). [“*C. scalaris* and *C. distenta* are forms of the same species – one contracted and the other distended.”]. De Toni (1895: 141) also stated that *C. distenta* was “Anne forma laxa *C. scalaris* J. Ag.?”

Treatment as separate species seems to begin with Laing (1899) who reported both forms as species without comment, and continued to do so in further publications (Laing 1926). Lindauer (1947) followed Laing in listing these as separate species (using the generic name *Blossevillea* (following Gardner 1913). Lindauer (1947) gave the distribution of *C. distenta* (as *Blossevillea distenta*) as Chatham Islands and Stewart Island, but noted that in Stewart Island “the habit of the plant is less typical” (p. 560). Lindauer *et al.* (1961) reported the two entities as separate species, as did Womersley (1964, 1987), Nelson *et al.* (1991, 1992) and Adams (1994).

Womersley (1964: 91) described *C. distenta* and *C. scalaris* as separate species, “differing in the more compressed and much longer receptacles, straighter and less robust axis, and looser branching ” [in *C. distenta*]. This is unhelpful as the dimensions of the axis and looseness of branching are variable in many species of *Cystophora* (for example, in *C. retorta*, Fig. 5.3), and receptacle dimensions provided by Womersley (1964) and Lindauer *et al.* (1961) for these species overlap. Receptacles are very variable in length in *C. scalaris* (Figs. 5.1 and 5.5), and from my observations compression appears to be related to the reproductive state of the

receptacle, rather than any underlying genetic difference. Receptacle dimensions are highly variable, even on a single thallus (Figs. 5.1 and 5.5).

Cystophora distenta was first described from the Chatham Islands by J. Agardh (1870). De Toni listed its distribution as the Chatham Islands and Bluff. Lindauer *et al.* (1961) extended its distribution to include Stewart Island and Frenchman's Pass (presumably French Pass in the Marlborough Sounds), and also noted that specimens were found in drift at Tauranga. Later specimens were reported from Cape Palliser on the southern North Island (Adams 1994) and Durville Island, on the northern tip of the South Island (Nelson *et al.* 1992). The species appears to be distributed on the mainland south of Cook Strait and on Stewart Island and the Chatham Islands. This distribution is the same as that of *C. scalaris* (Lindauer *et al.* 1961, Adams 1994).

As I am unable to separate *C. scalaris* and *C. distenta* using morphological or molecular data, I consider Agardh's (1870) assessment to be correct and regard *C. distenta* as a form of *C. scalaris*. The latter name is in more common usage (e.g., Morton & Miller 1968, Shears & Babcock 2007).

5.5.4 Amended species description

Cystophora scalaris J. Agardh *Öfversigt af Kongl. Vetenskaps-Academiens*

Förhandlingar 1870: 442. Womersley *Aus. J. Bot.* 1964: 82, Figs. 22, 23, pl. 7, fig. 2.

Figs. 5.5, 5.6.

Thallus brown, (0.1–)0.3–1 m long, with the primary axes bearing alternate, openly to densely branched laterals and secondary axes. *Holdfast* discoid-conical, 5–20 mm diameter, with a short, terete stipe; epilithic. *Primary axes* strongly flattened, 3–15 (–20) mm broad and 2–3 mm thick, strongly zigzag and scalariform to fairly straight, lower parts often denuded with scalariform residues, alternately distichously branched from the face of the axis, with each branch position producing 1(–2) indeterminate secondary axes from the centre of the face and/or (0–1) –2 determinate laterals from the edge of the scales, laterals 10–30 (–50) mm apart; secondary and tertiary axes similar, 20–80 cm long, initially terete, becoming flattened, strongly zig-zag and scalariform to fairly straight; lower parts of primary, secondary, and older tertiary axes usually denuded, with very prominent, close-set scalariform residues, and

markedly broader basal wings on the primary (and often secondary) axis at branch positions. *Laterals* 20–80 (–100) mm long, complanately branched, often bifurcate at their base, then alternately distichous with ramuli also alternately distichous, with acute to rounded axils; ultimate ramuli terete to compressed, 5–10 (–30) mm long, 1–3(–4) mm broad. Ramuli of lower laterals on main axis often enlarged and flattened. Vesicles sparse to numerous, replacing lower 1 to 2 or more ramuli of the lateral of each bifurcation, sub-spherical to slightly ovoid, stalked, mutic, 2–8 (–12) mm diameter.

Reproduction. Thalli monoecious, receptacles developed from ultimate ramuli, simple or occasionally branched, ovoid to compressed in section, 10–50 mm long, (0.5–)1–3 mm broad and 1–1.5 mm thick, elongate, margins undulate to strongly torulose. Conceptacles bisexual, ostioles in two rows along the margins of the receptacle, rarely in three rows.

Type locality Chatham Islands.

Lectotype LD (Herbarium Agardh, No. 1113).

Distribution New Zealand, from Wellington southwards; Chatham Islands, Auckland Islands.

Fig. 5.6. Organisation of ramuli and receptacles in *Cystophora*. Both specimens show indeterminate (upper) and determinate (lower) laterals. **A.** Complanate organisation in *C. scalaris*. **B.** Radial organisation in *C. retroflexa*. Scale bar = 20 mm.

5.6 Conclusion

Four species of *Cystophora* are present in New Zealand. These can be diagnosed with molecular and morphological characters. These species can be separated by the form and arrangement of receptacles (Table 5.2).

Table 5.2. Key to the New Zealand species of *Cystophora*

1. Branching from the edge of the main axes, receptacles leaf-like.....*C. platylobium*
1. Branching from the face of the axes, receptacles terete or flattened.....2
2. Receptacles swollen, smooth, terete with rounded apex, densely clustered on laterals; lower intertidal or shallow sub-tidal.....*C. torulosa*
2. Receptacles elongate, slightly to strongly flattened, apex blunt, pointed or with sterile awn, densely to openly clustered on laterals, sub-tidal.....3

3. Receptacles and ramuli arranged radially, usually narrow with long terminal awn*C. retroflexa*
3. Receptacles and ramuli distichous, forming complanate lateral branches; Receptacles slightly to strongly flattened, with blunt to acute tips.....*C. scalaris*

5.7 Acknowledgements

I thank the New Zealand Department of Conservation and Education NZ Study Abroad Scheme for funding assistance; Wendy Nelson, Tracy Farr, Kate Neill and the Department of Conservation Chatham Islands staff for assistance with collections and Alan Millar (National Herbarium of New South Wales) and Jenn Dalen (Museum of New Zealand Te Papa Tongarewa herbarium) for assistance with herbaria access.

5.8 References

- ADAMS, N. M. (1994). *Seaweeds of New Zealand: An illustrated guide*. Christchurch, New Zealand: Canterbury University Press.
- AGARDH, J. G. (1848). *Species, Genera et Ordines Algarum. Vol. I*. Lund, Sweden: C. W. K. Gleerup.
- AGARDH, J. G. (1870). Om Chatham-öarnes Alger. *Öfversigt af Kongl. Vetenskaps-Academiens Förhandlingar, Stockholm*, 5, 435–456.
- CLEMENT, M., POSADA, D. & CRANDALL, K. A. (2000). TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, 9, 1657–1659.
- DE TONI, G. B. (1895). *Sylloge algarum omnium hucusque cognitarum. Vol. III. Fucoideae*. Padua, Italy: Sumptibus auctoris.
- GARDNER, N. L. (1913). New Fucaceae. *University of California Publications in Botany*, 4, 317–374.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- KLEMM, M. F. & HALLAM, N. D. (1987). Branching pattern and growth in *Cystophora* (Fucales, Phaeophyta). *Phycologia*, 26, 252–261.
- LAING, R. M. (1899). Revised List of New Zealand Seaweeds. *Transactions and Proceedings of the New Zealand Institute*, 32, 57–40, plates V–VIII.
- LAING, R. M. (1926). A reference list of New Zealand Marine Algae. *Transactions of the New Zealand Institute*, 57, 126–185.
- LANJOUW, J., BAEHNI, C., ROBYNS, W., ROSS, R., ROUSSEAU, J., SCHOPF, J. M., SCHULZE, G. M., SMITH, A. C., VILMORIN, R. DE & STAFLEU, F. A. (1961). International code of botanical nomenclature, adopted by the ninth International Botanical Congress, Montreal, August 1959. *Regnum Vegetabile*, 23.
- LINDAUER, V. W. (1947). An annotated list of the brown seaweeds, Phaeophyceae, of New Zealand. *Transactions of the Royal Society of New Zealand*, 76, 542–566.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae. *Nova Hedwigia*, 3, 129–350.
- MCDERMOTT, C. J. & SHIMA, J. S. (2006). Ontogenetic shifts in microhabitat preference of the temperate reef fish *Forsterygion lapillum*: implications for population limitation. *Marine Ecology Progress Series*, 320, 259–266.

- MORTON J. E. & MILLER M. C. (1968). *The New Zealand Sea Shore*. London-Auckland: Collins.
- NELSON, W. A., ADAMS, N. M. & FOX, J. M. (1992). Marine Algae of the Northern South Island. A list of species. *National Museum of New Zealand Miscellaneous Series No. 26*. 1–79.
- NELSON, W. A., ADAMS, N. M. & HAY, C. H. (1991). Marine Algae of the Chatham Islands. *National Museum of New Zealand Miscellaneous Series No. 23*. 1–58.
- SCHIEL, D. R. (1990). Macroalgal assemblages in New Zealand: structure, interactions and demography. *Hydrobiologia*, 192, 59–76.
- SHEARS, N. T. & BABCOCK, R. C. (2007). *Quantitative description of New Zealand's shallow sub-tidal reef communities*. (Science for Conservation 280). Wellington, New Zealand: New Zealand Department of Conservation.
- SONDER, O. G. (1846). Algae L. Agardh. In: *Plantae Preissianae sive enumeratio plantarum quas in Australasia occidentali et meridionali-occidentali annis 1838–1841 collegit Ludovicus Preiss.* (Lehmann, C. Eds.), 2, 148–160. Hamburgi [Hamburg]: sumptibus Meissneri.
- TAYLOR, R. B. & COLE, R. G. (1994). Mobile epifauna on subtidal brown seaweeds in northeastern New Zealand. *Marine Ecology Progress Series*, 115, 271–282.
- WOMERSELY, H. B. S. (1964). The morphology and taxonomy of *Cystophora* and related genera (Phaeophyta). *Australian Journal of Botany*, 12, 53–110.
- WOMERSLEY, H. B. S. (1987). *The marine benthic flora of southern Australia. Part II*. Adelaide, Australia: South Australian Government Printing Division.
- ZUCCARELLO, G. C. & LOKHORST, G. M. (2005). Molecular phylogeny of the genus *Tribonema* (Xanthophyceae) using *rbcL* gene sequence data: monophyly of morphologically simple algal species. *Phycologia*, 44, 384–392.

Chapter 6

Species delimitation in brown algae – insights and applications to New Zealand brown algae.

6. 1 Abstract

Species delimitation in brown algae has traditionally proceeded by inferring genetic discontinuities from morphological characters, with additional data from culturing, crossing experiments, and studies of natural hybridisation. The development of molecular methods has provided an additional source of data, but few studies have proceeded to a phylogenetic concept of species. Rather, taxonomists have delimited species using a subjective assessment of available data. I review approaches to species delimitation in brown algae with reference to findings from previous chapters. Recent phylogeographic studies have produced a complex picture of species' history, with extensive changes in distribution, isolation and secondary contact, and a variety of speciation scenarios. I argue that this complexity means no single approach can delimit all species. Rather, an explicitly historical species concept is needed, with the history of species included in species descriptions.

6.2 Introduction

THE PREVIOUS CHAPTERS have investigated genetic diversity of *Carpophyllum* and *Cystophora* at different scales. Characterising genetic diversity is an essential task for biologists seeking to understand evolution, and different approaches have emerged to deal with the multiple scales of diversity. Phylogeneticists focus on historical relationships between species. Taxonomists use various data to infer discontinuities in genetic variation in order to delimit and describe species. Population geneticists study allele frequencies within and between populations. This thesis began with a detailed mitochondrial DNA phylogeography of *C. maschalocarpum* (Chapter 2), then an investigation into the relationship between two species of *Carpophyllum* (Chapter 3). I produced a broad phylogeny of *Cystophora* (Chapter 4), and dealt with the taxonomic separation of two pairs of species (Chapter 5). Here I review this work with reference to approaches to species delimitation in brown algae.

6.3 Species delimitation

Species are basic units of ecological studies and biodiversity assessments, and are a fundamental point of reference for nearly all biological science. Despite this, consistent delimitation of species continues to be problematic and sometimes contentious (Dayrat 2005, Will *et al.* 2005, De Queiroz 2007). The availability of genetic data and dissatisfaction with traditional methods (Tautz *et al.* 2003, Godfray *et al.* 2007) has prompted several attempts at developing objective methods of species delimitation (Sites & Marshall 2004, Shaffer & Thomson 2007, Pons *et al.* 2009, Wiens 2007). These include phylogenetic (tree-based) methods (Brower 1999, Templeton 2001, Wiens & Penkrot 2002) and population genetic methods, either estimating gene flow (Porter 1990) or genetic distance between populations (Good & Wake 1992, Highton 1990, 1995). Other approaches include DNA barcoding (Hebert *et al.* 2003a, Hebert & Gregory 2005) or deviation from expectations of a coalescence model (Pons *et al.* 2006).

How to delineate species is the central task of taxonomy. A second task, providing tools (names, descriptions, keys) for accurate species identification, is dependent on accurate species delineation. As the purpose of taxonomy changed from classification – the convenient ordering of life to meet the needs of biologists – to an attempt to explain biological reality, the focus has changed from phenetics to phylogenetics. Most taxonomists now accept a phylogenetic approach (Quicke 1993, Dayrat 2005). This theoretical position has been reinforced by the availability of molecular data that is easily analysed by phylogenetic methods. Taxonomy has become a study of history, specifically, the history of speciation. Reading history directly is difficult for most taxonomists, but is especially difficult for phycologists dealing with brown algae as the fossil record of brown algae is negligible (Silberfield *et al.* 2010), and well-preserved specimens amenable to ancient DNA work are almost unknown⁸. Therefore the history of speciation in brown algae must be inferred from extant data. Molecular data from brown algae has revealed hybridisation, cryptic speciation, intra-specific variation and incongruities between morphological and molecular data, increasing the complexity of defining species boundaries. An additional problem is applying

⁸ Dillehay *et al.* (2008) recovered well preserved 14 000-year-old fragments of seaweed from archaeological sites in South America so the possibility of ancient DNA work exists.

nomenclatural rules, which have a typological basis, to species delineated using molecular methods (Dayrat 2005). In the brown algae, species boundaries have often been described or suggested from molecular data, but authors have refrained from making appropriate nomenclatural changes.

Like other organisms, brown algal species have traditionally been delimited using morphological characters. Cryptic speciation, morphological plasticity, and complex life histories have often confounded morphological prescriptions, which have not always been congruent with other sources of information (e.g., molecular data, culturing, and natural and artificial crossing experiments). In addition, theoretical progress in formulating species concepts and methods of species delimitation has produced a more complex picture of species (Hey 2006, De Queiroz 2007, Mallet 2008). These approaches are discussed below.

6.4 Morphological approaches

Morphological approaches have been the mainstay of brown algal taxonomy (e.g., Agardh 1848, De Toni 1895, Setchell & Gardner 1925, Abbott & Hollenberg 1976, Fletcher 1987, Womersley 1987), and species delimitation using morphological characters has often been robust, despite rearrangements of higher taxonomic groups. Problems of morphological approaches often invoked (e.g., McDevit & Saunders 2009) include a limited number of characters, plasticity (Burrows & Lodge 1951, Mathieson *et al.* 1981, Henkel *et al.* 2007, Demes *et al.* 2009) and convergence (Harvey & Goff 2006, Jones *et al.* 2010, Tronholm *et al.* 2010a). Despite this, morphological characters continue to be given considerable weight, although very recent studies have prioritised molecular data (Macaya & Zuccarello 2010, Fraser *et al.* 2010b). Morphological methods are generally descriptive, although numerical taxonomic methods have been used in the brown algae (Russell & Fletcher 1975, Rice & Chapman 1985, Pérez-Ruzafa *et al.* 1993), but have largely been superseded by molecular data, with morphological characters mapped onto trees (e.g., De Clerck *et al.* 2006, Silberfield *et al.* 2010). Morphometric methods have also been used (e.g., Widdowson 1971, Rice 1989). Morphometrics are useful when used to determine diagnostic characters in

conjunction with molecular markers or culturing (Kraan *et al.* 2001, Roberson & Coyer 2004, Tronholm *et al.* 2010a).

In some genera, morphological plasticity has led to a plethora of synonyms and sub-species names. For example, there are 1046 names for species and varieties of *Fucus* and 869 names for *Sargassum* in Algaebase (Guiry & Guiry 2011). In *Cystophora*, morphological methods have resulted in oversplitting of some species and I have synonymised *C. scalaris*/*C. distenta* and *C. retroflexa*/*C. congesta*. Oversplitting probably results from limited sampling and a typological approach, which fail to capture a continuum of morphological variation. *A priori* concepts of species might result in conscious or unconscious collection bias for “typical” or “healthy” specimens, reinforcing selective sampling.

6.5 Culture experiments

Heteromorphic life histories and environmental plasticity complicates morphological studies. Culture studies can overcome some of these difficulties. Culture experiments have been used to test morphological characters. This is especially informative in species with heteromorphic life histories (e.g., Wynne 1969, Clayton 1976, 1978, Kogame 1996, 1997a, 1997b, Kogame & Yamagishi 1997) and multiple pathways in morphological development.

6.6 Crossing experiments and reproductive isolation

Complete reproductive isolation is often viewed as the ultimate step in speciation and the most fundamental test of species delimitation. Reproductive isolation can be tested directly, at least in species that are amenable to growth in culture, but is often inferred from morphological or phylogenetic data.

The Biological Species Concept (BSC, Mayr 1942, 1996) bases species delimitation on reproductive isolation, although most proponents allow some level of hybridization or gene flow (O’Brien & Mayr 1991, Coyne & Orr 2004). The BSC has a compelling logic, but becomes problematic where hybridization is

widespread⁹, such as in *Fucus* and *Carpophyllum*, and can be ahistorical (Velasco 2008). Reproductive isolation is usually inferred, generally from morphological discontinuities, and often with subjective values placed on different characters by taxonomists (Mayr 1996). Reproductive isolation can be tested directly using crossing experiments.

Crossing experiments have been used fairly extensively in a few genera of brown algae, especially *Ectocarpus* and some Laminariales. Often there has been a reluctance to formally apply results of crossing experiments to name new species or synonymise existing species, even where reproductive compatibility or isolation is clearly shown. For example, Müller (1988) declined to separate Chilean and Mediterranean isolates of *Ectocarpus siliculosus*, even after crosses showed hybrid sporophytes could not reproduce sexually and therefore could be considered separate species under the Biological Species Concept. In spite of supporting information (differences in development and morphology), Müller declined to separate these species “for practical reasons,” apparently because morphological diagnosis would be demanding. Conversely, Lewis *et al.* (1986) and Lewis & Neushul (1994) produced viable and fertile sporophytes from crosses of geographically separated *Macrocystis* specimens, then considered as separate species. However the authors considered that these should be maintained as separate species on the basis of morphological differences. In these cases the morphological species concept appears to have been given priority.

The complexity of crossing data was shown in *Alaria* (Kraan & Guiry 1998, 2000, Kraan *et al.* 2001), where crosses between some isolates of the same morphologically-determined species show limited viability, whereas distinct species can be fully interfertile. This is congruent with molecular data which shows greater genetic variation within *A. esculenta* compared to variation between other recognised *Alaria* species. Kraan & Guiry (2000) reject the morphological and biological species concepts for *Alaria*, and consider only a phylogenetic species concept is appropriate, but refrained from splitting reproductively isolated lineages into species.

⁹ O’Brien & Mayr (1991) accept hybridisation between biological species, as long as this does not “disintegrate the genetic integrity of the species”. It is not clear what this means for closely related species where gene flow from hybridisation is difficult to distinguish from incomplete lineage sorting.

Druehl *et al.* (2005) addressed some of the technical difficulties in conducting crossing experiments. Development of sporophytes in negative controls showed several Laminarean species are able to reproduce asexually via parthenogenesis or apogamy. Five species combinations produced putative hybrids in culture, but biparental inheritance of ITS DNA was confirmed in only one combination, raising doubt that progeny are hybrids. Druehl *et al.* (2005) recommend using molecular data (e.g., Coyer *et al.* 2002) to test hybrid status of cultured specimens. Parthenogenesis has also been reported in Ectocarpales (Bothwell *et al.* 2010), Chordariales (Peters 1987) and Fucales (Clayton *et al.* 1998) and might be common in brown algae. In addition to these problems, crossing experiments are limited to species that are amenable to growth in culture. My attempts to grow *Carpophyllum* in culture failed, and this difficulty has been reported by other workers (Dromgoole 1973, D. Schiel pers. comm.). This precludes performing artificial crosses, and investigating the fertility of hybrids. *Carpophyllum* hybrids form receptacles and zygotes, but their subsequent viability is unknown. One approach to these difficulties, and to determine whether potential for interspecific fertility demonstrated in culture is realised in nature, is to study natural hybrids.

6.7 Hybridisation as natural crossing experiments

Hybridisation appears to be fairly common in brown algae (Lewis 1996), and hybridisation can be regarded as a natural crossing experiment (Hewitt 1988), with assessments of the fitness and fertility of hybrids pertinent to the understanding of species limits (Arnold & Hodges 1995, Riesberg 1995).

Several problems arise in determining the prevalence of hybridisation and the importance of hybridisation to species boundaries. First, there are technical issues with identifying hybrids. Morphological intermediacy does not necessarily identify hybrids (Wilson 1992, Riesberg 1995) and Coyer *et al.* (2006b) showed morphologically similar salt marsh forms of *Fucus* could be interspecific hybrids or polyploid *F. vesiculosus*. A response to this problem is to identify hybrids by molecular markers, but Lane & Saunders (2005) showed that the presence of cryptic, epiphytic, gametophyte stages of Laminarialean species on sporophytes of related species could contaminate DNA samples, mimicking the signature of

hybrids. Wallace *et al.* (2004) identified *Fucus* hybrids by microsatellite admixture, but Engel *et al.* (2005) suggested this inference was confounded by inadequate characterisation of genetic variation in parent populations.

Secondly, hybridisation might be a consequence of taxonomic inflation rather than biological reality, where progeny of oversplit species are considered hybrids. *Macrocystis* species were considered to produce hybrids in culture (Lewis *et al.* 1986, Lewis & Neushul 1994), but recent studies have synonymised species (Westermeyer *et al.* 2007, Demes *et al.* 2009) and all forms of *Macrocystis* are now considered conspecific (Macaya & Zuccarello 2010).

Lastly, population genetic studies of hybrids are usually limited to one or a few populations (e.g., Wallace *et al.* 2004, Coyer *et al.* 2006b, 2006c, 2007). Different population of hybrids with the same parent species might be under different selective regimes and have different evolutionary trajectories (Butlin *et al.* 2008). Local hybrid zones might not be representative of patterns of hybridisation across species' ranges.

In vitro hybridisation has been widely reported in Laminariales (Liptack & Druehl 2000, Kraan & Guiry 2000), but natural hybrids might be rare (Druehl *et al.* 2005), although an extensive molecular survey of North Pacific *Alaria* species found evidence for widespread hybridisation (Lane *et al.* 2007). In this study there was no congruity between *cox1* clusters and morphology and no thresholds in genetic distance between groups. ITS sequences had many polymorphic sites and showed a continuum of variation rather than clustering as species level groups. Lane *et al.* (2007) suggest extensive hybridisation occurred after secondary contact between previously allopatric species, and ITS data indicates recombination following a breakdown of species barriers.

Extensive hybridisation raises the question of how species are maintained. In *Fucus*, where hybridisation is best studied (Scott & Hardy 1994, Coyer *et al.* 2002, Wallace *et al.* 2004, Billard *et al.* 2005, Coyer *et al.* 2006b, 2006c, Mathieson *et al.* 2006, Coyer *et al.* 2007, 2010), selfing acts to restrict introgression in some hermaphrodite species (Engel *et al.* 2005, Perrin *et al.* 2007). Selfing is correlated with height in the intertidal, with the inference that species in the upper intertidal have longer periods of emersion, and less time to disperse gametes, and gametes will often be lost from stands, and selfing is advantageous

under these conditions. Selection against hybrids should reinforce evolution of selfing or of mating barriers.

It is not clear how often species are lost through hybridization. This is unlikely to be obvious in the fossil record, even of species that leave good fossils, such as vertebrates or foraminiferans, as fusion of species may be difficult to distinguish from extinction of one lineage. In a changing environment (climate change, tectonics) a degree of allopatric divergence followed by secondary contact and fusion might be common (Butlin *et al.* 2008). Major changes in species ranges (Hoarau *et al.* 2007, Maggs *et al.* 2008, Fraser *et al.* 2009b) and demography (Graham *et al.* 2010) suggest a highly dynamic history of species distributions over the last 20 000 years. This in turn suggests a dynamic pattern of allopatry and sympatry (Coyer *et al.* 2010), where lineages might often diverge for a period, before reticulating following secondary contact.

In *Carpophyllum* hybridisation appears to be following different patterns in different areas. In Northland there is evidence for hybridisation between *C. angustifolium* and *C. maschalocarpum* that confounds phylogenetic methods. In the Bay of Plenty *C. angustifolium* and *C. maschalocarpum* lineages are distinguishable with nuclear markers, but *C. maschalocarpum* mitochondrial markers appear to have introgressed into populations of *C. angustifolium*. In *Carpophyllum*, all species are dioecious and sub-tidal. Strong selection might be responsible for the persistence of separate morphological species in areas of sympatry. In particular, *C. angustifolium* might be adapted for high wave exposure, with a strong terete stipe and a low drag morphology (Hodge 2009). I hypothesise that periodic extreme weather events would remove hybrids and *C. maschalocarpum* from exposed habitats, maintaining species boundaries. Further experimental work to determine the impacts of wave exposure (Thomsen & Wernberg 2005) on this system would be valuable. Hybrids between *C. maschalocarpum* and *C. flexuosum*, and between *C. maschalocarpum* and *C. plumosum*, have been reported (Dromgoole 1973, Lindauer *et al.* 1961) but seem more localised or morphologically cryptic than the widespread hybridisation found between *C. maschalocarpum* and *C. angustifolium*.

In *Cystophora* most species are hermaphrodite, and hybrids should be rare. Further investigation of the sister species *C. retorta* (hermaphrodite) and *C.*

siliquosa (dioecious) might show similar patterns of hybridisation and introgression to those found in *Fucus*. *Cystophora subfarcinata* and *C. retroflexa* are morphologically similar, often sympatric and both species are morphologically plastic. This might hide hybridisation between these species and further investigation might also reveal past or current hybridisation. Two specimens resembling *C. scalaris* were found north of the usual range of this species, one from Wilson Bay, Coromandel, the other from Ahuriri Street, Napier. Mitochondrial DNA sequences from these specimens grouped with *C. torulosa* and *C. retroflexa* (data not shown). Further data might confirm that these are *C. torulosa* × *C. retroflexa* hybrids.

Generally, the high frequency of natural hybridisation and absence of reproductive isolation in recognised species suggests reproductive isolation occurs at a late stage of speciation in brown algae, and is often preceded by ecological and morphological differentiation.

6.8 Molecular data

Early molecular studies focussed on characterising classes among the heterokontophyta (Bhattacharya *et al.* 1992, Daugbjerg & Andersen 1997a, b), or resolving ordinal or familial level phylogenetic relationships within the Phaeophyceae (Tan & Druehl 1993, 1994, Saunders & Druehl 1992, 1993, Rousseau *et al.* 1997, Peters & Clayton 1998, Siemer *et al.* 1998, Boo *et al.* 1999, Rousseau & de Reviers 1999a, 1999b, Serrão *et al.* 1999, Kawai & Sasaki 2000, Rousseau *et al.* 2000). Intra-familial relationships were investigated in *Ectocarpus* and *Kukuckia* (Stache-Crain *et al.* 1997), and *Laminariocolax* (Burhardt & Peters 1998), *Pelvetia* and *Pelvetiopsis* (Lee *et al.* 1998, 1999), and *Undaria* (Yoon & Boo 1999). This early work was reviewed by Druehl & Saunders (1992) and de Reviers & Rousseau (1999). It resulted in substantial reorganisation of the orders and families within the Phaeophyceae.

Phylogenetic resolution of the brown algal lineage has increased dramatically over the last decade. Taxon representation has increased considerably (Kawai *et al.* 2001, Rousseau *et al.* 2001, Draisma *et al.* 2002, Lee & Bae 2002, Cho *et al.* 2006, De Clerck *et al.* 2006), as have the number of molecular markers used (Kim *et al.* 2002, 2003, Cho *et al.* 2004, Lane *et al.* 2006,

Phillips *et al.* 2008, Kogishi *et al.* 2010). Recent studies have combined markers from nuclear, mitochondrial and chloroplast genomes (Bittner *et al.* 2008, Fraser *et al.* 2010b, Tronholm *et al.* 2010b, Silberfield *et al.* 2010).

A second focus of molecular studies has been resolving the systematic position of “difficult” or anomalous species, including *Notheia anomala* (Saunders & Kraft 1995), *Myagropsis myagroides* (Horiguchi & Yoshida 1998), *Halosiphon tomentosus* (Peters 1998), *Laminarionema elsbetiae* (Peters & Burkhardt 1998), *Desmarestia chordalis* (Peters *et al.* 2000), *Caepidium antarcticum* (Peters & Ramírez 2001), *Microzonina velutina* (Burrowes *et al.* 2003), *Stschapovia flagellaris* (Kawai & Sasaki 2004), *Phaeostrophion irregulare* (Kawai *et al.* 2005), *Petrospongium rugosum* (Cho & Boo 2006), *Discosporangium mesartrocarpum* (Kawai *et al.* 2007), and *Herpodiscus durvillaeae* (Heesch *et al.* 2008).

While not aimed at species differentiation, this phylogenetic work did offer relevant insights. In the Scytosiphonaceae, Kogame *et al.* (1999) found molecular differentiation was better correlated with the morphology of the sporophyte microthallus than with the morphology of the upright stage, which appears to be more morphologically plastic. Low variation in markers was shown in ITS sequences from *Desmarestia* (van Oppen *et al.* 1993), *Fucus* (Leclerc *et al.* 1998) and *Sargassum* (Stiger *et al.* 2000) and in mitochondrial markers (Coyer *et al.* 2006a); this suggests these markers are conserved or speciation is recent. The notion that morphological complexity is only associated with the most derived groups of brown algae was finally dispensed with (Rousseau & de Reviers 1999a, Draisma *et al.* 2001, Phillips *et al.* 2008).

Numerous studies have proposed new species supported by phylogenetic data, including *Asterocladon lobatum* (Müller *et al.* 1998), *Sargassum boreale* (Yoshida *et al.* 2000), *Chorda rigida* (Kawai *et al.* 2001), *Neoleptonema yongpilii* (Lee *et al.* 2002), *Newhousia imbricata* (Kraft *et al.* 2004), *Asterocladon interjectum* (Uwai *et al.* 2005), *Cladosiphon umezakii* (Ajisaka *et al.* 2007), *Aureophycus aleuticus* (Kawai *et al.* 2008), *Fucus radicans* (Bergström *et al.* 2005), *Dictyota cyanoloma* and *D. cymatophila* (Tronholm *et al.* 2010a, 2010b), and four *Padina* species (Ni-Ni-Win *et al.* 2010). Particularly revealing of the power of molecular techniques is the reinstatement of *Ectocarpus crouaniorum*, a third European species in this well-studied genus (Peters *et al.* 2010). In most

cases, phylogenetic data was used in support of morphological diagnoses, and to place new species in higher taxonomic groups. Recently, Tronholm *et al.* (2010a) separated the “pseudo-cryptic” *Dictyota cymatophila* from *D. dichotoma*, primarily by phylogenetic data. Phenological and morphometric data also separated species, but showed some overlap, and it is unlikely two species would have been diagnosed without phylogenetic data.

In the last decade an increasing number of studies have addressed the phylogeography of single species or closely related species over all or most of their range (Coyer *et al.* 2003, Cho *et al.* 2005, Harvey & Goff 2006, Hoarau *et al.* 2007, Lane *et al.* 2007, Fraser *et al.* 2009a, 2009b, 2009c, Pereyra *et al.* 2009, Tellier *et al.* 2009, Uwai *et al.* 2009, Coyer *et al.* 2010, Peters *et al.* 2010, Fraser *et al.* 2010, Kogishi *et al.* 2010, Neiva *et al.* 2010, Olsen *et al.* 2010, Cheang *et al.* 2010a, 2010b). These studies have sufficient detail to explore the history of species themselves (rather than a broad history of relationships between species). In many cases (e.g., Tellier *et al.* 2009, Coyer *et al.* 2010), multiple lineages have been found within current species, and some authors have proposed splitting species (Fraser *et al.* 2010b, Martin & Zuccarello in preparation).

These data should allow species delimitation to proceed from a morphological concept of species to a phylogenetic concept. In general this hasn't happened, as there has been a separation of phylogenetic analyses and taxonomic treatment of species. For example, Stache-Crain *et al.* (1997), in an early phylogenetic study that included worldwide sampling, found *Ectocarpus siliculosus* formed several clades which they suggested might have species status, but declined to create new species due to lack of morphological differentiation. Cho *et al.* (2005) found two species of *Colpomenia peregrina*, separated by phylogeographic and ecological data, and Cho *et al.* (2007) found phylogenetic evidence for two species of *Scytosiphon lomentaria*, but neither study established new species. Only a few authors have explicitly addressed phylogenetic species concepts. Harvey & Goff (2006) rejected the morphological species concept as misleading in the case of *Halidrys* and *Cystoseira* species from the Pacific Coast of North America, showing Pacific species had a common origin, separate from Atlantic members of these genera. Pacific species had independently converged to morphologies similar to Atlantic species. These morphologies were diverse, despite low genetic variation between Pacific species. The authors concluded that

revision using a phylogenetic species concept was needed. Fraser *et al.* (2009a) applied a phylogenetic species concept to *Durvillaea antarctica* and suggested that a morphologically variant form (the ‘cape’ morphology, where the thallus forms a broad sheet, rather than dividing into thongs) can be separated by morphological and molecular data and should be regarded as a separate species. This highlights the problems with incipient species. The ‘cape’ form forms a monophyletic group in *cox1* trees, but this renders *D. antarctica* polyphyletic, as the ‘thong form’ Chilean lineage is sister to the “cape” clade, while the New Zealand and southern ocean lineage is derived (Fraser *et al.* 2009b). There is no evidence for morphological separation of the Chilean and New Zealand clades, but a pure phylogenetic species concept would support this (Fraser *et al.* 2010b). Either several lineages must be separated solely on the basis of molecular data or *D. antarctica* should be regarded as a meta-species (*sensu* de Queiroz & Donoghue 1988), encompassing several distinct mitochondrial lineages but without ascribing species status to these lineages. Conversely, *D. chathamensis* is morphologically distinct from the allopatric *D. antarctica* lineage, but is not differentiated by most markers (only by *rbcL*, Fraser *et al.* 2010b). In these examples, the phylogenetic species concept has been invoked to support *suggestions* for species delimitation, and in support of other data, but no author has split species of brown algae under a purely phylogenetic species concept, although this has been done with green algae (Leliaert *et al.* 2009) and microalgae (Lilly *et al.* 2007, Coffroth & Santos 2005). Instead, phycologists have delimited species by considering the weight of available evidence. Recently, phylogenetic data has been given greater weight, with the synonymisation of some morphologically variable species (Macaya & Zuccarello 2010).

In part, the reluctance to apply a pure phylogenetic species concept relates to the difficulty in assigning hierarchical levels to phylogenetic lineages and the increasing power of molecular data to diagnose lineage differentiation (Avice & Wollenberg 2007). Phylogenies are limited in being based on a small sample of genetic information from a small sample of specimens, and generally produce coarse-scale lineages showing simplified relationships between species. Complex histories, with lineages within species lineages, as well as reticulation, raise a major question for taxonomic work: to what degree are speciation processes active within most lineages? At one extreme, lineages might undergo brief periods

of speciation, separated by long periods of stasis (with cohesive, well differentiated species), at the other extreme, lineages might be continually subdividing, with morphological differentiation, reproductive isolation and other properties exhibiting themselves at different levels throughout a lineage's history.

Recent studies in North Atlantic *Fucus* species demonstrate the difficulty of applying methods of species delimitation that assume simple speciation histories and test properties of species that are confounded by hybridisation and intraspecific morphological variation (Coyer *et al.* 2006a). Over 125 intraspecific taxa have been described in *Fucus* (Coyer *et al.* 2010), based on morphology. Various numbers of species can be recognised, depending on the degree of lineage separation and which property of a species-lineage is tested for (De Queiroz 2007). Phylogenetic approaches using ITS (Serrão *et al.* 1999) and mitochondrial markers (Coyer *et al.* 2006a) discerned two major clades, one containing *F. vesiculosus*, *F. spiralis*, *F. radicans*, *F. ceranoides* and *F. virsoides*, but were unable to separate these species. Cluster analyses using microsatellites separated *F. vesiculosus* from *F. spiralis* (Engel *et al.* 2005) and also separated *F. ceranoides* (Billard *et al.* 2005). Bergström *et al.* (2005) delimited a third species *Fucus radicans*, which is estimated to have diverged from *F. vesiculosus* as little as 400 years ago (Pereyra *et al.* 2009), driven by adaption to a low salinity habitat (the Baltic Sea), and this delimitation is supported by functional divergence in heat shock response (Lago-Lestón *et al.* 2010). Billard *et al.* (2010) then divided *F. spiralis* into two clusters, spatially segregated into a high shore and low shore varieties. Despite ecological, molecular and morphological differences, they did not interpret these as species, partly because there is less differentiation between the forms than between *F. spiralis* and *F. vesiculosus*. Most recently, Coyer *et al.* (2010) found that the southern part of the *F. spiralis* range forms a third cluster. This southern cluster hybridised with *F. vesiculosus* at some point in the past, producing the low shore cluster. Also, while a nuclear marker shows reciprocal monophyly between *F. ceranoides* and *F. vesiculosus*, the entire northern range of *F. ceranoides* is introgressed by mitochondrial DNA from *F. vesiculosus* (Neiva *et al.* 2010). Coyer *et al.* (2010) propose a “sliding window” concept of speciation for *Fucus*, where different species are delineated within a continuum of species separation.

These studies, and in-depth sampling of other species (Lane *et al.* 2007, Olsen *et al.* 2010) show an increasing ability to infer more detailed histories of species, and make the application of an informed historical species concept plausible in brown algae.

6.9 Species delimitation in *Carpophyllum*

Four species are currently recognised in *Carpophyllum* (Adams 1994, Lindauer *et al.* 1961), and my molecular data support continued recognition of these species. All species are morphologically and ecologically distinct, but genetic distances between markers are low. *C. maschalocarpum* and *C. angustifolium* can be separated by mitochondrial spacer data (Chapter 2) but with some reservations: (1) there is evidence for mitochondrial introgression in some populations (Chapter 3); (2) genetic distances within species are greater than distance between species; and (3) one cluster, associated with specimens from North Cape and Wekarua, is not clearly associated with either species (Chapter 2). Some genetic data was collected on *Carpophyllum plumosum* and *C. flexuosum* but was not included in the previous chapters. These data support the continued recognition of these species. For example I found unique ITS2 sequences for *C. plumosum* and *C. flexuosum* (but only one mutational step removed from each other and from *C. maschalocarpum*/*C. angustifolium* sequences), and mitochondrial spacer sequences that are genetically distant from other species.

The phylogenetic relationship between species is not completely clear. Markers such as ITS and *rbcLS* spacer show low variation (*rbcLS* spacer sequences are identical in *C. maschalocarpum* and *C. angustifolium*, with three changes separating these from *C. flexuosum* and six changes separating these from *C. plumosum*). *Carpophyllum angustifolium* and *C. maschalocarpum* are sister species according to all markers tested (the above and *rbcL*, *cox3*, *rps14-atp8*), and *C. plumosum* appears to be the earliest branching species. The history of *Carpophyllum* species is only partly elucidated by these data. Significant changes in the distribution of *C. maschalocarpum* appear to have taken place since the LGM, especially the colonisation of the southern North Island and South Island. The centre of genetic diversity of *C. angustifolium* appears to be Northland, but data are limited, whereas *C. maschalocarpum* is most genetically diverse in the

Bay of Plenty. Possibly these lineages separated in allopatry during periods of high sea level during the Pliocene (when the northern North Island was separated into islands), and are now hybridising following secondary contact. An alternative explanation is sympatric speciation from a common ancestor driven by strong selective pressure for different habitats, with *C. angustifolium* occupying exposed areas, *C. flexuosum* sheltered areas and *C. maschalocarpum* occupying intermediate environments. Further molecular data is needed to complete this picture, especially from *C. angustifolium*, as is ecophysiological data.

6.10 Species delimitation in *Cystophora*

Cystophora species have been delimited on the basis of morphological data. In New Zealand species I consider that some species have been over-split and I synonymise two pairs of names. I also suspect the Australian species *C. cuspidata* should be synonymised with *C. subfarcinata*, but sampling is inadequate to establish this. I was unable to find *C. cymodocea*, despite an extensive search of the type locality (Duttons Beach, Portland, Victoria) and suspect this is a distended form of *C. retroflexa* growing epiphytically. In all these cases, oversplitting, or potential oversplitting, appears to have arisen from limited characterisation of the range of morphological expression in species. Further molecular work can address some of these questions, and more work is needed, especially to clarify species boundaries in Australian species, and to further understand the history of the evolution and dispersal of *Cystophora* species.

6.11 Dispersal and history

Dispersal ability is expected to correlate strongly with gene flow, and be a major factor in algal speciation. Species with limited dispersal potential, with immotile gametes and/or negatively-buoyant thalli, should show limited gene flow between populations, and stronger signatures of genetic structure and founder effects (van den Hoek 1987). Genetic structure is strong in many species of brown algae (Chapter 2, Williams & DiFiori 1996, Kusmo & Druehl 2000, Cho *et al.* 2005, 2007, Hoarau *et al.* 2007, Fraser *et al.* 2009b, Cheang *et al.* 2010), but the relationship between genetic structure and dispersal is complex. *Carpophyllum*

maschalocarpum has positively buoyant thalli and is frequently found in drift. Some well separated populations – such as mainland New Zealand and the Chatham Islands, share mitochondrial haplotypes, whereas other relatively close populations are apparently isolated (Chapter 2). In *Cystophora torulosa*, *C. platylobium* and *C. retroflexa*, *cox1* haplotypes and ITS ribotypes are shared between New Zealand and Australia, as were haplotypes of a rapidly evolving mitochondrial spacer in *C. retroflexa*, suggesting multiple trans-Tasman dispersal (Chapter 4). In *Macrocystis pyrifera*—a species that has been shown to remain fertile after extended periods afloat (Macaya *et al.* 2005, Hernández-Carmona *et al.* 2006)—Macaya & Zuccarello (2010) found one *cox1* haplotype that is distributed throughout the southern hemisphere, but several other *cox1* haplotypes were sampled in single populations only. *Durvillaea* is especially interesting, having large buoyant thalli that appear capable of long-distance dispersal through the southern oceans resulting in genetically homogenous populations (Fraser *et al.* 2009b), but has limited north-south dispersal resulting in well differentiated lineages in lower latitudes. Also curious is that one lineage of *Durvillaea* (the “cape form”) is found from New Zealand’s sub-Antarctic islands and along the eastern coasts of New Zealand’s South Island, while other lineages are latitudinally restricted. The cape form has no obvious differences in dispersal potential. These buoyant species contrast with *Lessonia variegata* and *L. nigrescens*, species with heavy, negatively buoyant thalli, which show strong genetic differentiation between regions of New Zealand (Martin & Zuccarello in review) and Chile (Tellier *et al.* 2009), respectively.

These studies suggest brown algae with buoyant thalli have increased potential for long distance transport, but also that this potential is not always realised. Partly this results from geography. Dispersal is rapid along pathways such as the southern ocean west wind drift system (Waters 2008), and perhaps along the East Auckland Current (Chapter 2, although data from other species is needed to confirm this), but might be restricted elsewhere (Fraser *et al.* 2009b, 2010b, Chapter 2). But biological interactions such as density blocking (suggested by Fraser *et al.* 2009b, 2010b) might be important. Several areas remain to be addressed. One is the pathways taken by drifting algae. Data from *Carpophyllum* suggest relatively high gene flow between the Chatham Islands and the North Island. Data from drift devices show current transport between the mainland and

the Chatham Islands requires 30 to 50 days (Chiswell 2009), but windblown floating material might travel considerably faster, especially during extreme weather events, for example, material from a boat wrecked in Cook Strait has been recorded arriving at the Chatham Islands three weeks later (Young 1929). Direct measurements of seaweed transport rates (Harrold & Lisina 1989, Hawes 2008) should elucidate drift rates.

Several lines of evidence suggest *Cystophora* evolved in Australia and species dispersed to New Zealand relatively recently: (1) Species diversity in *Cystophora* is greatest in southern Australia, and the earliest diverging species (*C. moniliformis*) is endemic to southern Australia; (2) Several species are endemic to Australia, whereas only one species (*C. scalaris*) is endemic to New Zealand (Womersley 1964); and (3) two mitochondrial spacer haplotypes found in New Zealand specimens are shared with Australian specimens (Chapter 4). Sharing of mitochondrial spacer haplotypes suggests very recent dispersal, given the strong structuring shown in this marker in *Carpophyllum*. *Cystophora* is thus part of what Parsons (1985) termed the Australasian element of the New Zealand seaweed flora. All New Zealand species are vesiculate, whereas several Australian species (e.g., *C. moniliformis*, *C. siliquosa*, *C. xiphocarpa*) lack vesicles, or rarely develop vesicles (Womersley 1964). It seems likely that vesicles are associated with long distance dispersal, although several vesiculate Australian species are absent from New Zealand, despite tolerating similar temperature regimes (Adey & Steneck 2001). Whether successful trans-Tasman dispersal is a matter of chance or if there are other ecophysiological factors involved in dispersal or establishment is not known.

6.12 Genetic distance – molecular data without history

Genetic distance is a straightforward method of delineating species that uses phenetic, rather than historical criteria. Various methods of delineating species using genetic distance have been proposed (Sites & Marshall 2004), usually requiring population level data. These include comparing geographical distance with genetic distance (Good & Wake 1992), with significant deviations from isolation-by-distance suggesting more than one species. Highton (1989) suggests testing for a significant bimodal distribution of Nei's D parameter for population

differentiation, with a threshold suggested above which separate peaks represent multiple species. These methods have not been applied to brown algae. Both methods are essentially ahistorical, and might be confounded by strong structuring of populations, for example, the mtDNA spacer dataset for *Carpophyllum maschalocarpum* shows a bimodal distribution of Nei's D, this is a consequence of demographic expansion of the southern lineage, but there is no supporting evidence that this represents a speciation process. Similarly, the existence of particular dispersal pathways in *C. maschalocarpum* confounds the assumption of a linear relationship between geographical and genetic distance.

Using genetic distance might overcome the problem created by modern molecular data's power to diagnose extremely fine lineages, which could result in oversplitting of species under the phylogenetic species concept (Avice & Wollenberg 1997). But, there are problems that genetic distance might not be correlated with reproductive isolation (Ferguson 2002, Gourbière & Mallet 2009, Presgraves 2010). Speciation times appear very variable. Knowlton *et al.* (1993) suggested that many marine organisms had developed reproductive barriers since the closing of the Central American isthmus, c. 3 million years ago. Conversely, *Fucus radicans* might have attained some reproductive isolation in 400 years (Pereyra *et al.* 2009). In many cases, especially in higher latitudes, substantial demographic and geophysical changes (temperature, sea level changes) have occurred in the last 3 million years (Hewitt 2004, Maggs *et al.* 2008), creating complex patterns of isolation and contact (Coyer *et al.* 2010) that confound inferences based on genetic distance.

A recent development in distance methods is the emergence of DNA barcoding. Strictly, barcoding is a method of specimen identification, but the leading proponent of DNA barcoding, Paul Hebert, has described DNA barcoding as a method of species delimitation (Hebert & Gregory 2005) and assessing the barcoding method requires assessing the ability of mitochondrial data to delimit species.

Molecular tools have long been used for species identification in brown algae. Early studies used allozymes (Benzie *et al.* 2000, Hull *et al.* 2001), RAPDs (Ho *et al.* 1995) or sequence data (Kogame & Masuda 2001). Molecular identification continues to be a useful tool, especially with cryptic forms (Fox &

Swanson 2007), morphologically plastic (Kucera & Saunders 2008, Endo *et al.* 2009) or convergent species (Camus *et al.* 2005, Jones *et al.* 2010).

DNA barcoding distinguishes species using genetic distance between phenetic clusters inferred from short (c. 600 bp) mitochondrial DNA sequences. DNA barcoding has a compelling simplicity: a threshold between intraspecific and interspecific genetic variation is defined (usually ten-times, Golding *et al.* 2009) and specimens exceeding this threshold are considered to be separate species. Species can then be identified by simple sequence comparisons against a universal database.

Barcoding requires that genetic variation in the selected gene region can be adequately characterised. Intraspecific variation requires extensive sampling to understand the genetic structure – and therefore the species history. For example, extensive global sampling of *Durvillaea antarctica* by Fraser *et al.* (2010b) showed extensive *cox1* variation that was highly geographically structured, with sufficient lineage delineation to propose a species complex. One *cox1* haplotype was extensively distributed – around most of the southern oceans. In this situation, even widespread sampling, by the standards of barcoding studies, might return a single haplotype, even if samples were drawn from eight different countries. Such a result might suggest identification of *D. antarctica* by *cox1* barcoding was straightforward, but this sampling would not enable accurate identification of further specimens from northern localities.

Similarly in Chapter 2, one mitochondrial spacer haplotype is present over a wide part of the range of *Carpophyllum maschalocarpum*. This marker that is probably more variable than regions used for barcoding but it illustrates the point. A collection of 10 specimens might detect only a single haplotype. With slightly wider sampling one might get two haplotypes, only two steps removed from each other and conclude that intraspecific variation is low. With more sampling one might find a third haplotype, four steps removed from the others, or even further removed if you sample one of these haplotypes which appear to have entered the *C. maschalocarpum* mitochondrial genome following introgressive hybridisation with *C. angustifolium*, and assume cryptic speciation was taking place. Accurate characterisation of genetic variation requires sampling from the Bay of Plenty, where most of the variation is found.

Proponents of barcoding appeal to the speed and low cost of assembling sets of “barcodes” that can then be used as vouchers for subsequent species identification. According to proponents, barcoding will facilitate documenting as many species as possible before they go extinct (Baker *et al.* 2009), allow fast organisation of molecular data into putative species groups before formal description (Litaker *et al.* 2007, Borisenko *et al.* 2009), eliminate the need for the complex taxonomic training that is currently required for species description and identification (Sass *et al.* 2007), allow identification of incomplete specimens, (Schindel & Miller 2005), and provide a tool for non-specialists, for example in water quality assessments (Moniz & Kaczmarek 2009), identifying invasive species (Saunders 2009), correct identification of difficult and cryptic species (Starr *et al.* 2009), greater discrimination of cryptic species (Sherwood *et al.* 2008) and standardisation of molecular data (McDevit & Saunders 2010). The appeal of many of these rationales is low cost, rather than scientific rigour. This might be reasonable in some applications where a number of misidentifications are acceptable, especially where error rates can be estimated and reported, but is of questionable scientific value.

Only a few studies have attempted to validate the use of *cox1* for barcoding brown algae. Lane *et al.* (2007) attempted a *cox1* barcoding study of *Alaria* species from the north-east Pacific. Genetic variation was relatively low and there was no clear threshold between inter- and intra-clade variation. In addition, clusters of *cox1* mitotypes were not congruent with *a priori* morphological species assignments, or with ITS clades. ITS variation was relatively high, and did not resolve species level groups. This result was interpreted as showing widespread introgression following secondary contact by species that evolved in allopatry, with introgression and ITS recombination.

Kucera & Saunders (2008) used *cox1* to delineate North American samples of *Fucus*. Mitotypes separated into three clades. *Fucus serratus* from Canada were well demarcated from other *Fucus* species but this species is a recent (1887) introduction to North America and this deme would be expected to show low variation. *Fucus spiralis* and *F. vesiculosus* were not separated, which is not surprising given the low variation and extensive hybridisation between these species. *Fucus distichus* and north Pacific specimens identified in the fields as *F. spiralis* were also not separated, but the authors consider *F. spiralis* in this clade

to result from incorrect morphological assignment of specimens of *F. distichus* that has undergone morphological convergence to an *F. spiralis* form. The authors then made an *a posteriori* reassignment of these specimens to *F. distichus*, with hybridisation ruled out due to the absence of polymorphic sites in ITS sequences. This was not accompanied by any anatomical evidence, and it seems possible that these are *F. spiralis* with introgressed *F. distichus* mitochondria and ITS which has homogenised to a *F. distichus*-ribotype following an earlier hybridisation event. Widespread mitochondrial introgression has been reported in European *F. serratus* (Coyer *et al.* 2010) and *F. ceranoides* (Neiva *et al.* 2010). This study was reported as demonstrating the efficacy of *cox1* barcoding in brown algae, but it is unclear what hypothesis was being tested here, as genetic distance between clusters was low or absent, morphology and molecular delimitation was not congruent, and sampling was limited to North American demes of *Fucus* species which have their centre of diversity in Europe (Coyer *et al.* 2003, Hoarau *et al.* 2007, Coyer *et al.* 2006a, 2010).

Other studies of brown algae have shown broad variation in *cox1* sequences across orders (McDevit & Saunders 2009), but did not address species delimitation, which requires inclusion of sister species, and addressed biogeographical structure of populations with extensive sampling (McDevit & Saunders 2010). In general these studies have shown mean genetic distances between *cox1* sequences of sister species of brown algae are often less than the 10X threshold proposed for barcoding animals. Presumably a high threshold is to compensate for low sampling and distance methods' limitations in capturing evolutionary processes. In *Cystophora* intraspecific variation was of a similar order to interspecific variation between some sister species. Recognition of these species is supported by morphology and ITS sequences, and further sampling is likely to increase intraspecific variation. An alternative to genetic distances is character based barcoding methods, where a designated number of nucleotide positions or combination of positions is used to characterise species (DeSalle 2006, Rach *et al.* 2008). This requires extensive sampling to ensure proposed characters are fixed (Wiens & Servedio 2010).

Cytoplasmic introgression is an obvious problem for mitochondrial sequence based barcoding, and has been found in *Fucus* (Neiva *et al.* 2010, Coyer *et al.* 2010), *Alaria* (Lane *et al.* 2007) and *Carpophyllum angustifolium* (Chapter

4). Comparison of cytoplasmic markers with single copy nuclear markers (Neiva *et al.* 2010) should show introgression, but so far these markers have not been widely available for brown algae. Future work should show whether these are unusual cases, or whether cytoplasmic introgression is widespread in the brown algae.

Two developments suggest future barcoding studies will be better integrated with other approaches. One is multi-locus barcoding (Kress *et al.* 2009) which allows barcoding data to be analysed using phylogenetic methods. Kress *et al.* (2009) also focussed on a single locality (Barro Colorado Island, Panama), where species diversity is well characterised and sampling intra-species genetic diversity should be relatively straightforward. Other barcoding studies have employed additional markers to overcome uncertainty (Lane *et al.* 2007, McDevit & Saunders 2010). A second development is more widespread sampling, allowing barcoding data to be used for mitochondrial phylogeographic studies (McDevit & Saunders 2010). These trends might overcome the limitations of low sampling and single markers proposed in the original barcoding concept (Hebert *et al.* 2003a, 2003b).

6.13 Model based methods

Microsatellites analysis has been used to separate closely related brown algal species. Most studies use a Bayesian procedure (STRUCTURE) that simultaneously generates clusters and assigns specimens to clusters based on a model that assumes Hardy-Weinberg equilibrium and linkage equilibrium within populations (Pritchard *et al.* 2000). This approach has been used to separate species of *Fucus* (Coyer *et al.* 2003, Billard *et al.* 2005, Engel *et al.* 2005, Tatarenkov *et al.* 2007) where it has successfully discriminated species where there is low phylogenetic divergence and gene flow between species. This is useful but does not elucidate species histories, as there are difficulties in using microsatellites for phylogenetic inference (Noor *et al.* 2001, Buschiazzi & Gemmell 2006).

Knowles & Carstens (2007) suggested using coalescence to estimate the probabilities of gene trees under a particular history to evaluate the possibility of lineage splitting (i.e. speciation). This models species histories probabilistically

rather than inferring species history from indications of reciprocal monophyly in sampled genes. A method that combines population biology and phylogenetic approaches was developed by Pons *et al.* (2006). This method attempts to determine a threshold between inter-species level (phylogenetic) lineage branching and intraspecies (tokogenetic) branching, using a “Generalised Yule Mixed Coalescent model.” A maximum likelihood tree is inferred from sequence data, the ages of nodes are estimated, and a lineage-through-time plot constructed. A change in branching rates in this plot should correspond to a change from phylogenetic processes to coalescent processes. This method has not been used in brown algae. Leliaert *et al.* (2009) applied the method to green algae (*Boodlea*), but found estimated confidence intervals were very broad, allowing a wide range of species divisions. A preliminary attempt at applying this method to *cox1* data from *Cystophora*, with changes in branching rates estimated visually from the lineage-through-time plot, was not congruent with expected species boundaries, and different species boundaries were indicated with small changes in parameters.

6.14 Synthesising species concepts

Eukaryote algae have not featured greatly in the development of species concepts. Birds seem somewhat overrepresented, probably because of the influence of Ernst Mayr; terrestrial plants usually get a mention (hybridisation), as do prokaryotes (asexuality). None-the-less, debates over species concepts have influenced phycologists and are pertinent to any discussion of species delimitation.

At the end of the 1990s various attempts were made to synthesise species concepts (Avise & Wollenberg 1997, Mayden 1997, de Queiroz 1998). These attempts were anticipated by Hull (1965) who proposed understanding species as a cluster concept. Avise & Wollenberg (1997) suggested that there is no difficulty in combining the biological (BSC) and phylogenetic species concepts (PSC), as long as diagnosis of monophyly uses multiple lines of evidence. This “multi-locus PSC” would minimise conflicts between gene trees and species trees, and limit diagnosis of sub-species lineages. Mayden (1997) argues for a hierarchy of species concepts, with the highest order species concept not operable, but based on theory, but supported by various lower order concepts that are, to varying degrees, operable. This allows a unification of species concepts, with the

relationships between them clearly defined. De Queiroz (1998, 2007) suggests separating the definition of species from operational methods used to delimit species, as these are different philosophical concepts, albeit ones that can inform one another. In De Queiroz's scheme (2007), species are metapopulation lineages that acquire different properties (monophyly, reproductive isolation, etc.) at different times during the process of speciation. Species are delimited by testing for these various properties, but not all properties will coincide at one moment in the history of speciation, so no single property is required (or expected) in every species. Successful delimitation of species, therefore, will depend on the property tested and the history of lineage separation.

Piglucci (2003) develops a similar argument to De Queiroz, but one that is not based on history. They argue that the species problem is essentially a philosophical question that needs empirical data to answer it (rather than a scientific question with philosophical aspects). The question can be resolved by viewing species as a cluster concept *sensu* Wittgenstein, where there are shared properties between all species, none of which are necessary or sufficient alone.

These concepts are compatible with Coyer *et al.*'s (2010) sliding window concept of species for *Fucus*. They also suggest a weight of evidence approach to species delimitation, rather than a Popperian hypothesis testing approach. This is similar the actual practice by taxonomists dealing with brown algae. With the availability of various lines of evidence, it is becoming rare for a single property of a species to be privileged above all others. Rather, various properties are considered and weighed before a decision is made.

Uncertainty is likely to remain a problem for species delimitation. Contemporary data will rarely – if ever – support a unique hypothesis about historical events. The complexity of biological processes, the use of proxies (morphology, genetic distance) for speciation, inference from limited data and researchers' biases (splitters and lumpers) suggests species delimitation will remain an exercise in approximation, subject to revision.

I suggest no method is likely to encompass the diversity of evolutionary processes and historical scenarios found in speciation, and conclude that: (1) a weight of evidence approach should be taken in delimiting species, with no single method being privileged (this essentially continues the *status quo* in brown algal taxonomy) and; (2) different sets of species' properties will be found in different

species. These properties are best encapsulated by including species' history in species descriptions.

Contingency plays a large role in speciation processes, and the delimitation of species is an exploration of biological history. There are expected to be major differences in the speciation histories of different lineages, for example the fast speciation in *Fucus radicans* where reinforcement (selection for characters limiting hybridisation) might be expected to play a role, or the slow drift to reproductive isolation in allopatric species. Incorporating species histories into the description of species would overcome the lingering influence of the typological approach, where species are described by a name and morphological diagnosis. Explicitly including history as part of species descriptions would emphasise that recognition that a lineage has species status can reflect different speciation processes.

6.15 References

- ABBOTT, I. A. & HOLLENBERG, G. J. (1976). *Marine algae of California*. Stanford CA: Stanford University Press.
- ADAMS, N. M. (1994). *Seaweeds of New Zealand: An illustrated guide*. Christchurch, New Zealand: Canterbury University Press.
- ADEY, W. H. & STENECK, R. S. (2001). Thermogeography over time creates biogeographic regions: A temperature/space/time-integrated model and an abundance weighted test for benthic marine algae. *Journal of Phycology*, 37, 677–698.
- AGARDH, J. G. (1848). *Species genera et ordines algarum, seu descriptiones succinctae specierum, generum et ordinum, quibus algarum regnum constituitur. Volumen Primum. Algas fucoideas complectens*. Lundae [Lund], Sweden: C. W. K. Gleerup.
- AJISAKA, T., KIM, S. H., UWAI, S. & KAWAI, H. (2007). *Cladosiphon umezakii* sp. nov. (Ectocarpales, Phaeophyceae) from Japan. *Phycological Research*, 55, 193–202.
- ARNOLD, M. L. & HODGES, S. A. (1995). Are natural hybrids fit or unfit relative to their parents? *Trends in Ecology and Evolution*, 10, 67–71.
- AVISE, J. C. & WOLLENBERG, K. (1997). Phylogenetics and the origin of species. *Proceedings of the National Academy of Sciences of the USA*, 94, 7748–7755.
- BAKER, A. J., TAVERES, E. S. & ELBOURNE, R. F. (2009). Countering criticisms of single mitochondrial DNA gene barcoding in birds. *Molecular Ecology Resources*, 9, (Suppl. 1) 257–268.
- BENZIE, J. A. H., BALLMENT, E. & EDYVANE, K. (2000). Allozymes as genetic identification markers of *Sargassum* spp. (Phaeophyta) from the Great Barrier Reef, Australia. *Botanica Marina* 43, 169–179.
- BERGSTRÖM, L., TATARENKOV, A., JOHANNESSON, K. JÖNSSON, R. B. & KAUTSKY, L. (2005). Genetic and morphological identification of *Fucus radicans* sp. nov. (Fucales, Phaeophyceae) in the brackish Baltic Sea. *Journal of Phycology*, 41, 1025–1038.

- BHATTACHARYA, D., MEDLIN, L., WAINRIGHT, P. O., ARIZTIA, C. B., STICKEL, S. K. & SOGIN, M. L. (1992). Algae containing chlorophylls a + c are paraphyletic: Molecular evolutionary analysis of the chromophyta. *Evolution*, **46**, 1801–1817.
- BILLARD, E., DAGUIN, C., PEARSON, G. SERRÃO, E., ENGEL, C. & VALERO, M. (2005). Genetic isolation between three closely related taxa: *Fucus vesiculosus*, *F. spiralis* and *F. ceranoides* (Phaeophyceae). *Journal of Phycology*, **41**, 900–905.
- BILLARD, E., SERRÃO, E., PEARSON, G. A., DESTOMBE, C. & VALERO, M. (2010). *Fucus vesiculosus* and *spiralis* species complex: a nested model of local adaptation at the shore level. *Marine Ecology Progress Series*, **405**, 163–174.
- BILLARD, E., SERRÃO, E., PEARSON, G. A., ENGEL, C. R., DESTOMBE, C. & VALERO, M. (2005). Analysis of sexual phenotype and prezygotic fertility in natural populations of *F. spiralis*, *F. vesiculosus* (Fucaceae, Phaeophyta) and their putative hybrids. *European Journal of Phycology*, **40**, 397–407.
- BITTNER, L., PAYRI, C. E., COULOUX, A., CRUAUD, C., DE REVIERS, B. & ROUSSEAU, F. (2008). Molecular phylogeny of the Dictyotales and their position within the Phaeophyceae, based on nuclear, plastid and mitochondrial sequence data. *Molecular Phylogenetics and Evolution*, **49**, 211–226.
- BOO, S. M., LEE, W. J., YOON, H. S., KATO, A. & KAWAI, H. (1999). Molecular Phylogeny of Laminariales (Phaeophyceae) inferred from small subunit ribosomal DNA sequences. *Phycological Research*, **47**, 109–114.
- BORISENKO, A. V., SONES, J. E. & HEBERT, P. D. N. (2009). The front-end logistics of DNA barcoding: challenges and prospects. *Molecular Ecology Resources*, **9**, (Suppl. 1), 27–34.
- BOTHWELL, J. H., MARIE, D., PETERS, A. F., COCK, J. M. & COELHO, S. M. (2010). Role of endoreduplication and apomeiosis during parthenogenetic reproduction in the model brown alga *Ectocarpus*. *New Phytologist*, **188**, 111–121.
- BROWER, A. V. Z. (1999). Delimitation of phylogenetic species with DNA methods: A critique of Davis and Nixon's Population Aggregation Analysis. *Systematic Biology*, **48**, 119–213.
- BURKHARDT, E. & PETERS, A. F. (1998). Molecular evidence from nrDNA ITS sequences that *Laminariocolax* (Phaeophyceae, Ectocarpaceae *sensu lato*) is a worldwide clade of closely related kelp endophytes. *Journal of Phycology*, **34**, 682–691.
- BURROWES, R., ROUSSEAU, F., MÜLLER, D. G., DE REVIERS, B. (2003). Taxonomic placement of *Microzonina* (Phaeophyceae) in the Syringodermatales based on the rbcL and 28S nrDNA sequences. *Cryptogamie Algologie*, **24**, 63–73.
- BURROWS, E.M. & LODGE, S. (1951). Autecology and the species problem in *Fucus*. *Journal of the Marine Biological Association of the United Kingdom*, **30**, 161–176.
- BUSCHIAZZO, E. & GEMMELL, N. J. (2006). The rise, fall and renaissance of microsatellites in eukaryotic genomes. *BioEssays*, **28**, 1040–1050.
- BUTLIN, R. K., GALINDO, J. & GRAHAME, J. W. (2008). Sympatric, parapatric or allopatric: the most important way to classify speciation? *Philosophical Transactions of the Royal Society of London B Series*, **363**, 2997–3007.
- CAMUS, C., MEYNARD, A. P., FAUGERON, S. KOGAME, K. & CORREA, J. A. (2005). Differential life history phase expressions in two co-existing species of *Scytosiphon* (Phaeophyceae) in northern Chile. *Journal of Phycology*, **41**, 931–941.
- CHEANG, C. C., CHU, K. H. & ANG, P. O. (2010). Phylogeography of the marine macroalga *Sargassum hemiphyllum* (Phaeophyceae, Heterokonta) in northwestern Pacific. *Molecular Ecology*, **19**, 2933–2948.
- CHISWELL, S. M. (2009). Colonisation and connectivity by intertidal limpets among New Zealand, Chatham and Sub-Antarctic Islands. II. Oceanographic connections. *Marine Ecology Progress Series*, **388**, 121–135.

- CHO, G. Y. & BOO, S. M. (2006). Phylogenetic position of *Petrospongium rugosum* (Ectocarpales, Phaeophyceae): insights from the protein coding plastid *rbcL* and *psaA* gene sequences. *Cryptogamie, Algologie*, 27, 3–15.
- CHO, G. Y., BOO, S. M., NELSON, W. & CLAYTON, M. N. (2005). Genealogical partitioning and phylogeography of *Colpomenia peregrina* (Scytosiphonaceae, Phaeophyceae) based on plastid *rbcL* and nuclear ribosomal DNA internal transcribed spacer sequences. *Phycologia*, 44, 103–111.
- CHO, G. A., KOGAME, K., KAWAI, H. & BOO, S. M. (2007). Genetic diversity of *Scytosiphon lomentaria* (Scytosiphonaceae, Phaeophyceae) from the Pacific and Europe based on RuBisCo large subunit and spacer, and ITS nrDNA sequences. *Phycologia*, 46, 657–665.
- CHO, G. A., LEE, S. H. & BOO, S. M. (2004). A new brown algal order, Ishigeales (Phaeophyceae), established on the basis of plastid protein-coding *rbcL*, *psaA* and *psbA* region comparisons. *Journal of Phycology*, 40, 921–936.
- CHO, G. Y., ROUSSEAU, F., REVIERS, B. DE & BOO, S. M. (2006). Phylogenetic relationships within the Fucales (Phaeophyceae) assessed by the photosystem I coding *psaA* sequences. *Phycologia*, 45, 512–519.
- CLAYTON, M. N. (1976). The morphology, anatomy and life history of a complanate form of *Scytosiphon lomentaria* (Scytosiphonales, Phaeophyta) from Southern Australia. *Marine Biology*, 38, 201–208.
- CLAYTON, M. N. (1978). Morphological variation and life history in cylindrical forms of *Scytosiphon lomentaria* (Scytosiphonaceae: Phaeophyta) from Southern Australia. *Marine Biology*, 47, 349–357.
- CLAYTON, M. N., KEVEKORDES, K., SCHOENWAEELDER, M. E. A., SCHMID, C. E. & ASHBURNER, C. M. (1998). Parthenogenesis in *Hormosira banksii* (Phaeophyceae). *Botanica Marina*, 40, 23–30.
- COFFROTH, J. A. & SANTOS, S. R. (2005). Genetic diversity of symbiotic dinoflagellates in the genus *Symbiodinium*. *Protist*, 156, 19–34.
- COYER, J. A., HOARAU, G., COSTA, J. F., HOGERDIJK, B., BILLARD, E., VALERO, M., ET AL. (2010). Evolution and species diversification within the intertidal brown macroalgae *Fucus spiralis*/*F. vesiculosus* species complex in the North Atlantic. *Molecular Phylogenetics and Evolution*, doi:10.1016/j.ympev.2010.11.015.
- COYER, J. A., HOARAU, G., OUDOT-LE SECQ, M-P., STAM, W. T. & OLSEN, J. L. (2006a). A mtDNA-based phylogeny of the brown algal genus *Fucus* (Heterokontophyta; Phaeophyta). *Molecular Phylogenetics and Evolution*, 39, 209–222.
- COYER, J. A., HOARAU, G., PEARSON, G. A., SERRAO, E. A., STAM, W. T. & OLSEN, J. L. (2006b). Convergent adaptation to a marginal habitat by homoploid hybrids and polyploid ecads in the seaweed genus *Fucus*. *Biology Letters*, 2, 405–408.
- COYER, J. A., HOARAU, G., SKAGE, M., STAM, W. T. & OLSEN, J. L. (2006c). Origin of *Fucus serratus* (Heterokontophyta; Fucaceae) populations in Iceland and the Faroes: a microsatellite based assessment. *European Journal of Phycology*, 41, 235–246.
- COYER, J. A., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2007). Hybridization and introgression in a mixed population of the intertidal seaweeds *Fucus evanescens* and *F. serratus*. *Journal of Evolutionary Biology*, 20, 2322–2333.
- COYER, J. A., PETERS, A. F., HOARAU, G., STAM, W. T. & OLSEN, J. L. (2002). Hybridization of the marine seaweeds *Fucus serratus* and *Fucus evanescens* (Heterokontophyta: Phaeophyceae) in a 100-year-old zone of secondary contact. *Proceedings of the Royal Society of London B Series*, 269, 1829–1834.
- COYER, J. A., PETERS, A. F., STAM, W. T. & OLSEN, J. L. (2003). Post ice-age recolonization and differentiation of *Fucus serratus* L. (Phaeophyceae; Fucaceae) populations in Northern Europe. *Molecular Ecology*, 12, 1817–1829.
- COYNE, J. A. & ORR, H. A. (2004). *Speciation*. Sunderland, MA: Sinauer Associates.

- DAUGBJERG, N. & ANDERSEN, R. A. (1997a). Phylogenetic analyses of the *rbcL* sequences from Haptophytes and Heterokont algae suggest their chloroplasts are unrelated. *Molecular Biology and Evolution*, *14*, 1242–1251.
- DAUGBJERG, N. & ANDERSEN, R. A. (1997b). A molecular phylogeny of the heterokont algae based on analyses of chloroplast encoded *rbcL* sequence data. *Journal of Phycology*, *33*, 1031–1041.
- DAYRAT, B. (2005). Towards integrative taxonomy. *Biological Journal of the Linnean Society*, *85*, 407–415.
- DE CLERCK, O., LELIAERT, F., VERBRUGGEN, H., LANE, C. E., CAMPOS DE PAULA, J., PAYO, D. A. & COPPEJANS, E. (2006). A revised classification of the Dictyoteae (Dictyotales, Phaeophyceae) based on *rbcL* and 26S ribosomal DNA sequence analysis. *Journal of Phycology*, *42*, 1271–1288.
- DE QUEIROZ, K. (1998). The general lineage concept of species, species criteria, and the process of speciation: A conceptual unification and terminological recommendations. In Howard, D. J. & Berlocher, S. H. (eds.) *Endless Forms: Species and Speciation* (Chapter 5, pp. 57–75). Oxford, United Kingdom: Oxford University Press.
- DE QUEIROZ, K. (2007). Species concepts and species delimitation. *Systematic Biology*, *56*, 879–886.
- DE QUEIROZ, K., & DONOGHUE, M. J. (1988). Phylogenetic systematics and the species problem. *Cladistics*, *4*, 317–338.
- DE REVIERS, B. & ROUSSEAU, F. (1999). Towards a new classification of the brown algae. *Progress in Phycological Research*, *13*, 107–201.
- DE TONI, G. B. (1895). *Sylloge algarum omnium hucusque cognitarum. Vol. III. Fucoideae*. Padua, Italy: Sumptibus auctoris.
- DEMES, K. W., GRAHAM, M. H. & SUSKIEWICZ, T. S. (2009). Phenotypic plasticity reconciles incongruous molecular and morphological taxonomies: The giant kelp, *Macrocystis* (Laminariales, Phaeophyceae), is a monospecific genus. *Journal of Phycology*, *45*, 1266–1269.
- DESALLE, R. (2006). Species discovery versus species identification in DNA barcoding efforts: Response to Rubinoff. *Conservation Biology*, *20*, 1545 – 1547.
- DILLEHAY, T. D., RAMÍREZ, C., PINO, M., COLLINS, M. B., ROSSEN, J. & PINO-NAVARRO, J. D. (2008). Monte Verde: Seaweed, food, medicine, and the peopling of South America. *Science*, *320*, 784–786.
- DRAISMA, S. G. A., OLSEN, J. L., STAM, W. T., & VAN REINE, P. W. F. (2002). Phylogenetic relationships within the Sphacelariales (Phaeophyceae): *rbcL*, RUBISCO spacer and morphology. *European Journal of Phycology*, *37*, 385–401.
- DRAISMA, S. G. A., VAN REINE, P. W. F., STAM, W. T. & OLSEN, J. L. (2001). A reassessment of phylogenetic relationships within the Phaeophyceae based on Rubisco large subunit and ribosomal DNA sequences. *Journal of Phycology*, *37*, 586–603.
- DROMGOOLE, F. I. (1973). *A contribution to the biology of the genus Carpophyllum Grev*. Unpublished PhD. Thesis. Auckland, New Zealand: University of Auckland.
- DRUEHL, L. D. & SAUNDERS, G. W. (1992). Molecular explorations in kelp evolution. *Progress in Phycological Research*, *8*, 47–83.
- DRUEHL, L. D., COLLINS, J. D., LANE, C. E. & SAUNDERS, G. W. (2005). An evaluation of methods used to assess intergeneric hybridization in kelp using Pacific Laminariales (Phaeophyceae). *Journal of Phycology*, *41*, 250–262.
- ENDO, H., PARK, E.-J., SATO, Y., MIZUTA, H. & SAGA, N. (2009). Intraspecific diversity of *Undaria pinnatifida* (Harvey) Suringar (Laminariales, Phaeophyta) from Japan, China and Korea, based on the *cox1* gene and ITS2 sequences. *Fisheries Science*, *75*, 393–400.
- ENGEL, C. R., DAGUIN, C. & SERRÃO, E. (2005). Genetic entities and mating systems in hermaphroditic *Fucus spiralis* and its close dioecious relative *F. vesiculosus* (Fucaceae, Phaeophyceae). *Molecular Ecology*, *14*, 2033–2046.

- FERGUSON, J. W. H. (2002). On the use of genetic divergence for identifying species. *Biological Journal of the Linnean Society*, 75, 509–516.
- FLETCHER, R. L. (1987). *Seaweeds of the British Isles. Vol. 3. Fucophyceae (Phaeophyceae)*. London: British Museum (Natural History).
- FOX, C. H. & SWANSON, A. K. (2007). Nested PCR detection of microscopic life-stages of laminarian macroalgae and comparison with adult forms along intertidal height gradients. *Marine Ecology Progress Series*, 332, 1–10.
- FRASER, C. I., HAY, C. H., SPENCER, H. G. & WATERS, J. M. (2009a). Genetic and morphological analyses of the southern Bull Kelp *Durvillaea antarctica* (Phaeophyceae: Durvilliales) in New Zealand reveals cryptic species. *Journal of Phycology*, 45, 436–443.
- FRASER, C. L., NIKULA, R., SPENCER, H. G. & WATERS, J. M. (2009b). Kelp genes reveal effects of subantarctic sea ice during the Last Glacial Maximum. *Proceedings of the National Academy of Sciences of the USA*, 106, 3249–3253.
- FRASER, C. I., SPENCER, H. G. & WATERS, J. M. (2009c). Glacial oceanographic contrasts explain the phylogeography of Australian bull kelp. *Molecular Ecology*, 18, 2287–2296.
- FRASER, C. I., THIEL, M., SPENCER, H. G. & WATERS, J. M. (2010a). Contemporary habitat discontinuity and historic glacial ice drive genetic divergence in Chilean kelp. *BMC Evolutionary Biology*, 19, 203. doi:10.1186/1471-2148-10-203.
- FRASER, C. I., WINTER, D. J., SPENCER, H. G. & WATERS, J. M. (2010b). Multigene phylogeny of the southern bull-kelp genus *Durvillaea* (Phaeophyceae: Fucales). *Molecular Phylogenetics and Evolution*, 57, 1301–1311.
- GODFRAY, H. C. J., CLARK, B. R., KITCHING, I. J., MAYO, S. J. & SCOBLE, M. J. (2007). The web and the structure of taxonomy. *Systematic Biology*, 56, 943–955.
- GOLDING, G. B., HANNER, R. & HEBERT, P. D. N. (2009). Preface. *Molecular Ecology Resources*, 9, (Suppl. 1) iv–vi.
- GOOD, D. A. & WAKE, D. B. (1992). *Geographic variation and speciation in the Torrent Salamanders of the genus Rhyacotriton (Caudata: Rhyacotritonidae)* (University of California Publications in Zoology No. 126). California: University of California Press.
- GOURBIÈRE, S. & MALLET, J. (2009). Are species real? The shape of the species boundary with exponential failure, reinforcement and the “missing snowball”. *Evolution*, 64, 1–24.
- GRAHAM, M. H., KINLAN, B. P. & GROSBERG, R. K. (2001). Post-glacial redistribution and shifts in productivity of giant kelp forests. *Proceedings of the Royal Society of London B Series*, 277, 399–406.
- GUIRY, M. D. & GUIRY, G. M. (2011). *Algaebase*. Retrieved February 1, 2011, from <http://www.algaebase.org>.
- HARROLD, C. & LISINA, S. (1989). Radio-tracking rafts of giant kelp: local production and regional transport. *Journal of Experimental Marine Biology and Ecology*, 130, 237–251.
- HARVEY, J. B. J. & GOFF, L. J. (2006). A reassessment of species boundaries in *Cystoseira* and *Halidrys* (Phaeophyceae, Fucales) along the North American West Coast. *Journal of Phycology*, 42, 707–720.
- HAWES, N. A. (2008). *Nearshore dispersal and reproductive viability of intertidal furoid algae: How effective is drift in local to regional dispersal?* Unpublished masters thesis. Christchurch, New Zealand: University of Canterbury.
- HEBERT, P. D. N. & GREGORY, T. R. (2005). The promise of DNA barcoding for taxonomy. *Systematic Biology*, 54, 852–859.
- HEBERT, P. D. N., CYWINSKA, A., BALL, S. L. & DE WAARD, J. R. (2003a). Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London B Series*, 270, 313–321.

- HEBERT, P. D. N., RATNASINGHAM, S. & DE WAARD, J. R. (2003b). Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society of London B Series*, 270, (Suppl. 1) S96–S99. doi:10.1098/rsbl.2003.0025.
- HEESCH, S., PETERS, A. F., BROOM, J. E. S. & HURD, C. L. (2008). Affiliation of the parasitic brown alga *Herpodiscus durvillaeae* with the Sphacelariales based on DNA sequence comparisons and morphological observations. *European Journal of Phycology*, 43, 283–295.
- HENKEL, S. K., HOFMANN, G. E. & WHITMER, A. C. (2007). Morphological and genetic variation in *Egregia menziesii* over a latitudinal gradient. *Botanica Marina*, 50, 159–170.
- HERNÁNDEZ-CARMONA, G., HUGHES, B. & GRAHAM, M. H. (2006). Reproductive longevity of drifting kelp *Macrocystis pyrifera* (Phaeophyceae) in Monterey Bay, USA. *Journal of Phycology*, 42, 1199–1207.
- HEWITT, G. M. (1988). Hybrid zones – natural laboratories for evolutionary studies. *Trends in Ecology and Evolution*, 3, 158–167.
- HEWITT, G. M. (2004). Genetic consequences of climatic oscillations in the Quaternary. *Philosophical Proceedings of the Royal Society of London B Series*, 359, 183–195.
- HEY, J. (2006). On the failure of modern species concepts. *Trends in Ecology and Evolution*, 21, 447–450.
- HIGHTON, R. (1989). Biochemical evolution in the slimy Salamanders of the *Plethodon glutinosus* complex in the eastern United States. Part 1. Geographical protein variation. *Illinois Biological Monographs*, 57, 1–78.
- HIGHTON, R. (1990). Taxonomic treatment of genetically differentiated populations. *Herpetologica*, 46, 114–121.
- HIGHTON, R. (1995). Speciation in Eastern North American Salamanders of the Genus *Plethodon*. *Annual Review of Ecology and Systematics*, 26, 579–600.
- HO, C. L., PHANG, S. M. & PANG, T. (1995). Molecular characterisation of *Sargassum polycystum* and *S. siliculosum* (Phaeophyta) by polymerase chain reaction using random amplified polymorphic DNA (RAPD) primers. *Journal of Applied Phycology*, 7, 33–41.
- HOARAU, G., COYER, J. A., VELDSINK, J. H., STAM, W. T. & OLSEN, J. L. (2007). Glacial refugia and recolonization pathways in the brown seaweed *Fucus serratus*. *Molecular Ecology*, 16, 3606–3616.
- HODGE, F. J. (2009). *Hybridisation in the brown algae Carpophyllum: Investigating morphology, distribution and wave exposure*. Unpublished masters thesis. Wellington, New Zealand: Victoria University of Wellington.
- HORIGUCHI, T. & YOSHIDA, T. (1998). The phylogenetic affinities of *Myagropsis myagroides* (Fucales, Phaeophyceae) as determined from 18S rDNA sequences. *Phycologia*, 37, 237–245.
- HULL, D. L. (1965). The effect of essentialism on taxonomy—two thousand years of stasis. *The British Journal for the Philosophy of Science*, 25, 314–326 (part 1), 16, 1–18 (part II).
- HULL, S. L., SCOTT, G. W. & JOHNSON, L. J. (2001). An investigation into the genetic variation in four fucal species using cellulose acetate electrophoresis. *Botanica Marina*, 44, 119–123.
- JONES (KAIN), J., BUCHANAN, J., BOO, S. M. & LEE, K. M. (2010). *Colpomenia bullosa* crust masquerading as *Ralfsia verrucosa* in southeast Australia. *Phycologia*, 49, 617–627.
- KAWAI, H., HANYUDA, T., DRAISMA, S. G. A. & MÜLLER, D. G. (2007). Molecular phylogeny of *Discosporangium mesartrocarpum* (Phaeophyceae) with a reinstatement of the order Discosporangiales. *Journal of Phycology*, 43, 186–194.
- KAWAI, H., HANYUDA, T., LINDBERG, M. & LINDSTROM, S. C. (2008). Morphology and molecular phylogeny of *Aureophycus aleuticus* Gen. et Sp. Nov. (Laminariales, Phaeophyceae) from the Aleutian Islands. *Journal of Phycology*, 44, 1013–1021.
- KAWAI, H. & SASAKI, H. (2000). Molecular phylogeny of the brown algal genera *Akkesiphycus* and *Halosiphon* (Laminariales), resulting in the circumscription of the new families Akkesiphycaceae and Halosiphonaceae. *Phycologia*, 39, 416–428.

- KAWAI, H. & SASAKI, H. (2004). Morphology, life history and molecular phylogeny of *Stschapovia flagellaris* (Tilopteridales, Phaeophyceae) and the erection of the Stschapoviaceae fam. nov. *Journal of Phycology*, 40, 1156–1169.
- KAWAI, H., SASAKI, H., MAEDA, Y. & HENRY, E. C. (2005). Morphology and molecular phylogeny of *Phaeostrophion irregulare* (Phaeophyceae) with a proposal for Phaeostrophiaceae fam. nov., and a review of the Ishigeaceae. *Phycologia*, 44, 169–182.
- KAWAI, H., SASAKI, H., MAEDA, Y. & ARAI, S. (2001). Morphology, life history, and molecular phylogeny of *Chorda rigida* sp. nov. (Laminariales, Phaeophyceae) from the Sea of Japan and the genetic diversity of *Chorda filum*. *Journal of Phycology*, 37, 130–142.
- KIM, S.-K. & KAWAI, H. (2002). Taxonomic revision of *Chordaria flagelliformis* (Chordariales, Phaeophyceae) including novel use of the intragenic spacer region of rDNA for phylogenetic analysis. *Phycologia*, 41, 328–339.
- KIM, S.-H., PETERS, A. F. & KAWAI, H. (2003). Taxonomic revision of *Sphaerotrichia divaricata* (Ectocarpales, Phaeophyceae), with a reappraisal of *S. firma* from the north-west Pacific. *Phycologia*, 42, 183–192.
- KNOWLES, L. L. & CARSTENS, B. C. (2007). Delimiting species without monophyletic gene trees. *Systematic Biology*, 56, 887–895.
- KNOWLTON, N., WEIGT, L. A., SOLÓRZANO, L. A., MILLS, D. E. K. & BERMINGHAM, E. (1993). Divergence in proteins, mitochondrial DNA, and reproductive compatibility across the Isthmus of Panama. *Science*, 260, 1629–1632.
- KOGAME, K. (1996). Morphology and life history of *Scytosiphon canaliculatus* comb. nov. (Scytosiphonales, Phaeophyceae) from Japan. *Phycological Research*, 44, 85–94.
- KOGAME, K. (1997a). Sexual reproduction and life history of *Petalonia fascia* (Scytosiphonales, Phaeophyceae). *Phycologia*, 36, 389–394.
- KOGAME, K. (1997b). Life histories of *Colpomenia sinuosa* and *Hydroclathrus clathratus* (Scytosiphonaceae, Phaeophyceae) in culture. *Phycological Research*, 45, 227–231.
- KOGAME, K., HORIGUCHI, T. & MASUDA, M. (1999). Phylogeny of the order Scytosiphonales (Phaeophyceae) based on DNA sequences of *rbcL*, partial *rbcS*, and partial LSU nrDNA. *Phycologia*, 38, 496–502.
- KOGISHI, K., KITAYAMA, T., MILLER, K. A., HANYUDA, T. & KAWAI, H. (2010). Phylogeography of *Cutleria cylindrica* (Cutleriales, Phaeophyceae) in northeastern Asia, and the identity of an introduced population in California. *Journal of Phycology*, 46, 553–558.
- KOGAME, K. & MASUDA, M. (2001). Crustose sporophytes of *Colpomenia bullosa* (Scytosiphonaceae, Phaeophyceae) in nature. *Cryptogamie, Algologie*, 22, 201–208.
- KOGAME, K. & YAMAGISHI, Y. (1997). The life history and phenology of *Colpomenia peregrina* (Scytosiphonales, Phaeophyceae) from Japan. *Phycologia*, 36, 337–344.
- KRAAN, S. & GUIRY, M. D. (1998). *Strain selection in the edible brown seaweed Alaria esculenta: Genetic fingerprinting and hybridization studies under laboratory conditions* (Marine Resources Series No. 8). Ireland: Marine Institute.
- KRAAN, S. & GUIRY, M. D. (2000). Sexual hybridization experiments and phylogenetic relationships as inferred from Rubisco spacer sequences in the genus *Alaria* (Phaeophyceae). *Journal of Phycology*, 36, 190–198.
- KRAAN, S., RUENESS, J., GUIRY, M. D. (2001). Are North Atlantic *Alaria esculenta* and *A. grandifolia* (Alariaceae, Phaeophyceae) conspecific? *European Journal of Phycology*, 36, 35–42.
- KRAFT, G. T., SAUNDERS, G. W., ABBOTT, I. A. & HAROUN, R. J. (2004). A uniquely calcified brown alga from Hawaii: *Newhousia imbricata* gen. et sp. nov. (Dictyotales, Phaeophyceae). *Journal of Phycology*, 40, 383–394.
- KRESS, W. J., ERICKSON, D. L., JONES, F. A., SWENSON, N. G., PEREZ, R., SANJUR, O. & BERMINGHAM, E. (2009). Plant DNA barcodes and a community phylogeny of a tropical forest

- dynamics plot in Panama. *Proceedings of the National Academy of Sciences of the U.S.A.*, doi: 10.1073/pnas.0909820106
- KUCERA, H. & SAUNDERS, G. W. (2008). Assigning morphological variants of *Fucus* (Fucales, Phaeophyceae) in Canadian waters to recognized species using DNA barcoding. *Botany*, 86, 1065–1079.
- KUSUMO, H. T. & DRUEHL, L. D. (2000). Variability over space and time in the genetic structure of the winged kelp *Alaria marginata*. *Marine Biology*, 136, 397–409.
- LAGO-LESTÓN, A, MOTA, C., KAUTSKY, L. & PEARSON, G. A. (2010). Functional divergence in heat shock response following rapid speciation of *Fucus* spp. in the Baltic Sea. *Marine Biology*, 157, 683–688.
- LANE, C. E., LINDSTROM, S. C. & SAUNDERS, G. W. (2007). A molecular assessment of northeast Pacific *Alaria* species (Laminariales, Phaeophyceae) with reference to the utility of DNA barcoding. *Molecular Phylogenetics and Evolution*, 44, 634–648.
- LANE, C. E., MAYES, C. M., DRUEHL, L. & SAUNDERS, G. W. (2006). A multi-gene molecular investigation of the kelp (Laminariales, Phaeophyceae) resolves competing phylogenetic hypotheses and supports substantial taxonomic re-organization. *Journal of Phycology*, 42, 493–512.
- LANE, C. E. & SAUNDERS, G. W. (2005). Molecular investigation reveals epi/endophytic extrageneric kelp (Laminariales, Phaeophyceae) gametophytes colonizing *Lessoniopsis littoralis* thalli. *Botanica Marina*, 48, 426–436.
- LECLERC, M. C., BARRIEL, V., LECOINTRE, G. & DE REVIERS, B. (1998). Low divergence in rDNA ITS sequences among five species of *Fucus serratus* (Phaeophyceae) suggests a very recent radiation. *Journal of Molecular Evolution*, 46, 115–120.
- LEE, E. Y., PEDERSEN, P. M., & LEE, I. K. (2002). *Neoleptonema yongpili* E.-Y. Lee & I.K. Lee, gen. et sp. nov. (Phaeophyceae), based on morphological characters and RuBisCO spacer sequences. *European Journal of Phycology*, 37, 237–245.
- LEE, W. J. & BAE, K. S. (2002). Phylogenetic relationship among several genera of Dictyotaceae (Dictyotales, Phaeophyceae) based on 18S rRNA and partial rbcL gene sequences. *Marine Biology*, 140, 1107–1115.
- LEE, Y. K., YOON, H. S., & BOO, S. M. (1998). Phylogenetic relationships between *Pelvetia* and *Pelvetiopsis* (Phaeophyceae) based on small subunit ribosomal DNA sequences. *Journal of Plant Biology*, 41, 103–109.
- LEE, Y. K., YOON, H. S., MOTOMURA, T., KIM, Y. J. & BOO, S. M. (1999). Phylogenetic relationships between *Pelvetia* and *Pelvetiopsis* (Fucaceae, Phaeophyta) inferred from sequences of the RuBisCo spacer region. *European Journal of Phycology*, 34, 205–211.
- LELIAERT, F., VERBRUGGEN, H., WYSOR, B. & DE CLERCK, O. (2009). DNA taxonomy in morphologically plastic taxa: Algorithmic species delimitation in the *Boodlea* complex (Chlorophyta: Cladophorales). *Molecular Phylogenetics and Evolution*, 53, 122–133.
- LEWIS, R. J. (1996). Hybridization of brown algae: Compatibility and speciation. In Chaudhary, B. R. & Agrawal, S. B. (Eds.) *Cytology, Genetics and Molecular Biology of Algae*. (pp. 275–289). Amsterdam: SPB Academic Publishing.
- LEWIS, R. J. & NEUSHUL, M. (1994). Northern and southern hemisphere hybrids of *Macrocystis* (Phaeophyceae). *Journal of Phycology*, 30, 346–353.
- LEWIS, R. J., NEUSHUL, M. & HARGER, B. W. W. (1986). Interspecific hybridization of the species of *Macrocystis* in California. *Aquaculture*, 57, 203–210.
- LILLY, E. L., HALANYCH, K. M. & ANDERSON, D. M. (2007). Species boundaries and global biogeography of the *Alexandrium tamarens* complex (Dinophyceae). *Journal of Phycology*, 43, 1329–1338.
- LINDAUER, V. W., CHAPMAN, V. J. & AIKEN, M. (1961). The Marine Algae of New Zealand II: Phaeophyceae. *Nova Hedwigia*, 3, 129–350.

- LIPTACK, M. K. & DRUEHL, L. D. (2000). Molecular evidence for an interfamilial laminariales cross. *European Journal of Phycology*, 35, 135–142.
- LITAKER, R. W., ZANDERSEA, M. W., KIBLER, S. R., REECE, K. S., STOKES, N. A., LUTZONI, F. M., ET AL. (2007). Recognizing dinoflagellate species using ITS rDNA sequences. *Journal of Phycology*, 43, 344–355.
- MACAYA, E. C. & ZUCCARELLO, G. C. (2010). DNA barcoding and genetic divergence in the giant kelp *Macrocystis* (Laminariales). *Journal of Phycology*, 46, 736–742.
- MACAYA, E., BOLTAÑA, S., BUSCHMANN, A., HINOJOSA, I., MACCHIAVELLO, J., VALDIVIA, N., ET AL. (2005). Presence of sporophylls in floating kelp rafts of *Macrocystis* spp. (Phaeophyceae) along the Chilean Pacific Coast. *Journal of Phycology*, 41, 913–922.
- MAGGS, C. A., CASTILHO, R., FOLTZ, D., HENZLER, C., JOLLY, M. T., KELLY, J., ET AL. (2008). Evaluating signatures of glacial refugia for North Atlantic benthic marine taxa. *Ecology*, 89, (Suppl. 1) S108–S122.
- MALLET, J. (2008). Hybridization, ecological races and the nature of species: empirical evidence for the ease of speciation. *Philosophical Transactions of the Royal Society of London B Series*, 363, 2971–2986.
- MATHIESON, A. C., DAWES, C. J., WALLACE, A. L. & KLEIN, A. S. (2006). Distribution, morphology, and genetic affinities of dwarf embedded *Fucus* populations from the Northwest Atlantic Ocean. *Botanica Marina*, 49, 283–303.
- MATHIESON, A. C., NORTON, T. A. & NEUSHUL, M. (1981). The taxonomic implications of genetic and environmentally induced variations in seaweed morphology. *The Botanical Review*, 47, 313–347.
- MAYDEN, R. L. (1997). A hierarchy of species concepts: The denouement in the saga of the species problem. In Claridge, M. F., Dawah, H. A. & Wilson, M. R. *Species: The units of biodiversity*. (pp. 381–384). London: Chapman and Hall.
- MAYR, E. (1942). *Systematics and the origin of species*. New York: Columbia University Press.
- MAYR, E. (1996). What is a species and what is not? *Philosophy of Science*, 63, 262–277.
- MCDEVIT, D. C. & SAUNDERS, G. W. (2009). On the utility of DNA barcoding for species differentiation among brown macroalgae (Phaeophyceae) including a novel extraction protocol. *Phycological Research*, 57, 131–141.
- MCDEVIT, D. C. & SAUNDERS, G. W. (2010). A DNA barcode examination of the Laminariaceae (Phaeophyceae) in Canada reveals novel biogeographical and evolutionary insights. *Phycologia*, 49, 235–248.
- MONIZ, B. J. M. & KACZMARSKA, I. (2009). Barcoding diatoms: Is there a good marker? *Molecular Ecology Resources*, 9, (Suppl. 1) 65–74.
- MÜLLER, D. G. (1988). Studies on sexual compatibility between *Ectocarpus siliculosus* (Phaeophyceae) from Chile and the Mediterranean Sea. *Helgoländer wissenschaftliche Meeresuntersuchungen*, 42, 469–476.
- MÜLLER, D. G., PARODI, E. R. & PETERS, A. F. (1998). *Asterocladon lobatum* gen. et sp. nov., a new brown alga with a stellate chloroplast arrangement, and its systematic position judged from nuclear rDNA sequences. *Phycologia*, 37, 425–432.
- NEIVA, J., PEARSON, G. A., VALERO, M. & SERRAO, E. A. (2010). Surfing the wave on a borrowed board: range expansion and spread of introgressed organellar genomes in the seaweed *Fucus ceranoides* L. *Molecular Ecology*, 19, 4812–4822.
- NI-NI-WIN, HANYUDA, T., ARAI, S., UCHIMURA, M., PRATHEP, A., DRAISMA, S. G. A., SOE-HTUN & KAWAI, H. (2010). Four new species of *Padina* (Dictyotales, Phaeophyceae) from the western Pacific Ocean, and reinstatement of *Padina japonica*. *Phycologia*, 49, 136–153.
- NOOR, M. A. F., KLIMAN, R. M. & MACHADO, C. A. (2001). Evolutionary history of microsatellites in the obscure group of *Drosophila*. *Molecular Biology and Evolution*, 18, 551–556.

- O'BRIEN, S. J. & MAYR, E. (1991). Bureaucratic mischief: Recognizing Endangered species and subspecies. *Science*, 251, 1187–1188.
- OLSEN, J. L., ZECHMAN, F. W., HOARAU, G., COYER, J. A., STAM, W. T., VALERO, M. & ABERG, P. (2010). The phylogeographic architecture of the furoid seaweed *Ascophyllum nodosum*: an intertidal 'marine tree' and survivor of more than one glacial-interglacial cycle. *Journal of Biogeography*, 37, 842–856.
- PARSONS, M. J. (1985). New Zealand seaweed flora and its relationships. *New Zealand Journal of Marine and Freshwater Research*, 19, 131–138.
- PEREYRA, R. T., BERGSTRÖM, L., KAUTSKY, L. & JOHANNESSON, K. (2009). Rapid speciation in a newly opened postglacial marine environment, the Baltic Sea. *BMC Evolutionary Biology*, 9, 70. doi: 10.1186/1471-2148-9-70
- PÉREZ-RUZAFÁ, I., GALLARDO, T. & GOMEZCANCIO, R. (1993). Numerical taxonomy of some taxa of the genus *Fucus* in the Iberian Peninsula. *Hydrobiologia*, 261, 81–90.
- PERRIN, C., DAGUIN, C., VAN DE VLIET, M., ENGEL, C. R., PEARSON, G. A. & SERRÃO, E. A. (2007). Implications of mating system for genetic diversity of sister algal species: *Fucus spiralis* and *Fucus vesiculosus* (Heterokontophyta, Phaeophyceae). *European Journal of Phycology*, 42, 219–230.
- PETERS, A. F. (1987). Reproduction and sexuality in the Chordariales (Phaeophyceae). A review of culture studies. *Progress in Phycological Research*, 5, 223–263.
- PETERS, A. F. (1998). Ribosomal DNA sequences support taxonomic separation of the two species of *Chorda*: reinstatement of *Halosiphon tomentosus* (Lyngbye) Jaasund (Phaeophyceae, Laminariales). *European Journal of Phycology*, 33, 65–71.
- PETERS, A. F. & BURKHARDT, E. (1998). Systematic position of the kelp endophyte *Laminarionema elsbetiae* (Ectocarpales *sensu lato*, Phaeophyceae) inferred from nuclear ribosomal DNA sequences. *Phycologia*, 37, 114–120.
- PETERS, A. F. & CLAYTON, M. N. (1998). Molecular and morphological investigations of three brown algal genera with stellate plastids: Evidence for Scytothamnales ord. nov. (Phaeophyceae). *Phycologia*, 37, 106–113.
- PETERS, A. F. & RAMÍREZ, M. E. (2001). Molecular phylogeny of small brown algae, with special reference to the systematic position of *Caepidium antarcticum* (Adenocystaceae, Ectocarpales). *Cryptogamie, Algologie*, 22, 187–200.
- PETERS, A. F., RAMÍREZ, M. E. & RÜLKE, A. (2000). The phylogenetic position of the subantarctic marine macroalga *Desmarestia chordalis* (Phaeophyceae) inferred from nuclear ribosomal ITS sequences. *Polar Biology*, 23, 95–99.
- PETERS, A. F., VAN WIJK, S. J., CHO, G. Y., SCORNET, D., HANYUDA, T., KAWAI, H., *ET AL.* (2010). Reinstatement of *Ectocarpus crouaniorum* Thuret in Le Jolis as a third common species of *Ectocarpus* (Ectocarpales, Phaeophyceae) in Western Europe, and its phenology at Roscoff, Brittany. *Phycological Research*, 58, 157–170.
- PHILLIPS, N., BURROWES, R., ROUSSEAU, F., DE REVIERS, B. & SAUNDERS, G. W. (2008). Resolving evolutionary relationships among the brown algae using chloroplast and nuclear genes. *Journal of Phycology*, 44, 394–405.
- PIGLUCCI, M. (2003). Species as a family resemblance concept. *BioEssays*, 25, 596–602
- PONS, J., BARRACLOUGH, T. G., GOMEZ-ZURITA, J., CARDOSO, A., DURAN, D. P., HAZELL, S., *ET AL.* (2006). Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology*, 55, 595–609.
- PORTER, A. H. (1990). Testing nominal species boundaries using gene flow statistics: The taxonomy of two hybridizing Admiral butterflies (*Limenitis*: Nymphalidae). *Systematic Biology*, 39, 131–147.
- PRESGRAVES, D. C. (2010). The molecular evolutionary basis of species formation. *Nature Reviews Genetics*, 11, 175–180.

- PRITCHARD, J. K., STEPHENS, M. & DONNELLY, P. (2000). Inference of population structure using multilocus genotype data. *Genetics*, 155, 945–959.
- QUICKE, D. L. J. (1993). *Principles and techniques of contemporary taxonomy*. Glasgow, United Kingdom: Blackie Academic & Professional.
- RACH, J., DESALLE, R., SARKAR, I. N., SCHIERWATER, B. & HADRY, H. (2008). Character-based DNA barcoding allows discrimination of genera, species and populations in Odonata. *Philosophical Transactions of the Royal Society of London B Series*, 275, 237–247.
- RICE, E. L. (1989). A statistical morphological analysis and taxonomic revision of the genus *Xiphophora* (Fucaceae). *Botanical Journal of the Linnean Society*, 100, 355–387.
- RICE, E. L. & CHAPMAN, A. R. O (1985). A numerical taxonomic study of *Fucus distichus* (Phaeophyta). *Journal of the Marine Biological Association of the United Kingdom*, 65, 433–459.
- RIESEBERG, L. H. (1995). The role of hybridization in evolution: Old wine in new skins. *American Journal of Botany*, 82, 955–953.
- ROBERSON, L. M. & COYER, J. A. (2004). Variation in blade morphology of the kelp *Eisenia arborea*: incipient speciation due to local water motion? *Marine Ecology Progress Series*, 282, 115–128.
- ROUSSEAU, F. & DE REVIERS, B. (1999a). Circumscription of the order Ectocarpales (Phaeophyceae): bibliographical synthesis and molecular evidence. *Cryptogamie, Algologie*, 20, 5–18.
- ROUSSEAU, F. & DE REVIERS, B. (1999b). Phylogenetic relationships within the Fucales (Phaeophyceae) based on combined partial SSU and LSU rDNA sequence data. *European Journal of Phycology*, 34, 53–64.
- ROUSSEAU, F., BURROWES, R., PETERS, A. F., KUHLENKAMP, R., & DE REVIERS, B. (2001). A comprehensive phylogeny of the Phaeophyceae based on nrDNA resolves the earliest divergences. *Comptes Rendues de la Academie des Sciences de Paris – Sciences de la vie*, 324, 305–319.
- ROUSSEAU, F., DE REVIERS, B., LECLERC, M.-C., ASENSI, A. & DELÉPINE, R. (2000). Adenocystaceae fam. nov. (Phaeophyceae) based on morphological and molecular evidence. *European Journal of Phycology*, 35, 35–43.
- ROUSSEAU, F., LECLERC, M.-C. & REVIERS, B. DE (1997). Molecular phylogeny of European Fucales (Phaeophyceae) based on large subunit rDNA sequence comparisons. *Phycologia*, 36, 438–446.
- RUSSELL, G. & FLETCHER, R. L. (1975). A numerical taxonomic study of the British Phaeophyta. *Journal of the Marine Biological Association of the United Kingdom*, 55, 763–783.
- SASS, C., LITTLE, D. P., STEVENSON, D. W. & SPECHT, C. D. (2007). DNA Barcoding in the Cycadales: Testing the potential of proposed barcoding markers for species identification of Cycads. *PLoS ONE*, 2, e1154. doi:10.1371/journal.pone.0001154.
- SAUNDERS, G. W. (2009). Routine DNA barcoding of Canadian Gracilariales (Rhodophyta) reveals the invasive species *Gracilaria vermiculophylla* in British Columbia. *Molecular Ecology Resources*, 9, (Suppl. 1) 140–150.
- SAUNDERS, G. W. & DRUEHL, L. D. (1992). Nucleotide sequences of the small sub-unit ribosomal RNA genes from selected Laminariales (Phaeophyta): implications for kelp evolution. *Journal of Phycology*, 28, 544–549.
- SAUNDERS, G. W. & DRUEHL, L. D. (1993). Revision of the kelp family Alariaceae and the taxonomic affinities of *Lessoniopsis* Reinke (Laminariales, Phaeophyta). *Hydrobiologia*, 260/261, 689–697.
- SAUNDERS, G. W. & KRAFT, G. T. (1995). The phylogenetic affinities of *Notheia anomala* (Fucales, Phaeophyceae) as determined from partial small sub-unit RNA sequences. *Phycologia*, 34, 383–389.

- SCHINDEL, D. E. & MILLER, S. E. (2005). DNA barcoding a useful tool for taxonomists. *Nature*, 435, 17.
- SCOTT, G. W. & HARDY, F. G. (1994). Observations of the occurrence of hybrids between two sympatric species of fucoid algae. *Cryptogamie, Algologie*, 15, 297–305.
- SERRÃO, E. A., ALICE, L. A. & BRAWLEY, S. H. (1999). Evolution of the Fucaceae (Phaeophyceae) inferred from nrDNA-ITS. *Journal of Phycology*, 35, 382–394.
- SETCHELL, W. A. & GARDNER, N. L. (1925). The marine algae of the Pacific coast of North America, Part III, Melanophyceae. *University of California Publications in Botany*, 8, 383–898.
- SHAFFER, H. B. & THOMSON, R. (2007). Delimiting Species in Recent Radiations. *Systematic Biology*, 56, 896–906.
- SHERWOOD, A. R., VIS, M. L., ENTWISLE, T. J., NECCHI, O. JR. & PRESTING, G. G. (2008). Contrasting intra versus interspecies DNA sequence variation for representatives of the Batrachospermales (Rhodophyta): Insights from a DNA barcoding approach. *Phycological Research*, 56, 269–279.
- SIEMER, B. L., STAM, W. T., OLSEN, J. L., & PEDERSEN, P. M. (1998). Phylogenetic relationships of the brown algal orders Ectocarpales, Chordariales, Dictyosiphonales and Tilopteridales (Phaeophyceae) based on Rubisco large subunit and spacer sequences. *Journal of Phycology*, 34, 1038–1048.
- SILBERFELD, T., LEIGH, J. W., VERBRUGGEN, H., CRUAUD, C., DE REVIERS, B. & ROUSSEAU, F. (2010). A multi-locus time-calibrated phylogeny of the brown algae (Heterokonta, Ochrophyta, Phaeophyceae): Investigating the evolutionary nature of the “brown algal crown radiation”. *Molecular Phylogenetics and Evolution*, 56, 659–674.
- SITES, J. W. JR. & MARSHALL, J. C. (2004). Operational criteria for delimiting species. *Annual Review of Ecology, Evolution, and Systematics*, 35, 199–227.
- STACHE-CRAIN, B., MÜLLER, D. G. & GOFF, L. J. (1997). Molecular systematics of *Ectocarpus* and *Kuckuckia* (Ectocarpales, Phaeophyceae) inferred from phylogenetic analysis of nuclear and plastid encoded DNA sequences. *Journal of Phycology*, 33, 152–168.
- STARR, J. R., NACZI, R. F. C. & CHOUINARD, B. N. (2009). Plant DNA barcodes and species resolution in sedges (*Carex*, Cyperaceae). *Molecular Ecology Resources*, 9, (Suppl. 1) 151–163.
- STIGER, V., HORIGUCHI, T., YOSHIDA, T., COLEMAN, A. W. & MASUDA, M. (2000). Phylogenetic relationships of *Sargassum* (Sargassaceae, Phaeophyceae) with reference to a taxonomic revision of section Phyllocystae based on ITS-2 nrDNA sequences. *Phycological Research*, 48, 251–260.
- TAN, I. H. & DRUEHL, L. D. (1993). Phylogeny of the Northeast Pacific brown algae (Phaeophyceae) orders as inferred from 18S rRNA gene sequences. *Hydrobiologia*, 260/261, 699–704.
- TAN, I. H. & DRUEHL, L. D. (1994). A molecular analysis of *Analipus* and *Ralfsia* (Phaeophyceae) suggests the order Ectocarpales is polyphyletic. *Journal of Phycology*, 30, 721–729.
- TATARENKOV, A., JÖNSSON, R. B., KAUTSKY, L. & JOHANNESSEN, K. (2007). Genetic structure in populations of *Fucus vesiculosus* (Phaeophyceae) over spatial scales from 10 m to 800 km. *Journal of Phycology*, 43, 675–685.
- TAUTZ, D., ARCTANDER, P., MINELLI, A., THOMAS, R. H. & VOGLER, A. P. (2003). A plea for DNA taxonomy. *Trends in Ecology and Evolution*, 18, 70–74.
- TELLIER, F., MEYNARD, A. P., CORREA, J. A. FAUGERON, S. & VALERO, M. (2009). Phylogeographic analyses of the 30 degrees S south-east Pacific biogeographic transition zone establish the occurrence of a sharp genetic discontinuity in the kelp *Lessonia nigrescens*: Vicariance or parapatry? *Molecular Phylogenetics and Evolution*, 53, 679–693.
- TEMPLETON, A. R. (2001). Using phylogeographic analyses of gene trees to test species status and processes. *Molecular Ecology*, 10, 779–791.
- THOMSEN, M. S. & WERNBERG, T. (2005). What affects the forces required to break or dislodge macroalgae? *European Journal of Phycology*, 40, 139–148.

- TRONHOLM, A., SANSON, M., AFONSO-CARILLO, J., VERBRUGGEN, H. & DE CLERCK, O. (2010a). Niche partitioning and the coexistence of two cryptic *Dictyota* (Dictyotales, Phaeophyceae) species from the Canary Islands. *Journal of Phycology*, *46*, 1075–1087.
- TRONHOLM, A., STEEN F., TYBERGHEIN L., LELIAERT, F., VERBRUGGEN H., SIGUAN M. A. R. & DE CLERCK, O. (2010b). Species delimitation, taxonomy and biogeography of *Dictyota* in Europe (Dictyotales, Phaeophyceae). *Journal of Phycology*, *46*, 1301–1321.
- UWAI, S., KOGAME, K., YOSHIDA, G., KAWAI, H. & AJISAKA, T. (2009). Geographical genetic structure and phylogeography of the *Sargassum horneri/filicinum* complex in Japan, based on the mitochondrial *cox3* haplotype. *Marine Biology*, *156*, 901–911.
- UWAI, S., NAGASATO, C., MOTOMURA, T. & KOGAME, K. (2005). Life history and molecular phylogentic relationships of *Asterocladon interjectum* sp. nov. (Phaeophyceae). *European Journal of Phycology*, *40*, 179–194.
- VAN DEN HOEK, C. (1987). The possible significance for long range dispersal for the biogeography of seaweeds. *Helgoländer wissenschaftliche Meeresuntersuchungen*, *41*, 261–272.
- VAN OPPEN, M. J. H., OLSEN, J. L., STAM, W. T., VAN DEN HOEK, C. & WIENCKE, C. (1993). Arctic-Antarctic disjunctions in the benthic seaweeds *Acrosiphonia arcta* (Chlorophyta) and *Desmarestia viridis/willii* (Phaeophyta) are of recent origin. *Marine Biology*, *115*, 381–386.
- VELASCO, J. D. (2008). Species concepts should not conflict with evolutionary history, but often do. *Studies in History and Philosophy of Biological and Biomedical Sciences*, *39*, 407–414.
- WALLACE, A. L., KLEIN, A. S. & MATHIESON, A. C. (2004). Determining the affinities of salt marsh fucoids using microsatellite markers: Evidence of hybridization and introgression between two species of *Fucus* (Phaeophyta) in a Maine estuary. *Journal of Phycology*, *40*, 1013–1027.
- WATERS, J. M. (2008). Driven by the West Wind Drift? A synthesis of southern temperate marine biogeography, with new directions for dispersalism. *Journal of Biogeography*, *35*, 417–427.
- WESTERMEIER, R., PATIÑO, D. & MÜLLER, D. G. (2007). Sexual compatability and hybrid formation between the giant kelp species *Macrocystis pyrifera* and *M. integrifolia* (Laminariales, Phaeophyta) in Chile. *Journal of Applied Phycology*, *19*, 215–221.
- WIDDOWSON, T. B. (1971). A taxonomic revision of the genus *Alaria* Greville. *Syesis*, *4*, 11–49.
- WIENS, J. J. (2007). Species delimitation: New approaches for discovering diversity. *Systematic Biology*, *56*, 875–878.
- WIENS, J. J. & PENKROT, T. A. (2002). Delimiting species using DNA and morphological variation and discordant species limits in Spiny Lizards (Sceloporus). *Sytematic Biology*, *51*, 69–91.
- WIENS, J. J. & SERVEDIO, M. R. (2000). Species delimitation in systematics: inferring diagnostic differences between species. *Proceedings of the Royal Society of London B Series*, *267*, 631–636.
- WILL, K. W., MISHLER, B. D. & WHEELER, Q. D. (2005). The perils of DNA barcoding and the need for integrative taxonomy. *Systematic Biology*, *54*, 844–851.
- WILLIAMS, S. L. & DI FIORI, R. E. (1996). Genetic diversity and structure in *Pelvetia fastigiata* (Phaeophyta: Fucales): does a small effective neighbourhood size explain fine-scale genetic structure? *Marine Biology*, *126*, 371–382.
- WILSON, P. (1992). On inferring hybridisation from morphological intermediacy. *Taxon*, *41*, 11–23.
- WOMERSELY, H. B. S. (1964). The morphology and taxonomy of *Cystophora* and related genera (Phaeophyta). *Australian Journal of Botany*, *12*, 53–110.
- WOMERSLEY, H. B. S. (1987). *The marine benthic flora of southern Australia. Part II*. Adelaide, Australia: South Australian Government Printing Division.
- WYNNE, M. J. (1969). Life history and systematic studies of some Pacific North American Phaeophyceae (brown algae). *University of California Publications in Botany*, *50*, 1–88.

- YOON, H. S. & BOO, S. M. (1999). Phylogeny of Alariaceae (Phaeophyta) with special reference to *Undaria* based on sequences of the RuBisCo spacer region. *Hydrobiologia*, 398–399, 47–55.
- YOSHIDA, T., STIGER, V. & HORIGUCHI, T. (2000). *Sargassum boreale* sp. nov. (Fucales, Phaeophyceae) from Hokkaido, Japan. *Phycological Research*, 48, 125–131.
- YOUNG, M. W. (1929). Marine fauna of the Chatham Islands. *Transactions and Proceedings of the New Zealand Institute*, 60, 136–166.

Chapter 7

Appendices

7.1 Appendix 1. Alignment of mitochondrial spacer sequences from *Carpophyllum maschalocarpum*. A single specimen is shown for each haplotype/site. Specimen names are truncated as: FieldnumberSitenameHaplotype.

B560Opap01 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B213Kiri02 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B051Dors03 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A737Kiri04 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A738Kiri05 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B307Whal06 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B256Piha07 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B590Ahur08 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B228Wait09 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B320Wait10 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A679Waik11 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B255Rive12 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A865Leig13 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B392Math14 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A847Maral4 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A809Hybr15 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A808Hybr15 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A835Maral6 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A791Maral7 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A855Maral8 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A795Maral9 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B041Sail20 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A399Step21 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A938Otan22 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A799Cang23 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B289Dors24 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B031Sail25 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A940Otan26 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B032Sail27 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B345Sail28 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B033Sail29 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A422NCap30 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A420NCap31 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A421NCap32 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A374Weka33 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A890Hoop34 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B348Whal35 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A344Cang36 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B315Whal37 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A834Mara38 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B223Wait39 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B367Ahip40 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A947Otan43 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A555Cang44 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A566Fras45 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A932Sghy46 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A353Weka47 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A801Cang48 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A378Cang49 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A356Weka50 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B499Horo52 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A408Cang53 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B378Math55 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B379Math56 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B394Math57 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B501Make58 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B521Make59 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B507Make60 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B500Horo61 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B487Horo62 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B435Kait63 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B490Horo64 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A901Hoop65 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B377Math66 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A885Hoop67 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B454Kait67 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B525Mat168 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B565Opap69 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 B035Sail70 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A856Mara71 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC
 A766Mara72 GTAAAGTTTTTCAGTGGGGGTGAAATTTAATCACTTTGATAGGCGGGGGGTGTAAGAGCTGTGAGGTTTTTAGCTGACC

Appendix 1. (Continued)

B560Pap01 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B213Kiri02 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCCAGGCTAGAGTGTCTATGT
 B051Dors03 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCTTAGGCTAGAGTGTCTATGT
 A737Kiri04 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTGGGCTAGAGTGTCTATGT
 A738Kiri05 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B307Whal06 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGGCTGTCTATGT
 B256Piha07 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGACTAGAGTGTCTATGT
 B590Ahur08 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGACTAGAGTGTCTATGT
 B228Wait09 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B220Wait10 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGACTAGGCTGTCTATGT
 A679Waik11 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGACTAGGCTGTCTATGT
 B255Rive12 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGGAGGCTAGAGTGTCTATGT
 A865Leig13 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTAGAGTGTCTATGT
 B392Math14 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A847Mara14 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A809Hybr15 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAGGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A808Hybr15 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A835Mara16 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCAGGCTCGAGTGTCTATGT
 A791Mara17 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCAGGCTCGAGTGTCTATGT
 A855Mara18 CGTACTAATCCTAAGAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A795Mara19 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCGGGCTCGAGTGTCTATGT
 B041Sail20 CGTACTAATCCTAAAAAATATTGAGGCAAATTGCCTTTAAGCTAAACACGCGAAACCGCGGGCTCGAGTGTCTATGT
 A399Step21 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 A938Otan22 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAACTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 A799Cang23 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 B289Dors24 CGTACTAATCCTAAAAAATATTGAGGCAAATTATATTTAAGCTAAACACGCGAAACCGTAGGCCGAGTGTCTATGT
 B031Sail25 CGTACTAATCCTAAAAAATATTGAGGCAAATTACGCTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 A940Otan26 CGTACTAATCCTAAAAAATATTGAGGCAAATAACGTTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 B032Sail27 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 B034Sail28 CGTACTAATCCTAAAAAATATTGAGGCAAATCGCCTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 B033Sail29 CGTACTAATCCTAAAAAATATTGAGGCAAATAGCTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A422Ncap30 CGTACTAATCCTAAAAAATATTGAGTCAAATAACTTTTAAGCTAAACACGCGAAACCGTCGGCCCGGCTGTCTATGT
 A420Ncap31 CGTACTAATCCTAAAAAATATTGAGTCAAATAACTTTTAAGCTAAACATGCGAAACCGTCGGCCCGGCTGTCTATGT
 A421Ncap32 CGTACTAATCCTAAAAAATATTGAGTCAAATAAATTTTAAGCTAAACACGCGAAACCGTCGGCCCGGCTGTCTATGT
 A374Weka33 CGTACTAATCCTAAAAAATATTGAGTCAAATAACTTTTAGGCTAAACACGCGAAACCGTCGGCCCGGCTGTCTATGT
 A890Hoop34 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCCTGACCAGGTGTCTATGT
 B348Whal35 CGTACTAATCCTAAAAAATGTTGAGGCAAATTAC - TTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 A344Cang36 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B315Whal37 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTAGAGTGTCTATGT
 A834Mara38 CGTACTAATCCTAAAAAATATTGAGGCAAAGTACCTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 B223Wait39 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B367Ahip40 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A947Otan43 CGTACTAATCCTAAAAAATATTGAGGCAAATAGCCTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A555Cang44 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCCCCGACCGGCTGTCTATGT
 A566Fras45 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCCCCGACAGGTGTCTATGT
 A932SGhy46 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACACCTGCACAGGTGTCTATGT
 A353Weka47 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCCCCACACAGGTGTCTATGT
 A801Cang48 CGTACTAATCCTAAAAAATATTGAGGCAAATCACTTTTAGGCTAAACACGCGAAACCGCCCCGCACATGTGTCTATGT
 A378Cang49 CGTACTAATCCTAAAAAATATTGAGGCAAATCGCTTTTAAGCTAAACACGCGAAACCGCCTGCATAGGTGTCTATGT
 A356Weka50 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCCTACAAAGGACACCTGTGT
 B499Horo52 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A408Cang53 CGTACTAATCCTAAAAAATATTGAGGCAAATCACTTTTAAGCTAAACACGCGAAACCGTACACACAGGGCACCTGTGT
 B378Math55 CGTACTAATCCTAAAAAATGTTGAGGCAAATTACTTTTAGGCTAAACACGCGAAACCGTGGGCTAGAGTGTCTATGT
 B379Math56 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 B394Math57 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACAAGCGAAACCGCAGGCTAGAGTGTCTATGT
 B501Make58 CGTACTAATCCTAAAAAATATTGAGGCAAATCACTTTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 B521Make59 CGTACTAATCCTAAAAAATATTGAGGCAAATCACTTTTAAGCTAAACATGCGAAACCGTAGGCTCGAGTGTCTATGT
 B507Make60 CGTACTAATCCTAAAAAATATTGAGGCAAATCACC - TTAAGCTAAACATGCGAAACCGTAGGCTCGAGTGTCTATGT
 B500Horo61 CGTACTAATCCTAAAAAATATTGAGGCAAATTACGCTTAAGCTAAACACGCGAAACCGTAGGCTCGGGTGTCTATGT
 B487Horo62 CGTACTAATCCTAAAAAATATTGAGGCAAATTACGCTTAAGCTAAACACGCGAAACCGTGGGCTCGGGTGTCTATGT
 B435Kait63 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTACGCTCGAGTGTCTATGT
 B490Horo64 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTAGAGTGTCTATGT
 A901Hoop65 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACACCTGACCAGGTGTCTATGT
 B377Math66 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCCTGACCAGGTGTCTATGT
 A885Hoop67 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCCTGACCAGGTGTCTATGT
 B454Kait67 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGCCTGACAGGTGTCTATGT
 B525Mat168 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACACGCGAAACCGTAGGCTCGAGTGTCTATGT
 B565Pap69 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAGCACGCGAAACCGCAGGCTCGAGTGTCTATGT
 B035Sail70 CGTACTAATCCTAAAAAATATTGAGGCAAATCGCCTTTAAGCTAAACACGCGAAACCGCGGGCTCGAGTGTCTATGT
 A856Mara71 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGATAAACACGCGAAACCGCAGGCTCGAGTGTCTATGT
 A766Mara72 CGTACTAATCCTAAAAAATATTGAGGCAAATTACTTTTAAGCTAAACATGCGAAACCGCAGGCTCGAGTGTCTATGT

Appendix 1. (Continued)

B560Opap01 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B213Kiri02 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B051Dors03 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A737Kiri04 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A738Kiri05 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B307Whal06 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B256Piha07 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B590Ahur08 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B228Wait09 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B220Wait10 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A679Waik11 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B255Rivel2 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A865Leigl3 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B392Math14 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A847Mara14 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A809Hybr15 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A808Math15 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A835Mara16 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A791Mara17 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A855Mara18 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A795Mara19 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B041Sail20 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A399Step21 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A938Otan22 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A799Cang23 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B289Dors24 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B031Sail25 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A940Otan26 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B032Sail27 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B034Sail28 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B033Sail29 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A422NCap30 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A420NCap31 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A421NCap32 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A374Weka33 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A890Hoop34 GGCTTTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B348Whal35 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A344Cang36 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B315Whal37 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A834Mara38 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B223Wait39 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B367Ahip40 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A947Otan43 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A555Cang44 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A566Fras45 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A932SGhy46 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A353Weka47 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A801Cang48 AGCTCTATTAAATTTTAAATCAAATTAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A378Cang49 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A356Weka50 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B499Horo52 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A408Cang53 GGCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B378Math55 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B379Math56 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B394Math57 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B501Make58 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B521Make59 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B507Make60 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B500Horo61 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B487Horo62 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B435Kait63 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B490Horo64 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A901Hoop65 GGCTTTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B377Math66 GGCTTTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A885Hoop67 GGCTTTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B454Kait67 GGCTTTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B525Mat168 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B565Opap69 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 B035Sail70 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A856Mara71 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC
 A766Mara72 GCCTCTATTAAATTTTAAATCAAAGTAAAGTTATGTCCGTATTTTGTGAAACGTAGTTTACC

A799Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A801Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A805Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A829Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A845Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A833Cangu ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A809Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A758Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A917Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A924Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A922Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A916Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A895Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A909Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A908Chybr ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A796Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A836Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A766Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A855Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A792Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC
A853Cmasc ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCC

A799Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A801Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A805Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A829Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A845Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A833Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCCGAGGCCAGTGGACGG
A809Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A758Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A917Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A924Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A922Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A916Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A895Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A909Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A908Chybr CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A796Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A836Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A766Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A855Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A792Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG

Appendix 2. (Continued)

A853Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A725Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A728Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A716Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCCGTGGACGG
A713Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A730Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A399Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A345Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCCGTGGACGG
A398Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A408Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A659Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A555Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A378Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A379Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGGAGGCCAGTGGACGG
A350Cangu CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A106Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A597Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A509Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A598Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A243Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A240Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A342Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A569Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A220Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A111Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A404Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A418Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A389Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A353Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
A575Cmasc CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCGYGAAGGCCCGTGGACGG

[illegible]

Appendix 2. (Continued)

A404C_{masc} TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
A418C_{masc} TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
A389C_{masc} TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
A353C_{masc} TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
A575C_{masc} TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT

7.3 Appendix 3. Alignment of cloned ITS2 sequences from *Carpophyllum*. Each alignment shows 8–12 sequences recovered from cloned DNA from one specimen.

7.3.1 Sequences from A347, Waterfall Reef, Leigh.

A347_30 ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3C ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3D ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3G ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3H ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3P ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3K ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3L ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3a ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT
 A347_3I ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAAAT

A347_30 CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3C CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3D CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3G CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3H CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3P CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3K CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3L CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3a CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT
 A347_3I CGCCACAGCTTCGGGTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCCGTGT

A347_30 GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT
 A347_3C GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT
 A347_3D GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT
 A347_3G GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT
 A347_3H GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT
 A347_3P GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAG - AAACCGACCGTCCGGGCATTTAT
 A347_3K GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAG - AAACCGACCGTCCGGGCATTTAT
 A347_3L GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAG - AAACCGACCGTCCGGGCATTTAT
 A347_3a GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAG - AAACCGACCGTCCGGGCATTTAT
 A347_3I GTATTTTCACGTACATACTGCCAGCTCGTCCCTGAGTTACCGAAGCCTAGAGAGGAAACCGACCGTCCGGGCATTTAT

A347_30 TCTCT?TGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3C TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3D TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3G TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3H TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3P TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3K TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3L TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3a TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTTCTGTTGGGGCG
 A347_3I TCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCTCCAGAAGATCCGTTGTTGACGGCGCTCCCTTTCTGTTGGGGCG

A347_30 GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3C GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3D GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3G GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3H GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3P GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGGCGGTGGGCAGTCTC
 A347_3K GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3L GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3a GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC
 A347_3I GGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTGCCTGAGAGGCCCGTGGACGGTGGGCAGTCTC

A347_30 GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3C GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3D GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3G GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3H GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3P GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3K GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3L GAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3a GAGAATACCGGAGGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A347_3I GAGAATACCGGAGAGAGACCGGTGATAACGACGATGCCATACCCCCGAT

7.3.2 Sequences from A810, Maraehako Bay

A810_4R ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A810_4O ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A810_4K ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A810_4G ATCATCGAAACTTCGAACGCATCTTGCCTCCCGGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA

7.3.4 Sequences from A868, Waterfall Reef, Leigh

A868_5N ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5R ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5G ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5L ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5B ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5D ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5H ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA
 A868_5K ATCATCGAAACTTCGAACGCATCTTGCGCTCCCGGATATGCCTGGGAGCATGCTTGTCTGGGGAGGAGGAGGCGAAAA

A868_5N ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5R ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5G ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGAGGAGTGGTGCG
 A868_5L ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5B ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5D ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5H ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG
 A868_5K ATCGCCACAGCTTCGGGTTTCGATCTCGATCTCGAGGCGGTGGAGCGGACTCTGAGTGTTCGGGGGGCAGTGGTGCG

A868_5N GTGTGTATTTTCACGTACATACTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAGAGAAACCGACCGTCCGGG
 A868_5R GTGTGTATTTTCACGTACATACTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAGAGAAACCGACCGTCCGGG
 A868_5G GTGTGTATTTTCACGTACATACTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAGAGAAACCGACCGTCCGGG
 A868_5L GTGTGTATTTTCACGTACATACTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAGAGAAACCGACCGTCCGGG
 A868_5B GTGTGTATTTTCACGTACATGCTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAG--AAACCGACCGTCCGGG
 A868_5D GTGTGTATTTTCACGTACATGCTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAG--AAACCGACCGTCCGGG
 A868_5H GTGTGTATTTTCACGTACATGCTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAG--AAACCGACCGTCCGGG
 A868_5K GTGTGTATTTTCACGTACATGCTGCCAGCTCGTCCCCTGAGTTCACCGAAGCCTAGAGAG--AAACCGACCGTCCGGG

A868_5N CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5R CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5G CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5L CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5B CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5D CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5H CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT
 A868_5K CATTTATCTCTTTGCGGCGCTTAAGGCAGGTTCACTTTGCGTCCTCCAGAAGATCCGTTGTTGACGGCGCCCCCTTT

A868_5N CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
 A868_5R CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
 A868_5G CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
 A868_5L CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
 A868_5B CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCTGGAGGCCCGTGGACGG
 A868_5D CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCCGGAGGCCCGTGGACGG
 A868_5H CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCCGGAGGCCCGTGGACGG
 A868_5K CGTGGGGCGGGGACACGACGGGTTGCTGGGGATGAGCGCGGGTGGCCTTGCTGCGTCGCGCCGGAGGCCCGTGGACGG

A868_5N TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5R TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5G TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5L TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5B TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5D TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5H TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT
 A868_5K TGGGCAGTCTCGAGAATACCGGAGAGAGACCGGTGATAATGACGATGCCATACCCCCGAT

[illegible]

Appendix 4. (Continued)

[illegible]

[illegible]

Appendix 4. (Continued)[illegible]

B0013	Cretorta	TGGCTGTTACCTCCATCTTTAATACTTCTTTAGCCTCTTCATTAGTAGAGAAGCGGGGCAGGTACTGGTTGGACTGTT
B601	Ccephalorni	TACCCCCCTTAAGTGGTATTCAAGCACACTCGGGACCATCGGTTGATTAGCTATTTTTAGCCTCCATTATCGGGT
B091	Ccephalorni	TACCCCCCTTAAGTGGTATTCAAGCACACTCGGGACCATCGGTTGATTAGCTATTTTTAGCCTCCATTATCGGGT
B327	Ccarpoconflu	TACCCGCCGTTAAgTGGTATCCAAGCACACTCGGGTCCCTCTGTCGATTAGCTATTTTTAGTCTTCATTATACAGGT
B162	Ccarpoconflu	TACCCGCCGTTAAgTGGTATCCAAGCACACTCGGGTCCCTCTGTCGATTAGCTATTTTTAGTCTTCATTATACAGGT
B174	Lquericifoli	TACCTCCCTTAAGTGGTATTCAAGCACACTCGGGACCATCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B318	Cmonilifera	TATCCTCCCTTAAGCGGTATTCAAGCCCATTcAGGTCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B092	Cmonilifera	TATCCTCCCTTAAGCGGTATTCAAGCCCATTcAGGTCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B095	Cmonilifera	TATCCTCCCTTAAGCGGTATTCAAGCCCATTcAGGTCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B102	Cmonilifera	TATCCTCCCTTAAGCGGTATTCAAGCCCATTcAGGTCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B120	Cmonilifera	TATCCTCCCTTAAGCGGTATTCAAGCCCATTcAGGTCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B093	Cmoniliform	TACCCCCCTTAAGCGGTATTCAAGCCCATTcAGGACCATCGGTCGATTAGCTATCTTTAGTCTTCATTATCTGGT
B163	Cmoniliform	TACCCCCCTTAAGCGGTATTCAAGCCCATTcAGGACCATCGGTCGATTAGCTATCTTTAGTCTTCATTATCTGGT
B094	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B100	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B101	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B099	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B123	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B124	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B166	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B108	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B617	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B430	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
A615	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B137	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B169	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B425	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B426	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B335	Ccongesta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B436	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
A154	Cscalaris	TATCCTCCCTTAAGTAGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
A080	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B602	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
A085	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B556	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
A196	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B412	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B121	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B109	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B170	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B415	Csiliquosa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B111	Cpolycystid	TATCCTCCCTTAAGTGAATTCAAGCCCATTcAGGGCCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B112	Cpolycystid	TATCCTCCCTTAAGTGAATTCAAGCCCATTcAGGGCCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B148	Cpolycystid	TATCCTCCCTTAAGTGAATTCAAGCCCATTcAGGGCCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B171	Cpolycystid	TATCCTCCCTTAAGTGAATTCAAGCCCATTcAGGGCCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
A168	Cplatylobi	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B140	Cplatylobi	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B104	Cplatylobi	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B116	Ccongesta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B115	Ccongesta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B152	Ccongesta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B455	Ccongesta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B129	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B131	Cretorta	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGACCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B161	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B626	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B146	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B146	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B128	Csubfarccus	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B127	Csubfarccus	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCATCGGTTGATTAGCTATTTTTAGTCTTCATTATCCGGT
B159	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT
B134	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B168	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B136	Csubfarcina	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTTGATTAGCTATTTTTAGTCTTCATTATCTGGT
B626	Cretroflexa	TATCCTCCCTTAAGTGGTATTCAAGCCCATTcAGGGCCGTCGGTCGATTAGCTATTTTTAGTCTTCATTATCTGGT

[illegible]

[illegible]

[illegible]

[illegible]

7.5 Appendix 5. Alignment of ITS sequences from *Cystophora*. Specimen names are truncated to FieldnumberSpecies.

```

B174Lquercifoli ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAG-----TC
B090Cmoniliform ACCCCCTCACCCGTCGGGGATGAGCAAGCGAGCGAGCGAGCGAGCGAGTGAAGCGAGCGA-----ACGAGCCACC
B114Cexpansa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----ATGGACCATC
B109Cretorta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B131Cretorta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B415Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B118Csiliquosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B143Csiliquosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B100Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B146Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B153Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B110Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B127Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B168Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B136Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B161Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B103Ctorulosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B125Ctorulosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B179Ctorulosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B165Ctorulosa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B171Cpolycystid ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B112Cpolycystid ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCACC
B148Cpolycystid ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCACC
B111Cpolycystid ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B102Cmonilifera ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B119Cmonilifera ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B318Cmonilifera ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----AAGGACCATC
B095CMONILIFERA ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCGA-----TAGGACCATC
B092Cmonilifera -----
B159Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----ATC
A154Cscalaris -----GGGGACGAGCGAGCGA-----TCGA-----ATC
A292Cscalaris ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
A469Cscalaris ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B178Cscalaris ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B321Csp ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B336Cdistenta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B337Cdistenta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B336Cscalaris ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----ATC
B104Cplatylobiu ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCTAACTAATATCGATAGAAAGGAAAGGACCATC
A168Cplatylobiu ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGAGCGAGCTAACTAATATCGATAGAAAGGAAAGGACCATC
B149Cycongesta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----TCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG---ATC
B152Ccongesta ACCCACTCACCCGTCGGGGATGAGCGAGCGAGCGAGCGAGCGA-----ATC
B101Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----ATC
A080Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----ATC
A196Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----ATC
A085Cretroflexa ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGAGCGA-----ATC
B335Ccongesta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----ATC
B166Csubfarcina ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGA-----TCGA-----AAGGAC----
B116Ccongesta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----TCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG---ATC
B115Ccongesta ACCCACTCACCCGTCGGGGACGAGCGAGCGAGCGAGCGA-----ATC
B121Cretroflexa -----GCGAGCGAGCGA-----TCGA-----ATC

B174Lquercifoli ACGC-----CCGCCGCCCG-----TACCCGGGTGCGATGAACGAG---AAGCGGGGGG
B090Cmoniliform ACGCTCCGCTCGCCCGGCCCGCCCGCCCGCCCGCCCGTTCGTTCCCGGGTGCGATGAACGAGG---AGCGGGGGG
B114Cexpansa ACGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B109Cretorta AGGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B131Cretorta AGGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B415Cretroflexa AGGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B118Csiliquosa AGGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B143Csiliquosa AGGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B100Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B146Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B153Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B110Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B127Csubfarcina ACGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B168Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B136Csubfarcina -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B161Cretroflexa -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B103Ctorulosa ACGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B125Ctorulosa -----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B179Ctorulosa ACGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B165Ctorulosa ACGCTCCGCTCGCCCGGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B171Cpolycystid ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B112Cpolycystid ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG

```

Appendix 5. (Continued)

B148Cpolycystid	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B111Cpolycystid	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B102Cmonilifera	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B119Cmonilifera	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B318Cmonilifera	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B095CMONILIFERA	ACGCTCCGTTTCGCTCG-----TTCCG-----
B092Cmonilifera	-----TCGCCCGT-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B159Csubfarcina	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A154Cscalaris	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A292Cscalaris	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A469Cscalaris	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B178Cscalaris	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B321Csp	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B336Cdistenta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B337Cdistenta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B336Cscalaris	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B104Cplatylobiu	ACGCTCCGCTCGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A168Cplatylobiu	ACGCTCCGCTCGCCCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B149Cycongesta	ACGCTCCGCTCGCCCG-----GTTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B152Ccongesta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B101Cretroflexa	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A080Cretroflexa	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A196Cretroflexa	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
A085Cretroflexa	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B335Ccongesta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B166Csubfarcina	-----GCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B116Ccongesta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B115Ccongesta	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B121Cretroflexa	ACGCTCCGCTCGCCCG-----TTCCCGGGTGCGATGAACGAGGAGAAGCGGGGGG
B174Lquercifoli	ACGGAGCTGACG-GCTTCGTCGCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGACTTGGGTGCTGCCGCGGTAC
B090Cmoniliform	AGGAAGCTTTCTTGCTCCGTCGCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGACTTGGGTGCTGCCGCGGTAC
B114Cexpansa	ACGGAGCTTTCTTGCTTCGTCGCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGGAGGCTTGGGTGCTGCCGCGGTAC
B109Cretorta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B131Cretorta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B415Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B118Csiliquosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B143Csiliquosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B100Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B146Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B153Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B110Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B127Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B168Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B136Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B161Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B103Ctorulosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B125Ctorulosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B179Ctorulosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B165Ctorulosa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B171Cpolycystid	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B112Cpolycystid	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B148Cpolycystid	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B111Cpolycystid	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B102Cmonilifera	GCGGAGCAATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B119Cmonilifera	GCGGAGCAATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B318Cmonilifera	GCGGAGCAATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B095CMONILIFERA	-----
B092Cmonilifera	GCGGAGCAATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B159Csubfarcina	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A154Cscalaris	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A292Cscalaris	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A469Cscalaris	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B178Cscalaris	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B321Csp	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B336Cdistenta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B337Cdistenta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B336Cscalaris	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B104Cplatylobiu	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A168Cplatylobiu	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B149Cycongesta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B152Ccongesta	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
B101Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A080Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A196Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC
A085Cretroflexa	GCGGAGCTATCTTGCTTCGTCCTCTCGTGAAAGGACCCCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCGCGGTAC

B135Ccongesta	GCGGAGCTATCTTGCTTCGTCGCCGCTCGTGAAAGGACCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCCGGTAC
B166Csubfarcina	GCGGAGCTATCTTGCTTCGTCGCCGCTCGTGAAAGGACCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCCGGTAC
B116Ccongesta	GCGGAGCTATCTTGCTTCGTCGCCGCTCGTGAAAGGACCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCCGGTAC
B115Ccongesta	GCGGAGCTATCTTGCTTCGTCGCCGCTCGTGAAAGGACCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCCGGTAC
B121Cretroflexa	GCGGAGCTATCTTGCTTCGTCGCCGCTCGTGAAAGGACCCCTCCCTTCAGTGTGGCAGGCTTGGGTGCTGCCCGGTAC
B174Lquercifoli	TGGAGTGGGGAGGCTCGGGAGCGCCGCAACCC - TCTCCAGTGGGGCCGCCTTGTGGGGCGGGGAGGGCCCGAGGT
B090Cmoniliform	TGGAGTGGGGAGGCTCGGGAGCGCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGTGGCGGAGAGGGCTCGAGGT
B114Cexpansa	TGGACGGGGGAGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B109Cretorta	TGGACGGGGGAGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B131Cretorta	TGGACGGGGGAGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B415Cretroflexa	TGGCGGGGGGAGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B118Csiliquosa	TGGACGGGGGAGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B143Csiliquosa	TGGACGGGGGAGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B100Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B146Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B153Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B110Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B127Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B168Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B136Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B161Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B103Ctorulosa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B125Ctorulosa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B179Ctorulosa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B165Ctorulosa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B171Cpolycystid	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B112Cpolycystid	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B148Cpolycystid	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B111Cpolycystid	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B102Cmonilifera	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B119Cmonilifera	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B138Cmonilifera	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B095Cmonilifera	-----ACCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B092Cmonilifera	TGGACGGGGGCGGCTCGGGAGCGCCCG - ACCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B159Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A154Cscalaris	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A292Cscalaris	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A469Cscalaris	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B178Cscalaris	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B321Csp	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B336Cdistenta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B337Cdistenta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B336Cscalaris	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B104Cplatylobiu	TGGACGGGGGCGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A168Cplatylobiu	TGGACGGGGGCGGCTCGGGAGCGCCCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B149Cpycongesta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B152Ccongesta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B101Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A080Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A196Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
A085Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B335Ccongesta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B166Csubfarcina	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B116Ccongesta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B115Ccongesta	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B121Cretroflexa	TGGACGGGGGCGGCTCGGGAGCGCCCGCAACCCCTCTCCGAGTGGGGCCGCCTTGGCGGGCGGAGAGGGCCCGAGGT
B174Lquercifoli	--CGTGCATCG--CACAATGCGCT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCG--
B090Cmoniliform	--CACTGC-----ACGCGGT-GCACTCCTCGCGAAGATCCAGTGGCTCCGTCGGTTCGATCGGTTCGGTTCGGTTC
B114Cexpansa	--CACTGCATCGCGCAGCAGCGGTTGGCACTCT??CGAAGATCTCAGTGGCTCGGTTCG--TCGGTTCGGTTCGGTTC
B109Cretorta	--CACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCGA-----
B131Cretorta	--CACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCGA-----
B415Cretroflexa	--CACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCGA-----
B118Csiliquosa	--CACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCGA-----
B143Csiliquosa	--CACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAGATCCAGTGGCTCGGTTCGA-----
B100Csubfarcina	CTCACTGCATCGCGCAGCAGCGGTT-GCACTCCTCGCGAAG

Appendix 5. (Continued)

B179Ctorulosa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CC
 B165Ctorulosa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B171Cpolycystid --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B112Cpolycystid --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B148Cpolycystid --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B111Cpolycystid --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B102Cmonilifera --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CCGGCC
 B119Cmonilifera --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CC
 B318Cmonilifera --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CC
 B095CMONILIFERA --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CCGGCC
 B092Cmonilifera --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----CCGGCC
 B159Csubfarcina --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A154Cscalaris --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A292Cscalaris --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A469Cscalaris --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B178Cscalaris --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B321Csp --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B336Cdistanta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B337Cdistanta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B336Cscalaris --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B104Cplatylobiu --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A168Cplatylobiu --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B149Cycongesta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B152Cycongesta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B101Cretroflexa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A080Cretroflexa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A196Cretroflexa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 A085Cretroflexa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B335Cycongesta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B166Csubfarcina --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B116Cycongesta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B115Cycongesta --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----
 B121Cretroflexa --CACTGCATCGCGCACGACGCGGT-GCACTCCTCGCGAAGATCCCAGTGGCTCGGTCCG-----

 B174Lquercifoli --CCGGTCGGTCGGCTGGAGCCAGGTTGTCTCTTCCCTCCGCAACCATCGTTATCTTTTGTGCTTGCCGCCGC---
 B090Cmoniliform --GGTCGGTCGGTCGGCTGGAGCCACAGTGTCTCTTCCCTTGCAACACCATTAATCTTTTCGTTGCTTGCCGCCGCCCTT
 B14Cexpansa --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATTAATTAATCTTTTCGTTGCAATGCCGCCGC---
 B109Cretorta --TCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATCATTATTTTCGTTGCTTGCCGCCGCCGC---
 B131Cretorta --TCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATCATTATTTTCGTTGCTTGCCGCCGCCGC---
 B415Cretroflexa --TCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATCATTATTTTCGTTGCTTGCCGCCGCCGC---
 B118Csiliquosa --TCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATCATTATTTTCGTTGCTTGCCGCCGCCGC---
 B143Csiliquosa --TCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTTGCAACATCATTATTTTCGTTGCTTGCCGCCGCCGC---
 B100Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B146Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B153Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B110Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B127Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B168Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B136Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B161Cretroflexa --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B103Ctorulosa --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B125Ctorulosa --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B179Ctorulosa --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B165Ctorulosa --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B171Cpolycystid --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B112Cpolycystid --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B148Cpolycystid --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B118Cpolycystid --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B111Cpolycystid --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B102Cmonilifera --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B119Cmonilifera --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B318Cmonilifera --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B095CMONILIFERA --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B092Cmonilifera --GGTCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B159Csubfarcina --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 A154Cscalaris --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 A292Cscalaris --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 A469Cscalaris --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B178Cscalaris --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B321Csp --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B336Cdistanta --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B337Cdistanta --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B336Cscalaris --CCGGTCGGTCGGCTCGAGCTGGAGTGTCTCTTCCCTCGCAACATTAATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B104Cplatylobiu --TCGGTCGGTCGGCTGACGCCGAGTGTCTCTTCCCTTGCAACATGATTAATTTTCGTTGCTTGCCGCCGCCGC---
 A168Cplatylobiu --TCGGTCGGTCGGCTGACGCCGAGTGTCTCTTCCCTTGCAACATGATTAATTTTCGTTGCTTGCCGCCGCCGC---
 B149Cycongesta --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---
 B152Cycongesta --CCGGTCGGTCGGCTGGAGCCGAGTGTCTCTTCCCTCGCAACATCATTACTTTTCGTTGCTTGCCGCCGCCGC---

Appendix 5. (Continued)

B101Cretroflexa	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
A080Cretroflexa	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
A196Cretroflexa	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
A085Cretroflexa	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B335Ccongesta	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B166Csubfarcina	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B116Ccongesta	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B115Ccongesta	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B121Cretroflexa	--CCGGTCGGTCGGCCTGGAGCCGGAGTGTCTCTTCCCCTCGCAACATCATTACTTTTCGTTGCCTGCCGCCGC---
B174Lquercifoli	GCCGTCGTCCTCCG-AAAAGGTGATGGTGGCGGTGCGAGCAGTGGCAGC---GATT---GTTGCGGGAGGGAGAGCGA
B090Cmoniliform	GCCATCGTCCCC-AAAAGATGATGGTGGTGGTCTGTA-----TTGGCTGTTGTGGGAGGGAGAGTGA
B114Cexpansa	GCCATCATCCCCAAAAGGTGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGGGA
B109Cretorta	GCCATCGTCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGTGA
B131Cretorta	GCCATCGTCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGTGA
B415Cretroflexa	GCCATCGTCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGTGA
B118Csiliquosa	GCCATCGTCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGTGA
B143Csiliquosa	GCCATCGTCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGTGGGAGGGAGAGTGA
B100Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B146Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B153Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B110Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B127Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B168Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B136Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B161Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B103Ctorulosa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B125Ctorulosa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B179Ctorulosa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B165Ctorulosa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B171Cpolycystid	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B112Cpolycystid	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B148Cpolycystid	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B111Cpolycystid	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B102Cmonilifera	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B119Cmonilifera	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B318Cmonilifera	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B095CMONILIFERA	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B092Cmonilifera	GCCATCGCCCC-AAAATATGATGGTGGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B159Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
A154Cscalaris	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
A292Cscalaris	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
A469Cscalaris	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B178Cscalaris	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B321Csp	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B336Cdistenta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B337Cdistenta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B336Cscalaris	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B104Cplatylobiu	ACCATCGTCCC-GAAATATGACGTTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGCGGGAGGGAGAGTGA
A168Cplatylobiu	ACCATCGTCCC-AAAATATGACGTTGGCGGTGCGGAGTGATGGAGA-----TTAGCTGTTGCGGGAGGGAGAGTGA
B149Cycongesta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B152Ccongesta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B101Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
A080Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
A196Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
A085Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B335Ccongesta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B166Csubfarcina	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGA---TTATTAGCTGTTGTGGGAGGGAGAGTGA
B116Ccongesta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B115Ccongesta	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B121Cretroflexa	GCCATCG-----ATGGTGTGCGGTGCGGAGTGATGGAGACTATTATTAGCTGTTGTGGGAGGGAGAGTGA
B174Lquercifoli	ATTAGAATGAATGGGCAAGTAACACGTCGCGGAGGT--CACGGCGTGGGGCCGGGCGCGGTGCGGATC-CATTA-----
B090Cmoniliform	ATTAGAATGAATGGGCAAGTAGCAGTCGTGGAGAT--CAGGGCGTGGGGCCGGGCGCGATCGGATC-----
B114Cexpansa	GTTAGAATGAATGGGCAAGTAGCAGTCGTGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCG-----
B109Cretorta	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCGGATC-TACT-----
B131Cretorta	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCGGATC-TACT-----
B415Cretroflexa	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCGGATC-TACT-----
B118Csiliquosa	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCGGATC-TACT-----
B143Csiliquosa	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGGCGCGATCGGATC-TACT-----
B100Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGTGTCATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----
B146Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGTGTCATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----
B153Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGTGTCATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----
B110Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGTGTCATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----
B127Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGT--CATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----
B168Csubfarcina	ATTAGAATGAATGGGCAAGTAGCTCGTCGGGAGGTGTCATGGCGTGGGGTCCGGTCCGATCGGATC-TACT-----

Appendix 5. (Continued)

B136Csubfarcina ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B161Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B103Ctorulosa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B125Ctorulosa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B179Ctorulosa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B165Ctorulosa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B171Cpolycystid ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATCCTACG-----
 B112Cpolycystid ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATCCTACG-----
 B148Cpolycystid ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATCCTACG-----
 B111Cpolycystid ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATCCTACG-----
 B102Cmonilifera ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACT-----
 B119Cmonilifera ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACT-----
 B318Cmonilifera ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACT-----
 B095CMONILIFERA ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACT-----
 B092Cmonilifera ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACT-----
 B159Csubfarcina ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 A154Cscalaris ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 A292Cscalaris ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 A469Cscalaris ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B178Cscalaris ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B321Csp ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B336Cdistenta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B337Cdistenta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B336Cscalaris ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGGCGCGATCGGATC-TACTACTGTT
 B104Cplatylobiu ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-----
 A168Cplatylobiu ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGGCGCGATCGGATC-----
 B149Cycongesta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B152Ccongesta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B101Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 A080Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 A196Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 A085Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B335Ccongesta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B166Csubfarcina ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGTGT CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B116Ccongesta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B115Ccongesta ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----
 B121Cretroflexa ATTAGAATGAATGGGCAAGTAGCTCGTCGCGGAGGT--CATGGCGTGGGGTCGGTCGCGATCGGATC-TACT-----

 B174Lquercifoli --TGTTGCCGTCGTGTGCGGCCTCAGTCGACCTTCTTGGCGGCGGCT---TGTCGGCGCGGTGGT---CGG-----
 B090Cmoniliform ---GTGTCCTGGTCGTGGGCGGCCTCAGTCGACCTTGGCGGCGGCGGTT---CGTCGGCGCGGTGGTGGGCGGTGCGCG
 B104Cexpansa ---TGTTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGGTGGTGGGTCGGGCGAG
 B109Cretorta GTTGTGTCTGGTCGTGGGCGGCCTTCTCGTTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGTGGTGGGTGGGCGAG
 B131Cretorta GTTGTGTCTGGTCGTGGGCGGCCTTCTCGTTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGTGGTGGGTGGGCGAG
 B415Cretroflexa GTTGTGTCTGGTCGTGGGCGGCCTTCTCGTTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGTGGTGGGTGGGCGAG
 B118Csiliquosa GTTGTGTCTGGTCGTGGGCGGCCTTCTCGTTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGTGGTGGGTGGGCGAG
 B143Csiliquosa GTTGTGTCTGGTCGTGGGCGGCCTTCTCGTTCGACCTTCCGCGGCGGCGGCG---CGTCGGCGCGTGGTGGGTGGGCGAG
 B100Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGTTTCAG
 B146Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B153Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B110Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B127Csubfarcina GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B168Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B136Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B161Cretroflexa GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B103Ctorulosa GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B125Ctorulosa GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B179Ctorulosa GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B165Ctorulosa GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B171Cpolycystid GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B112Cpolycystid GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B148Cpolycystid GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B111Cpolycystid GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B102Cmonilifera GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B119Cmonilifera GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B318Cmonilifera GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B095CMONILIFERA GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B092Cmonilifera GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGACCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B159Csubfarcina GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTTCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 A154Cscalaris GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 A292Cscalaris GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 A469Cscalaris GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B178Cscalaris GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B321Csp GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B336Cdistenta GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B337Cdistenta GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG
 B336Cscalaris GTTGTGTCTGGTCGTGGGCGGCCTCAGTCGTCCTTCCGCGGCGGCGGCGGCGCGCTGGCGCGTGGTGGGTGGGCGAG

Appendix 5. (Continued)

B104Cplatylobim -----TCTGGTCTGTTGGGCGGCCTTCGTGACCTTCGCGGCGGCGGCG---CGCTGGCGCGTTGGTGGGTGGGGCGAG
 A168Cplatylobiu -----TCTGGTCTGTTGGGCGGCCTTCGTGACCTTCGCGGCGGCGGCG---CGCTGGCGCGTTGGTGGGTGGGGCGAG
 B149Cycongesta GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B152Ccongesta GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B101Cretroflexa GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 A080Cretroflexa GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 A196Cretroflexa GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 A085Cretroflexa GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B335Ccongesta GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B166Csubfarcina GT-----GTCGTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B116Ccongesta GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B115Ccongesta GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG
 B121Cretroflexa GTTGTGTCTGGTCTGTTGGGCGGCCTCAGTCGTTCTTCGCGGCGGCGGCGGCGCGCTGGCGCGTTGGTGGGTGGGGCGAG

 B174Lquercifoli -----ACGTGCGT-----CCTGCTTCCT-----GCCGCGTGCAGCAAGTAATCATTACCAGTTT
 B090Cmoniliform GTACGCGTGTAGGCACGTGCGC-----CCAACCTCCTCC-----CCACGCGCGTGCAGGAGCAAAATTACCAGTTT
 B114Cexpansa GTACGCGGTACGTACGTACGTGCTCCCAACTCCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B109Cretorta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGGCRAAAATTACCAGTTT
 B131Cretorta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGGCAAAAATTACCAGTTT
 B415Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B118Csiliquosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B143Csiliquosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B100Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B146Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B153Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B110Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B127Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B168Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B136Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B161Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B103Ctorulosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B125Ctorulosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B179Ctorulosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B165Ctorulosa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B171Cpolycystid GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B112Cpolycystid GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B148Cpolycystid GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B111Cpolycystid GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B102Cmonilifera GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B119Cmonilifera GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B318Cmonilifera GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B095CMONILIFERA GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B092Cmonilifera GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B159Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 A154Cscalaris GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 A292Cscalaris GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 A469Cscalaris GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B178Cscalaris GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B321Csp GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B336Cdistenta GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B337Cdistenta GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B336Cscalaris GTAGGCGTGTACGTACGTGCGCGCGCCCAACTAACTCCCTCCTTTCCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B104Cplatylobiu GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TCTCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 A168Cplatylobiu GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCC-----TCTCCATCGCGTGCAGCGAGCAAAAATTACCAGTTT
 B149Cycongesta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B152Ccongesta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B101Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 A080Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 A196Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 A085Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B335Ccongesta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B166Csubfarcina GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B116Ccongesta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B115Ccongesta GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT
 B121Cretroflexa GTAGGCGTGTACGTACGTGCGC-----CCAACCTCCTCCTTC-----TTCCCATCGCGTGCAGCGAGCAAAATTACCAGTTT

 B174Lquercifoli ----CTCTGATCTGACGCGAGGGAGGGGGCGTGGCTGCCCGACGCGGTGGCTGCAGCCCC-TCC---CGGA----GC
 B090Cmoniliform GTTTCTCTGAACCTGACCCGAGGGAGGGA-----GGGGGTACAGGCCCC-TCCCCCGGA----TC
 B114Cexpansa TTTTCTCTGAACCTGACCCGAGGGA-----GGGGGTAC-GGCCCCCTCCTCCCGGA----GC
 B109Cretorta TTTTCTCTGGACTGACCCGAGGGA-----CGGGGGCTAC-GGCCCC-CCCTCCCGGA----GC
 B131Cretorta TTTTCTCTGGACTGACCCGAGGGA-----CGGGGGCTAC-GGCCCC-CCCTCCCGGA----GC
 B415Cretroflexa TTTTCTCTGGACTGACCCGAGGGA-----GGGGGTAC-GGCCCC-CCCTCCCGGA----GC
 B118Csiliquosa TTTTCTCTGGACTGACCCGAGGGA-----GGGGGTAC-GGCCCC-CCCTCCCGGA----GC
 B143Csiliquosa TTTTCTCTGGACTGACCCGAGGGA-----GGGGGTAC-GGCCCC-CCCTCCCGGA----GC
 B100Csubfarcina TTTTCTCTGGACTGACCCGAAGGA-----GGGGGTACAGGCCCC---TCCCGAGCGGTC
 B146Csubfarcina TTTTCTCTGGACTGACCCGAAGGA-----GGGGGTACAGGCCCC---TCCCGAGCGGTC

[illegible]

[illegible]

Appendix 5. (Continued)

A154Cscalaris	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
A292Cscalaris	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
A469Cscalaris	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B178Cscalaris	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B321Csp	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B336Cdistenta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B337Cdistenta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B336Cscalaris	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTTCGTCGT---TCCCCTGAGTTCACCC
B104Cplatylobiu	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGA-----GTACGTCGT---TCCCCTGAGTTCACCC
A168Cplatylobiu	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGA-----GTACGTCGT---TCCCCTGAGTTCACCC
B149Cycongesta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B152Ccongesta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B101Cretroflexa	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
A080Cretroflexa	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
A196Cretroflexa	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
A085Cretroflexa	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B335Ccongesta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B166Csubfarcina	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B116Ccongesta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B115Ccongesta	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B121Cretroflexa	AGGCGGTGGCGCGGACTCTGCGTGTTCGCGGGGACGAGACGACGACGAAAGTACGTCGT---TCCCCTGAGTTCACCC
B174Lquercifoli	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAAGGCCCCCCC
B090Cmoniliform	AAGGCTGGAGGGCAACCGACCGGCCGGCCCGGCCACGAGTTTCGCGCG---CTGGAGGCGTTTCACTTCA---TGCCCTTC
B114Cexpansa	AAGGCTGGAGGGCAACCGACCGGCCGGCCGACACGAGTCG-----AGGCGTTCGCTTCTAA---CGCCCTTC
B109Cretorta	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B131Cretorta	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B415Cretroflexa	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCGCGCGCGGAGGCGTTTCGCTTCAA---CGCCCTTC
B118Csiliquosa	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCGCGCGCGGAGGCGTTTCGCTTCAA---CGCCCTTC
B143Csiliquosa	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCGCGCGCGGAGGCGTTTCGCTTCAA---CGCCCTTC
B100Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B146Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B153Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B110Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B127Csubfarcina	AAGGCTGGAGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B168Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B136Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B161Cretroflexa	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B103Ctorulosa	AAGGCTGGAGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B125Ctorulosa	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B179Ctorulosa	AAGGCTGGAGAGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B165Ctorulosa	AAGGCTGGAGGAGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B171Cpolycystid	AAGGCTGGAGGGCAACCGACCGGCCGGCCT---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B112Cpolycystid	AAGGCTGGAGGGCAACCGACCGGCCGGCCT---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B148Cpolycystid	AAGGCTGGAGGGCAACCGACCGGCCGGCCT---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B111Cpolycystid	AAGGCTGGAGGGCAACCGACCGGCCGGCCT---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B102Cmonilifera	AAGGTTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B119Cmonilifera	AAGGTTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B318Cmonilifera	AAGGTTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B095CMONILIFERA	AAGGTTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B092Cmonilifera	AAGGTTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCGAGGCGTTTCGCTTCAA---CGCCCTTC
B159Csubfarcina	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
A154Cscalaris	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
A292Cscalaris	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
A469Cscalaris	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B178Cscalaris	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B321Csp	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B336Cdistenta	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B337Cdistenta	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B336Cscalaris	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B104Cplatylobiu	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCTTAGGCGTTTCGCTTCAA---CGCCCTTC
A168Cplatylobiu	AAGGCTGGAGGGCAACCGACCGGCCGGCCA---CGAGTTGCGGCG---CCTTAGGCGTTTCGCTTCAA---CGCCCTTC
B149Cycongesta	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B152Ccongesta	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
B101Cretroflexa	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
A080Cretroflexa	AAGGCTGGAGGGCAACCGACCGGCCGGCCG---CGAGTTGCGGCG---CCAGAGGCGTTTCGCTTCAA---CGCCCTTC
A196Cretroflexa	AAGGCT

Appendix 5. (Continued)

B114Cexpansa	GGACGAGCCGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGGGAAGCTTAGAC----
B109Cretorta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B131Cretorta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B415Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B118Csiliquosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B143Csiliquosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B100Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B146Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B153Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B110Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B127Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B168Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B136Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B161Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B103Ctorulosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B125Ctorulosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B179Ctorulosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B165Ctorulosa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B171Cpolycystid	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B112Cpolycystid	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B148Cpolycystid	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B111Cpolycystid	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B102Cmonilifera	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B119Cmonilifera	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B318Cmonilifera	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B095CMONILIFERA	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B092Cmonilifera	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B159Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A154Cscalaris	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A292Cscalaris	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A469Cscalaris	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B178Cscalaris	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B321Csp	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B336Cdistenta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B337Cdistenta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B336Cscalaris	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B104Cplatylobiu	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
A168Cplatylobiu	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAG-----
B149Cycongesta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B152Ccongesta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B101Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A080Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A196Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
A085Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B335Ccongesta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B166Csubfarcina	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B116Ccongesta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B115Ccongesta	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B121Cretroflexa	GGACGAGCTGTTGTTGACGGCGCCCCCTTCGCGGGGCGGGGACACGACGGGTCGCTCGGGGTGAT-----
B174Lquercifoli	GACAGCGTCGCTCGATCGCTC-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----CGAC-GA--
B090Cmoniliform	GACAGCGTCGCTCGATCGCTCGATCGCTCGATCGCTCGAGGGCCGGAAGGGAGGTAGGCTCGTG----CGACT----
B114Cexpansa	-----GAGG-CCGGAAGGGAGGTAGGCTCGTGA----GGACTG----
B109Cretorta	-----GAGGGCCGGAAGGGAGGTAACTCGTGA----GGACTG----
B131Cretorta	-----GAGGGCCGGAAGGGAGGTAACTCGTGA----GGACTG----
B415Cretroflexa	-----GAGGGCCGGAAGGGAGGTAACTCGTGA----GGACTG----
B118Csiliquosa	-----GAGGGCCGGAAGGGAGGTAACTCGTGA----GGACTG----
B143Csiliquosa	-----GAGGGCCGGAAGGGAGGTAACTCGTGA----GGACTG----
B100Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B146Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B153Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B110Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B127Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B168Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B136Csubfarcina	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B161Cretroflexa	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B103Ctorulosa	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B125Ctorulosa	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B179Ctorulosa	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B165Ctorulosa	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B171Cpolycystid	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B112Cpolycystid	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B148Cpolycystid	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B111Cpolycystid	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B102Cmonilifera	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT
B119Cmonilifera	-----GAGGGCCGGAAGGGAGGTAGGCTCGTGA----GGACTGACT

Appendix 5. (Continued)

B318Cmonilifera -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGA--
 B095CMONILIFERA -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGA--
 B092Cmonilifera -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGA--
 B159Csubfarcina -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A154Cscalaris -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A292Cscalaris -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A469Cscalaris -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B178Cscalaris -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B321Csp -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B336Cdistenta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B337Cdistenta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B336Cscalaris -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B104Cplatylobiu -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGT--
 A168Cplatylobiu -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGT--
 B149Cycongesta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B152Ccongesta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B101Cretroflexa -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A080Cretroflexa -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A196Cretroflexa -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 A085Cretroflexa -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B335Ccongesta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B166Csubfarcina -----GAGGGCCGGAAGGGAGGTAGGCTCGTGA---GGACTGACT--
 B116Ccongesta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B115Ccongesta -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--
 B121Cretroflexa -----GACGGCCGGAAGGGAGGTAGGCTCGTGAAAAAGGACTGA--

 B174Lquercifolia -CTT-----GAACGTA-----AAACAACG-ACGACGACGCCATACCCCCGAT
 B090Cmoniliformis ---TCTT-T-----AAGAAAGGAACG-ACG-ACGACGACGCCATACCCCCGAT
 B114Cexpansa -----AG-AACGAACGAACGAACGACGACGCCATACCCCCGAT
 B109Cretorta -----ACGAACGAACGACGACGCCATACCCCCGAT
 B131Cretorta -----ACGAACGAACGACGACGCCATACCCCCGAT
 B415Cretroflexa -----ACGAACGAACGAACGACGACGCCATACCCCCGAT
 B118Csiliquosa -----ACGAACGAACGAACGACGACGCCATACCCCCGAT
 B143Csiliquosa -----ACGAACGAACGAACGACGACGCCATACCCCCGAT
 B100Csubfarcina ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B146Csubfarcina ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B153Csubfarcina ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B110Csubfarcinata ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B127Csubfarcinata ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B168Csubfarcinata ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B136Csubfarcinata ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B161Cretroflexa ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B103Ctorulosa ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B125Ctorulosa ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B179Ctorulosa ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B165Ctorulosa ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B171Cpolycystidea -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B112Cpolycystidid -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B148Cpolycystidea -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B111Cpolycystidea -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B102Cmonilifera -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B119Cmonilifera -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B318Cmonilifera -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B095CMONILIFERA -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B092Cmonilifera -CTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B159Csubfarcinata -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACGACGACGCCATACCCCCGAT
 A154Cscalaris -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 A292Cscalaris -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 A469Cscalaris -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 B178Cscalaris -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 B336Cdistenta -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 B337Cdistenta -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 B336Cscalaris -CTTCTTCTTCTTAAGGAAGGA-GGGA-----AACGAACG-ACGACGACGCCATACCCCCGAT
 B104Cplatylobium -CTTCT-CTTCGT--CGGTCGAA-CGA---GCGAACGAACGAACGACGACGCCATACCCCCGAT
 A168Cplatylobium -CTTCT-CTTCGT--CGAACGAA-CGAACGAACGAACGAACGACGACGCCATACCCCCGAT
 B149Cycongesta -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B152Ccongesta -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B101Cretroflexa -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 A080Cretroflexa -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 A196Cretroflexa -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 A085Cretroflexa -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B335Ccongesta -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B166Csubfarcina ACTTCTT-----AACGAACGACGACGCCATACCCCCGAT
 B116Ccongesta -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B115Ccongesta -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT
 B121Cretroflexa -CTTCTTCTTCTTAAGGAAGGAAGGGA-----AACGAACAACGACGACGCCATACCCCCGAT

Appendix 6. (Continued)

B145Ccongesta	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B149Ccongesta	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B152Ccongesta	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B166Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B169Csubfarcina	TTTGTGAAAGTGGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B335Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAGAAGTTAAG
B412Csubfarcina	TTTGTGAAAGTGGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B416Csubfarcina	TTTGTGAAAGTGGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B425Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAGAAGTTAAG
B426Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAGAAGTTAAG
B430Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG
B436Cretroflexa	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAGAAGTTAAG
B455Ccongesta	TTTGTGAAAGTAGTTTACCTACAGTATAAGAGAAATTGTTTTTCACAAAAAGTTAAG

7.7 Appendix 7. Additional work.

During the period of work on this thesis I have also contributed to the following published papers:

- ZUCCARELLO, G. C., BUCHANAN, J. & WEST, J. A. (2006). Increased sampling for inferring phylogeographic patterns in *Bostrychia radicans*/*B. moritziana* (Rhodomeleaceae, Rhodophyta) in the eastern USA. *Journal of Phycology*, 42, 1349–1352.
- HODGE, F., BUCHANAN, J. & ZUCCARELLO, G. C. (2010). Hybridisation between the endemic brown algae *Carpophyllum maschalocarpum* and *Carpophyllum angustifolium* (Fucales): Genetic and morphological evidence. *Phycological Research*, 58, 239–247.
- JONES (KAIN), J., BUCHANAN, J., BOO, S. M. & LEE, K. M. (2010). *Colpomenia bullosa* crust masquerading as *Ralfsia verrucosa* in southeast Australia. *Phycologia*, 49, 617–627.
- ZUCCARELLO, G. C., BUCHANAN, J., WEST, J. A. & PEDROCHE, F. F. (2011). Genetic diversity of the mangrove associated alga *Bostrychia moritziana* (Ceramiales, Rhodophyta) from southern Central America. *Phycological Research*. doi: 10.1111/j.1440-1835.2010.00605.x

7.8 Appendix 8

Phylogeography of *Carpophyllum maschalocarpum*

Joe Buchanan & G. C. Zuccarello
School of Biological Sciences, Victoria University of Wellington

A highly variable 225 bp mitochondrial spacer region from 651 samples of the endemic fucoid alga *Carpophyllum maschalocarpum* was analysed by SSCP or sequencing. We identified 52 haplotypes, and found a further 15 haplotypes in the sister species *Carpophyllum angustifolium* and in hybrids between these species (Fig. 1).

1. Forty-one *C. maschalocarpum* haplotypes were restricted to a single population. Most other haplotypes were restricted to regions.
2. Haplotypic diversity was low in southern populations.
3. High diversity in three far north populations arises from hybridisation with *C. angustifolium*.
4. Connectivity was strong between East Coast and Northland populations and between the mainland and the Chatham Islands.
5. High diversity was also found in the Chatham Islands.

Fig 1. Statistical parsimony network inferred using TCS software (95% confidence limits). Area of circles is proportional to number of haplotypes sampled. Small black circles represent missing haplotypes.

1. Population differentiation and dispersal limitation
Population differentiation between populations with shared haplotypes was moderate to strong at distances over 50 kilometres (see Table 1 below), especially where there are habitat discontinuities. Despite large numbers of *C. maschalocarpum* thalli observed in drift (Kingsford 1992), gene flow is low. At several sites, all or most haplotypes sampled were restricted to that population.

Table 1. Pairwise F_{ST} estimates from mtDNA spacer data (below diagonal) and minimum round coast distances (above diagonal) for Bay of Plenty populations of *Carpophyllum maschalocarpum*.

	Sailors Grave	Maketu	Opape	Marahako Bay	Otaga
Sailors Grave	-	103 km	179 km	190 km	215 km
Maketu	0.54931**	-	87 km	117 km	151 km
Opape	0.46280**	0.84906**	-	48 km	86 km
Marahako Bay	0.42232**	0.68024**	N. S.	-	38 km
Otaga	0.12573*	0.26357**	0.11334*	0.12548**	-

Significance shown as: ** = $p < 0.01$ and * = $p < 0.05$ after sequential Bonferroni correction. N. S. = No significant difference.

2. Low southern diversity

Haplotypic diversity was low in southern populations, and a single haplotype was sampled in the South Island. The southern limit of *C. maschalocarpum* is correlated with the 15°C sea surface temperature isotherm for the warmest month (Fig. 3 below). Northward movement of this isotherm during the Last Glacial Maximum (LGM) would have restricted *C. maschalocarpum* populations to the northern half of the North Island. Stepping stone recolonisation of southern New Zealand following the LGM would reduce genetic diversity through repeated founder effects.

Fig. 3. Sea surface temperatures (warmest month). 15°C isotherms for present day and estimate for LGM (Barrows & Jiggins 2005).

Fig 2. Distribution of mtDNA haplotypes of *C. maschalocarpum*

5. Chatham Islands diversity

Seven haplotypes were sampled in Chatham Islands populations, four were endemic. The presence of endemic haplotypes suggests populations survived LGM cooling. Paleoclimate studies indicate northward movement of the sub-tropical convergence during the LGM might have been constrained by the Chatham Rise (Nelson *et al.* 2000), maintaining relatively warm water temperatures near the Chatham Islands, with cooler temperatures around mainland New Zealand. Four haplotype lineages were present on the Chatham Islands, indicating at least four separate colonisation events, over a 650 km habitat discontinuity.

3. Hybridisation in northern populations
Morphologically intermediate hybrid specimens were found in the Bay of Plenty and Northland populations where *C. maschalocarpum* and *C. angustifolium* are sympatric. All specimens from Hooper Point, North Cape and Wekaraa populations were hybrids, with either *C. angustifolium* and *C. maschalocarpum* haplotypes.

4. Northland and the East Coast connectivity
One haplotype was common in Northland (Cape Waiwiri and the Poor Knights) and the East Coast (Horoera and Kaiti Beach), but was absent from interjacent sites. The East Cape Current is a potential dispersal pathway connecting these populations. This current does not extend into the Bay of Plenty and the Hauraki Gulf. SAMOVA analysis (Dupanloup *et al.* 2002) grouped these populations together despite their spatial separation (Fig. 4 below).

Fig. 4. Grouping of populations inferred by SAMOVA. F_{ST} reached a maximum (0.8114) at K=14.

Conclusions

The population genetic structure of *C. maschalocarpum* reflects both contemporary processes (dispersal pathways, hybridisation) and historical events (climate change). Low connectivity between adjacent populations suggests dispersal limitation, but connections between Northland and the East Coast, and between mainland New Zealand and the Chatham Islands, show long distance dispersal by ocean currents.

References

- Barrows TC, Jiggins S (2005) Sea surface temperatures around the Australian margin and Indian Ocean during the last glacial maximum. *Quaternary Science Reviews* 24, 1017-1047.
- Dupanloup I, Schneider S, Excoffier L (2002) A simulated annealing approach to define the genetic structure of populations. *Molecular Ecology* 11, 2573-2581.
- Kingsford MJ (1992) Drift algae and small fish in coastal waters of northeastern New Zealand. *Marine Ecology Progress Series* 88, 41-55.
- Nelson CH, Handy RL, Neil HL, Hendy CH, Weaver JPS (2000) Last glacial jetting of cold waters through the Subtropical Convergence zone in the Southwest Pacific off eastern New Zealand, and some geological implications. *Palaeogeography, Palaeoclimatology, Palaeoecology* 156, 103-121.

Acknowledgements

Thanks to Wendy Nelson, Kate Neil, Tracy Farr, Erasmo Macaya, Nick Shears and Peter Martin for assistance with collections and the Department of Conservation for support.

Phycological Research

Phycological Research 2010; 58: 239–247

Hybridization between the endemic brown algae *Carpophyllum maschalocarpum* and *Carpophyllum angustifolium* (Fucales): Genetic and morphological evidence

Fiona J. Hodge, Joseph Buchanan and Giuseppe C. Zuccarello*

School of Biological Sciences, Victoria University of Wellington, Wellington 6140, New Zealand

SUMMARY

Hybridization is an important evolutionary process, which can have significant effects on biodiversity. While hybridization is well known in plants, less is known about the prevalence of hybridization in other kingdoms. Hybridization in the field has been confirmed in brown algae in a few cases, mainly in the northern hemisphere genus *Fucus*. Putative hybrids have been found in the New Zealand endemic species *Carpophyllum angustifolium* and *Carpophyllum maschalocarpum*. We used nuclear-encoded molecular data (ITS2) and morphometrics to confirm hybridization between *C. angustifolium* and *C. maschalocarpum*. Putative hybrid thalli were collected that had heterozygous ITS2, each copy corresponding to one of the parental species from that population. Morphological analysis also showed that the three classes (two parental species and hybrids) were easily distinguishable in these populations. It was found that the hybrids had an intermediate morphology to the parent species. Some individuals with *C. angustifolium* morphology had hybrid ITS2 ribotypes suggestive of backcrossing between the hybrids and *C. angustifolium*. Our data reveal another case of hybridization within the Fucales and suggests that further research on how these species remain separate is needed.

Key words: *Carpophyllum angustifolium*, *Carpophyllum maschalocarpum*, Fucales, hybrids, hybridization, ITS2, New Zealand, morphometrics, Phaeophyceae.

INTRODUCTION

Hybridization is an important evolutionary process that brings together diverged genetic lineages, through the crossing of two genetically distinct species. Hybridization challenges our species definitions and provides insight into speciation processes (Seehausen 2004). Often it also produces unique hybrid morphologies. Hybrids are generally expected to have intermediate

morphologies reflecting their mixed genotypes, but may also have morphologies similar to, novel from, or more extreme than the parental species (Coyer *et al.* 2002a; Seehausen 2004; Lihova *et al.* 2007). Identification of hybrids based on morphology alone can be difficult due to the large degree of intra-specific variation and phenotypic plasticity present in many taxa (Rieseberg & Ellstrand 1993; Scott & Hardy 1994). Furthermore, morphological intermediacy does not always indicate hybrids and may reflect patterns of plasticity or environmental variation (Mathieson *et al.* 1981).

Introgression, the backcrossing of hybrids with parental species, can result in a continuum of morphologies between the parental species (Albert *et al.* 1997). Introgression is of great evolutionary significance as it can result in the transfer of adaptations between species (Seehausen 2004). Other potential outcomes of introgression include the strengthening of reproductive barriers and further divergence of the parental species, or the development of new hybrid lineages (Grant *et al.* 2004; Seehausen 2004).

The Phaeophyceae, brown algae, are morphologically diverse. Individuals range from microscopic filaments to the more familiar large conspicuous fleshy seaweeds (Graham & Wilcox 2000). Even within species there is great morphological variation (Mathieson *et al.* 1981). Despite this variability putative hybrids, especially in the morphologically complex Fucales, have been observed in wild populations since the 1850s (references within Thuret 1854; Mathieson *et al.* 1981).

New Zealand has a distinct brown algal flora with high levels of endemism (Nelson 1994). The dominance of Fuclean algae in the shallow subtidal is considered one of the more unusual aspects of the flora (Schiel 1990). Fuclean algae in the Northern Hemisphere dominate the intertidal, and consequently have

*To whom correspondence should be addressed.

Email: joe.zuccarello@vuw.ac.nz

Communicating editor: C. Katsaros.

Received 17 December 2009; accepted 26 March 2010.

doi: 10.1111/j.1440-1835.2010.00583.x

been relatively well studied (Chapman 1995). The vast majority of research into hybridization in Phaeophyceae is on the Northern Hemisphere fucalean genus *Fucus* (Mathieson *et al.* 1981; Chapman 1995; Wallace *et al.* 2004; Engel *et al.* 2005; Perrin *et al.* 2005; Billard *et al.* 2005b; Coyer *et al.* 2006b). Research into hybridization has revealed hybrids with intermediate morphologies, and in some cases evidence of introgression has been reported (Coyer *et al.* 2002a; Wallace *et al.* 2004; Billard *et al.* 2005b; Mathieson *et al.* 2006). Little work has been done on hybridization in Fucales outside of *Fucus*, despite the well documented abundance, diversity and ecological importance of this order in the Southern Hemisphere (Clayton 1984).

In north-eastern New Zealand two species of Fucales (Sargassaceae) form dominant stands in the high subtidal: *Carpophyllum maschalocarpum* (Turner) Greville and *Carpophyllum angustifolium* J. Agardh (Dromgoole 1973; Schiel 1988; Schiel 1990). Based on intermediate morphotypes it has been hypothesized that hybridization is occurring between the two species (Dromgoole 1973).

The internal-transcribed-spacer (ITS) is part of the nuclear ribosomal cistron, and is an often used sequence in plant evolutionary studies (Feliner & Rosello 2007). ITS is multi-copy, and homogenization of the multiple ITS sequences can occur (Álvarez & Wendel 2003). Coyer *et al.* (2002b) showed that artificially produced F_1 hybrids of two *Fucus* species with distinct ITS ribotypes were all heterozygous in ITS. Introgressed and F_2 individuals will not necessarily have heterozygous ITS ribotypes (Álvarez & Wendel 2003). Combining ITS and morphometric data can allow non-subjective identification of hybrid and parental individuals, and help determine the presence of introgression and F_2 hybrids within a population.

The present study using combined ITS2 identification and morphometric analysis confirmed hybridization between the non-*Fucus* fucalean algae, *C. angustifolium* and *C. maschalocarpum*. The expansion of the study of hybridization from the one northern hemisphere model, to a related but endemic Fucalean alga, will provide additional understanding of the importance of hybridization in this class.

MATERIALS AND METHODS

Sampling

Samples were collected from two sites on the East Cape of the North Island of New Zealand, where putative *C. angustifolium* $\times$ *C. maschalocarpum* hybrids were known to occur (Fig. 1). The two sites are both within the greater Whanarua Bay, which faces north-west and contains a number of smaller sandy bays separated by large rocky islands. The first site is located at the

Fig. 1. The location of sampling (a) on the East Cape of the North Island of New Zealand, and (b) the two sites in the greater Whanarua Bay: Maraehako Bay (MH) and Uncles Bay (UB) (Google Earth 5.0).

south-eastern end of Maraehako Bay ($S37^{\circ}40.4'$ $E177^{\circ}47.8'$). The second site was located at Uncles Bay ($S37^{\circ}40.0'$ $E177^{\circ}48.0'$), which is the adjacent bay to the north-east. The two sites are approximately 500 m apart, and are separated by Motu Kaimeanui Island and two deep inlets. Sampling was conducted during the 2007–2008 summer.

Transect lines were laid along rock walls in the high subtidal, and quadrat locations were selected randomly from the top 0.5 m of *Carpophyllum* depth distribution. Quadrats were rejected if the rock face was not vertical, less than three individuals were present, or if the quadrat was less than 0.5 m from a previously sampled quadrat. All *Carpophyllum* individuals with holdfasts inside the 20 cm by 20 cm quadrats were collected. Sixty-nine quadrats were sampled in the two sites, with 280 *C. angustifolium*, 152 *C. maschalocarpum* and 32 hybrid adults collected in total (see Fig. 2 for typical morphologies). Only a subset that was characterized molecularly was used in this study ($n = 80$). Collected material was frozen at -20°C within a week of collection, and thawed in seawater prior to morphometric analysis.

Representative herbarium specimens of *C. angustifolium*, *C. maschalocarpum* and (putative) hybrids collected in East Cape during this study have been deposited in the Museum of New Zealand – Te Papa Tongarewa herbarium under the WELT numbers A029625–29, A029635–38 and A029630–34, respectively.

Morphometrics

Adults longer than 20 cm were morphologically identified, cleared of epiphytes, and measured. Only those individuals with complete haptera were used in mor-

Fig. 2. Typical morphology of (a) *Carpophyllum maschalocarpum*, (b) putative *Carpophyllum angustifolium* × *C. maschalocarpum* hybrids, and (c) *C. angustifolium*. All at the same scale, scale bar in (a) shows 50 mm.

Fig. 3. The location of the measurements taken from each individual are shown with arrows on a stylized axis. Stipe width, lamina length and width are illustrated at 20 cm from the apex.

phometrics so as to avoid pseudo-replication (i.e. the doubling up of measurements from the same individual). Individuals were measured using digital callipers and a meter rule. A small section of lamina was removed from each individual after processing and placed into a labeled tube with silica gel for genetic analysis. The total length of the individual, presence of vesicles longer than 8 mm, and the presence of secondary branches longer than 20 cm were also recorded.

The morphometric characteristics selected for measurement were stipe width and thickness; and lamina length, width and thickness (Fig. 3). The present study focused on the stipe and lamina characteristics of the primary axis (defined as the dominant axis, which arises directly from the holdfast). The stipe and lamina measurements were taken from approximately 5, 10, 15, and 20 cm from the tip of the axis. Lamina length was measured from the edge of the stipe where the lamina

forms to its tip. Lamina and stipe width were measured at the widest point, perpendicular to an imagined mid-line. Stipe and lamina thickness was measured at the thickest point. In some places herbivory had altered the morphology of the stipe and lamina; these could be detected by sudden and distinct changes in lamina or stipe outline. Measurements of stipe and lamina that were affected by herbivory were excluded, and if possible replaced by measurements from the nearest intact equivalent within 3 cm.

Digital images for the *C. angustifolium* (#2572, Botanical Museum Lund) and *C. maschalocarpum* (#000562635, The Natural History Museum, London Department of Botany) type specimens were obtained and measurements of lamina length, lamina width and stipe width were taken where possible.

Genetic analysis

Validity of the morphological identification for the hybrids and parental species were checked by genetically identifying 80 individuals. Individuals with unusual morphological characteristics were also checked genetically.

DNA was extracted from approximately 3 mg of dry tissue using a cetyltrimethylammonium bromide (CTAB) extraction protocol (Zuccarello & Lokhorst 2005). The ITS2 was amplified using a PTC-100 (Programmable Thermal Controller, MJ Research Inc. Waltham, MA, USA) polymerase chain reaction (PCR) machine, and the primers KG4 (CTTTTCCTCCGCT TAGTTATATG) and KP5 (ACAACGATGAAGAACGCAG) (Lane *et al.* 2006). The PCR conditions were an initial 94°C for 4 min, followed by five cycles of 1 min at 94°C, 1 min 59°C, 1 min 72°C with the annealing temperature decreased by 1°C each cycle. Then 30

cycles of 45 s 94°C, 45 s 53°C, 45 s 72°C, followed by 10 min at 72°C. Sequencing was done commercially (Macrogen Inc., Seoul, Korea). The chromatograms were visually examined and individuals identified as either a parental species or a heterozygous hybrid (where double peaks were present).

Cloning

One specimen from Maraehako Bay that was found to be heterozygous for ITS2 in direct sequencing was selected for cloning to determine whether both parental ribotypes could be obtained. The complete ITS1-5.8S-ITS2 cistron was amplified using KP1 and KG4 primers (Lane *et al.* 2006). PCR product was cloned using the pGEM-T Easy vector kit (Promega Corporation, Madison, WI, USA) following the manufacturer's protocols. Transformed colonies were sub-cultured and plasmid preparations carried out by alkali lysis (Sambrook *et al.* 1989). Inserts were sequenced commercially using pUC M13 forward and reverse primers.

Morphometric data analysis

The morphometric data contained a large number of missing values. Scatterplots showed no difference between the character traits measurements at 5, 10, 15 and 20 cm from the apex, so the measurements from each individual were averaged. There were some missing data for lamina characteristics after this averaging, as many lamina were excluded from measurement due to herbivory damage. The missing values were not randomly distributed, and *C. maschalocarpum* averages were based on fewer data points than the others due to greater levels of herbivory and exposure damage. Randomly removing data points to an equivalent level in *C. angustifolium* and hybrid data made little or no change to overall group averages (0.002% to 0.03% change). This suggests no bias in accuracy due to variation in the number of data points averaged between the species.

Cluster analysis finds natural groupings or clusters within the data (Fielding 2007). These were conducted with genetically confirmed data and type specimens using the software NTSYSpc version 2.11 software (Rohlf 2000). The unweighted pair-groups method average (UPGMA) clustering method was used; this is a common and accepted method (Fielding 2007) for this type of data.

Boxplots were used to identify morphological traits that showed difference. These characters were incorporated into a distance matrix using Gower's Index in Le Prodigiciel R 4.0 software (Casgrain & Legendre 1999). This index was used as it can cope with missing values (Montanari & Mignani 1994; Podani 1999). It has also been used in other studies of macroalgae morphology (Fowler-Walker *et al.* 2006). Cophenetic values were

calculated to test how well the cluster tree represents the data (Rohlf & Sokal 1981; Rohlf 2000).

ANOVA was used to compare the morphological groups found in the cluster analysis. Non-normal data were log transformed prior to analysis with ANOVA. Log transformed data that still did not meet ANOVA assumptions was analyzed using the non-parametric Kruskal–Wallis test. χ^2 tests were used to test for associations between wave exposure zones and the presence of vesicles and branches. Significance was determined at the 0.05 probability level.

RESULTS

ITS2 sequences and morphological identification

Only two ITS2 ribotypes were found, corresponding to the two morphologically identified parent species. The ribotypes differed by three different point mutations and one base pair indel (GenBank Numbers GU992939–GU992940). The point mutations appeared as double peaks in chromatograms from heterozygous individuals, and the indel appeared as a change from predominantly single peaks to all superimposed double peaks. This change occurred at the same base position in both forward and reverse sequences. All samples identified *a priori* as hybrids had heterozygous sequences (i.e. sequences of both *C. maschalocarpum* and *C. angustifolium*).

Twelve ITS2 sequences were obtained from cloned DNA from the heterozygous specimen. Multiple copies of the two ribotypes were found, corresponding to those expected from direct sequencing. Some sequences showed single base changes (autapomorphies) that were not previously detected by direct sequencing and were only present in a single clone from each specimen. We attribute these to copy errors during PCR with *Taq* polymerase.

Morphological identifications matched with genetic sequences in 93.8% of the individuals ($n = 80$). All *C. maschalocarpum* ($n = 25$) and hybrid ($n = 27$) morphological assignments were congruent with ITS2 sequences. 82.1% of the samples ($n = 28$) morphologically identified as *C. angustifolium* had matching *C. angustifolium* ITS2 sequences. The remaining 17.9% were five individuals with hybrid sequences; these will be described from here on as 'putative backcrosses'.

Cluster analysis

Stipe width, lamina length and lamina width were used in the cluster analyses. The cluster analysis tree of the genetically identified individuals ($n = 80$) and the type specimens is shown (Fig. 4). This had a cophenetic correlation of 0.84, indicating a 'good fit' (Rohlf 2000).

Fig. 4. Cluster analysis showing the morphology of a random sub-sample of genetically identified individuals, and the type specimens for the parental species (shown by asterisks). Individuals cluster into four morphological groups as labeled I to IV. Colors represent the identity of the individuals (see legend). Putative backcrosses are those morphologically identified as *Carpophyllum angustifolium* but with hybrid ITS2 sequences, as shown with arrows. The cophenetic value of this tree is $r = 0.84486$.

A cluster analysis of all individuals collected showed the groupings as seen for the genetically identified samples (Hodge 2009).

Genetically identified individuals clustered into four main morphological groups. Group I contained all of the genetically confirmed *C. angustifolium* ($n = 23$), two hybrids, and the putative backcrosses ($n = 5$). Group II included all of the other hybrids ($n = 25$), and one individual identified as *C. maschalocarpum*. Closer examination of this *C. maschalocarpum* individual reveals it was only just long enough to be considered 'adult' (>20 cm long) and therefore included in this study. Groups III and IV were composed of *C. maschalocarpum* only (III, $n = 19$; IV, $n = 6$). Hereafter Group I will be referred to as *C. angustifolium*, Group II as the hybrids, Groups III and IV combined as *C. maschalocarpum*. A major subdivision between the morphology of *C. angustifolium* and hybrid individuals, and the *C. maschalocarpum* individuals, was revealed in the cluster analysis. The hybrid cluster had a higher similarity coefficient with the *C. angustifolium* cluster (0.75) than with the *C. maschalocarpum* clusters (0.49).

A cluster analysis of all individuals collected, i.e. including specimens only morphologically identified, had a cophenetic correlation of greater than 0.90, indicating the cluster tree is a 'very good fit' of the data (Rohlf 2000). This cluster analysis showed the same group structure and membership of genetically identified individuals as the cluster tree of only genetically identified individuals (Hodge 2009). Consequently morphological comparisons were done between groups using data from both morphologically and genetically identified individuals.

Comparing morphologies

Comparisons of the morphologies of the *C. angustifolium* group (I), the hybrid group (II) and the pooled *C. maschalocarpum* groups (III and IV) found a range of differences between them (Table 1). Log stipe width ($P < 0.001$), lamina length ($P < 0.001$), and lamina width ($P < 0.001$) were different between the three groups (all pairwise Tukey Tests were $P < 0.05$) (Fig. 5). The pooled *C. maschalocarpum* group had a wider stipe, and a shorter and wider lamina than the *C. angustifolium* group. The hybrid group exhibited traits intermediate to the parents for each of these traits. There was no overlap in log stipe width for the three groups (Fig. 5a).

Stipe thickness was also different between the three groups ($P < 0.001$) with median stipe thickness being thickest for the *C. angustifolium* group (I), and thinnest for the hybrid group (II). Lamina thickness was also found to be non-uniform ($P < 0.01$). The *C. angustifolium* group (I) had significantly thicker lamina than the hybrids (II), and the pooled *C. maschalocarpum* group (III and IV) (Tukey tests: $P < 0.05$). The lamina thickness of the hybrid group (II) and the pooled *C. maschalocarpum* group (III and IV) were not significantly different (Tukey test: $P = 0.952$).

There were significant differences in total length, stipe width, stipe thickness, presence of vesicles between the two groups (III and IV) of *C. maschalocarpum* (Table 2). Group IV was generally longer in total length, had a wider and thicker stipe, and had a greater proportion of individuals with vesicles present than Group III. There were also non-significant trends ($0.05 < P < 0.10$) of Group IV having wider and thicker lamina, and more frequently having secondary branches. There was no difference between the lamina length of the two groups. There was no difference in the distributions between the two *C. maschalocarpum* groups between sites ($X^2 = 3.146$, d.f. = 1, $n = 146$, $P = 0.076$) or exposure zones.

DISCUSSION

Our molecular data confirms that *Carpophyllum angustifolium* and *C. maschalocarpum* hybridize in popula-

Table 1. Relative morphological differences between the three groups (tested with ANOVA and Kruskal–Wallis tests)

Character trait	Group I	Group II	Group III and IV	Statistics
Stipe width	Narrow	Intermediate	Wide	$F = 4843.291$, d.f. = 2, $n = 463$, $P < 0.001$
Lamina length	Long	Intermediate	Short	$F = 43.275$, d.f. = 2, $n = 240$, $P < 0.001$
Lamina width	Narrow	Intermediate	Wide	$F = 247.230$, d.f. = 2, $n = 240$, $P < 0.001$
Stipe thickness	Thick	Thin	Intermediate	$\chi^2 = 23.134$, d.f. = 2, $n = 463$, $P < 0.001$
Lamina thickness	Thick	Thin	Thin	$F = 21.22$, d.f. = 2, $n = 240$, $P < 0.001$

Fig. 5. The three groups had significantly different (a) log stipe widths, (b) lamina lengths and (c) lamina widths. The boxplots show the spread of data (line in box = mean, box = 75% of values measured, lines = 95% of values) with outliers marked by circles and extreme outliers marked with asterisks.

tions from East Cape. *Carpophyllum angustifolium*, *C. maschalocarpum* and their hybrids have distinct and distinguishable morphologies. Identifiable hybrid and parental morphotypes have been found within many *Fucus* hybrid species complexes; however, morphologically ambiguous individuals are often present (Scott & Hardy 1994; Coyer *et al.* 2002a; Kucera & Saunders 2008). No other detailed morphometric analysis of randomly sampled and molecularly identified individu-

als has been done in a Fuclean hybrid system. Subsequently the range and degree of overlap of parental and hybrid morphologies are unknown in other Fuclean hybrid systems.

Stipe width alone can distinguish between *C. angustifolium*, *C. maschalocarpum* and their hybrids, and provides a useful taxonomic character for field identification. Stipe width meets three requirements described for a useful macroalgal taxonomic character in macroalgae; it is constant, easy to observe and non-destructive (Mathieson *et al.* 1981). Stipe width of the parent species and hybrids was easy to measure and rarely affected by herbivory or exposure damage (*pers. obs.*). An investigation into the spatial variability of *F. spiralis* and *F. vesiculosus* (Coleman & Muhlin 2008) found stipe width is also a valuable character for identification of *Fucus* species.

Carpophyllum angustifolium × *C. maschalocarpum* hybrids generally had an intermediate morphology to the parent species. Hybrids had intermediate stipe width, lamina length, lamina width and vesicle presence. However, the thickness of hybrid lamina was novel, being thinner than both parents. Furthermore the proportion of hybrid individuals with secondary branching was indistinguishable from *C. maschalocarpum*. Intermediate hybrid morphologies are usually found in hybrid systems although not necessarily in all cases (Rieseberg & Ellstrand 1993; Seehausen 2004; Coyer *et al.* 2006b). Intermediate morphology is used to identify putative hybrid individuals for further investigation in *Fucus* (Scott & Hardy 1994; Coyer *et al.* 2002a). Despite this there are examples of morphological traits in hybrids being aligned to one of the parental species. For example, hybridization between the dioecious *Fucus vesiculosus* and the hermaphroditic *Fucus spiralis* results in hybrids with parental sexual phenotypes rather than intermediate sexual allocation (Billard *et al.* 2005b). In salt marsh habitats *F. vesiculosus* × *F. spiralis* hybrids are morphologically indistinguishable from polyploid *F. vesiculosus* individuals (Coyer *et al.* 2006b).

The presence of hybrid genotypes with *C. angustifolium* morphotypes suggests introgression might be occurring. Hybrid ITS genotypes with parental morphology were also found in the *F. vesiculosus* × *F. spiralis* species complex, and were also cautiously interpreted

Table 2. Relative morphological differences between the *Carpophyllum maschalocarpum* clusters III and IV

Significance	Morphological trait	Group III	Group IV	Statistics
Statistically significant at 5% level	Total length	Shorter	Longer	$F = 6.803$, d.f. = 1, $n = 146$, $P = 0.010$
	Stipe width	Narrower	Wider	$F = 250.906$, d.f. = 1, $n = 146$, $P < 0.001$
	Stipe thickness	Thinner	Thicker	$F = 14.188$, d.f. = 1, $n = 146$, $P < 0.001$
	Proportion with vesicles	Low	High	$\chi^2 = 11.026$, d.f. = 1, $n = 146$, $P = 0.001$
Trends in the data (significant at the 10% level)	Lamina width	Narrower	Wider	$F = 2.885$, d.f. = 1, $n = 49$, $P = 0.096$
	Lamina thickness	Thinner	Thicker	$F = 3.837$, d.f. = 1, $n = 49$, $P = 0.056$
	Proportion with secondary axes present	Low	High	$\chi^2 = 3.161$, d.f. = 1, $n = 146$, $P = 0.075$
Not significant and no trends.	Lamina length	Same	Same	$F = 1.517$, d.f. = 1, $n = 49$, $P = 0.224$

as indicating introgression (Kucera & Saunders 2008). Introgression is an important evolutionary process that has the potential to integrate genetic material from one species into another (Coyer *et al.* 2007). Introgression has been identified in *Fucus* hybrid complexes, and is considered a significant factor in the evolution of *Fucus* lineages (Wallace *et al.* 2004; Coyer *et al.* 2006a, 2007). Further investigation with nuclear markers lacking the unusual properties of ITS (Álvarez & Wendel 2003) are needed to confirm introgression and the degree of introgression.

Fucus species seem to have maintained their genetic integrity despite extensive hybridization. For example, hybrids made up 13% of *Fucus serratus* and *Fucus evanescens* in one population; however, nuclear DNA introgression was only 1.5% in the two species, and both species maintain different distributions and morphologies (Coyer *et al.* 2002a, 2007). Coyer *et al.* (2006a) suggested different parental mating systems might act as a barrier to extensive introgression, and protect the parental species integrity. Hybridization in *Fucus* appears to occur between dioecious (having separate male and female individuals) and hermaphroditic species (e.g. Wallace *et al.* 2004; Billard *et al.* 2005a; Mathieson *et al.* 2006; Coyer *et al.* 2007). All members of the genus *Carpophyllum* are dioecious (Lindauer *et al.* 1961). Extensive introgression, and the loss of parental species genetic integrity, could be more likely in *Carpophyllum* hybrid systems if Coyer *et al.* (2006a) hypothesis is correct.

The absence of individuals with *C. maschalocarpum* morphology and hybrid genotypes suggests introgression may be asymmetrical, with backcrossing only between hybrids and *C. angustifolium*. Asymmetries in angiosperm reproduction are thought to be common in a broad range of taxa (Tiffin *et al.* 2000). Asymmetrical hybridization and introgression has been found in *Fucus* spp. (Coyer *et al.* 2002a, 2007). Determining the symmetry of introgression is important for predicting its implications on parental species integrity (Coyer *et al.* 2007).

The morphometric analysis distinguished variation within *C. maschalocarpum*, which could be related to

demography. The differences between the two *C. maschalocarpum* clusters (i.e. total length, stipe width, stipe thickness, vesicles) were characteristics thought to be associated with maturity (Dromgoole 1965). *Carpophyllum maschalocarpum* has pulsed peak recruitment, and has been shown to recruit in cohorts after bare rock is opened up by disturbance (Schiel 1980, 1988).

This work on *Carpophyllum* combining morphometric data with ITS2 identification found generally distinct and distinguishable parental and hybrid morphologies. Further investigation needs to be done using microsatellites and maternally inherited DNA in conjunction with morphology to confirm and determine the extent and symmetry of introgression, and identify backcrosses, F_2 and later hybrids.

ACKNOWLEDGMENTS

We thank Carlos Lehnebach for statistical help, and Tracey Taylor, Jen Van der Veur, and Aaron Packard for field assistance.

REFERENCES

- Albert, M. E., Dantonio, C. M. and Schierenbeck, K. A. 1997. Hybridization and introgression in *Carpobrotus* spp. (Aizoaceae) in California. I. Morphological evidence. *Am. J. Bot.* **84**: 896–904.
- Álvarez, I. and Wendel, J. F. 2003. Ribosomal ITS sequences and plant phylogenetic inference. *Mol. Phylogenet. Evol.* **29**: 417–34.
- Billard, E., Daguin, C., Pearson, G., Serrão, E., Engel, C. and Valero, M. 2005a. Genetic isolation between three closely related taxa: *Fucus vesiculosus*, *F. spiralis*, and *F. ceranoides* (Phaeophyceae). *J. Phycol.* **41**: 900–5.
- Billard, E., Serrão, E. A., Pearson, G. A., Engel, C. R., Destombe, C. and Valero, M. 2005b. Analysis of sexual phenotype and prezygotic fertility in natural populations of *Fucus spiralis*, *F. vesiculosus* (Fucaceae, Phaeophyceae) and their putative hybrids. *Eur. J. Phycol.* **40**: 397–407.
- Casgrain, P. and Legendre, P. 1999. *The R Package*. version 4.0.

- Chapman, A. R. O. 1995. Functional ecology of fucoid algae – 23 years of progress. *Phycologia* **34**: 1–32.
- Clayton, M. N. 1984. Evolution of the Phaeophyta with particular reference to the Fucales. In Round, F. E. and Chapman, D. J. (Eds) *Progress in Phycological Research*, Vol. 3. Biopress Ltd, Bristol, pp. 11–46.
- Coleman, M. A. and Muhlin, J. F. 2008. Patterns of spatial variability in the morphology of sympatric fucoids. *North-east Nat.* **15**: 111–22.
- Coyer, J. A., Peters, A. F., Hoarau, G., Stam, W. T. and Olsen, J. L. 2002a. Hybridization of the marine seaweeds, *Fucus serratus* and *Fucus evanescens* (Heterokontophyta: Phaeophyceae) in a 100-year-old zone of secondary contact. *Proc. R. Soc. Biol. Sci. Ser. B* **269**: 1829–34.
- Coyer, J. A., Peters, A. F., Hoarau, G., Stam, W. T. and Olsen, J. L. 2002b. Inheritance patterns of ITS1, chloroplasts and mitochondria in artificial hybrids of the seaweeds *Fucus serratus* and *F. evanescens* (Phaeophyceae). *Eur. J. Phycol.* **37**: 173–8.
- Coyer, J. A., Hoarau, G., Oudot-Le Secq, M.-P., Stam, W. T. and Olsen, J. L. 2006a. A mtDNA-based phylogeny of the brown algal genus *Fucus* (Heterokontophyta; Phaeophyta). *Mol. Phylogenet. Evol.* **39**: 209–22.
- Coyer, J. A., Hoarau, G., Pearson, G. A., Serrão, E. A., Stam, W. T. and Olsen, J. L. 2006b. Convergent adaptation to a marginal habitat by homoploid hybrids and polyploid ecads in the seaweed genus *Fucus*. *Biol. Lett.* **2**: 405–8.
- Coyer, J. A., Hoarau, G., Stam, W. T. and Olsen, J. L. 2007. Hybridization and introgression in a mixed population of the intertidal seaweeds *Fucus evanescens* and *F. serratus*. *J. Evol. Biol.* **20**: 2322–33.
- Dromgoole, F. I. 1965. Contribution to the ecology of *Carpophyllum maschalocarpum* (Turn.) Grev. MSc thesis, University of Auckland.
- Dromgoole, F. I. 1973. A contribution to the biology of the genus *Carpophyllum* Grev. PhD thesis, University of Auckland.
- Engel, C. R., Daguin, C. and Serrão, E. 2005. Genetic entities and mating systems in hermaphroditic *Fucus spiralis* and its close dioecious relative *F. vesiculosus* (Fucaceae, Phaeophyceae). *Mol. Ecol.* **14**: 2033–46.
- Feliner, G. N. and Rossello, J. A. 2007. Better the devil you know? Guidelines for insightful utilization of nrDNA ITS in species-level evolutionary studies in plants. *Mol. Phylogenet. Evol.* **44**: 911–9.
- Fielding, A. H. 2007. *Cluster and Classification Techniques for the Biosciences*. Cambridge University Press, Cambridge.
- Fowler-Walker, M. J., Wernberg, T. and Connell, S. D. 2006. Differences in kelp morphology between wave sheltered and exposed localities: morphologically plastic or fixed traits? *Mar. Biol.* **148**: 755–67.
- Graham, L. E. and Wilcox, L. W. 2000. *Algae*. Prentice Hall, Upper Saddle River, NJ.
- Grant, P. R., Grant, B. R., Markert, J. A., Keller, L. F. and Petren, K. 2004. Convergent evolution of Darwin's finches caused by introgressive hybridization and selection. *Evolution* **58**: 1588–99.
- Hodge, F. J. 2009. Hybridisation in the brown alga *Carpophyllum*: investigating morphology, distribution and wave exposure. MSc thesis, Victoria University of Wellington.
- Kucera, H. and Saunders, G. W. 2008. Assigning morphological variants of *Fucus* (Fucales, Phaeophyceae) in Canadian waters to recognized species using DNA barcoding. *Botany* **86**: 1065–79.
- Lane, C. E., Mayes, C., Druehl, L. D. and Saunders, G. W. 2006. A multi-gene molecular investigation of the kelp (Laminariales, Phaeophyceae) supports substantial taxonomic re-organization. *J. Phycol.* **42**: 493–512.
- Lihova, J., Kucera, J., Perny, M. and Marhold, K. 2007. Hybridization between two polyploid *Cardamine* (Brassicaceae) species in north-western Spain: Discordance between morphological and genetic variation patterns. *Ann. Botany.* **99**: 1083–96.
- Lindauer, V. W., Chapman, V. J. and Aiken, M. 1961. The marine algae of New Zealand: Phaeophyceae. *Nova Hedwigia* **3**: 129–350.
- Mathieson, A. C., Norton, T. A. and Neushul, M. 1981. The taxonomic implications of genetic and environmentally induced variations in seaweed morphology. *Bot. Rev.* **47**: 313–47.
- Mathieson, A. C., Dawes, C. J., Wallace, A. L. and Klein, A. S. 2006. Distribution, morphology, and genetic affinities of dwarf embedded *Fucus* populations from the Northwest Atlantic Ocean. *Bot. Mar.* **49**: 283–303.
- Montanari, A. and Mignani, S. 1994. Notes on the bias of dissimilarity indices: the case of archaeological classification. *Questiio* **18**: 39–49.
- Nelson, W. A. 1994. Distribution of macroalgae in New Zealand – an archipelago in space and time. *Bot. Mar.* **37**: 221–33.
- Perrin, C., De Vliet, M. V., Daguin, C., Engel, C., Pearson, G. A. and Serrão, E. A. 2005. Population genetic structure of dioecious versus hermaphrodite *Fucus* species, hybridisation along sympatric and allopatric zones. *Phycologia* **44**: 78–9.
- Podani, J. 1999. Extending Gower's general coefficient of similarity to ordinal characters. *Taxon* **48**: 331–40.
- Rieseberg, L. H. and Ellstrand, N. C. 1993. What can molecular and morphological markers tell us about plant hybridization? *Crit. Rev. Plant Sci.* **12**: 213–41.
- Rohlf, F. J. 2000. *NTSYS-pc: Numerical Taxonomy and Multivariate Analysis System*. Version 2.11. Exeter Software, New York.
- Rohlf, F. J. and Sokal, R. R. 1981. Comparing numerical taxonomic studies. *Syst. Zool.* **30**: 459–90.
- Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. *Molecular Cloning. A Laboratory Manual*, 2nd edn. Cold Springs Harbor Laboratory, Cold Springs Harbor, NY.
- Schiell, D. R. 1980. A demographic and experimental evaluation of plant and herbivore interactions in subtidal algal stands. PhD thesis, University of Auckland.

- Schiel, D. R. 1988. Algal interactions on shallow subtidal reefs in northern New Zealand – a review. *N. Z. J. Mar. Freshwat. Res.* **22**: 481–9.
- Schiel, D. R. 1990. Macroalgal assemblages in New Zealand – structure, interactions and demography. *Hydrobiologia* **192**: 59–76.
- Scott, G. W. and Hardy, F. G. 1994. Observations of the occurrence of hybrids between two sympatric species of Furoid algae. *Cryptogam. Algal.* **15**: 297–305.
- Seehausen, O. 2004. Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**: 198–207.
- Thuret, G. 1854. Recherches sur la fecondation des Fucaees. *Ann. Sci. Nat. Bot.* **4**: 197–214.

- Tiffin, P., Olsen, M. S. and Moyle, L. C. 2000. Asymmetrical crossing barriers in angiosperms. *Proc. R. Soc. Biol. Sci. Ser. B* **268**: 861–7.
- Wallace, A. L., Klein, A. S. and Mathieson, A. C. 2004. Determining the affinities of salt marsh fucoids using microsatellite markers: Evidence of hybridization and introgression between two species of *Fucus* (Phaeophyta) in a Maine estuary. *J. Phycol.* **40**: 1013–27.
- Zuccarello, G. C. and Lokhorst, G. M. 2005. Molecular phylogeny of the genus *Tribonema* (Xanthophyceae) using *rbcL* gene sequence data: monophyly of morphologically simple algal species. *Phycologia* **44**: 384–92.